20497

COPPER 4 SULFUR 60 SEVIN'7 BORON PEANUT DUST

NET WEIGHT 50 LBs.

FUNGICIDE-NUTRITIONAL-INSECTICIDE

DIRECTIONS FOR USE

CROP: PEANUTS

PESTS: Leafspot, aid in correcting the deficiency of boron, Mexican bean beetles, Thrips, Leaf-hoppers, Blister beetle, Bean leaf beetles.

Apply undiluted as a dust at the rate of 15 pounds per acre when spots first appear on old leaves at the base of the plants or when insects first appear. Only one application should be made per season. Dusting should be done with a machine capable of depositing a thin uniform coating of dust on all above ground plant surfaces. Dusting should only be done when the foliage is dry. Severe burning may result if application is made when the leaves are wet. Use of more than the recommended dosage may result in reduced yields.

Applications of dusting or spraying materials containing sulfur should not be made just prior to orduring periods of excessively high temperatures.

U. S. D. A. Reg. No. 279-2800	
ACTIVE INGREDIENTS:	
*Carbaryl (1-Naphthyl methyl-	
-carbamate)	7.0%
**Suifur	60.0%
***Copper expressed as metallic	4.0%
INERT INGREDIENTS: ****	29 .0 %
-	100 0%

*Sevin U.S.Pat. Nos. 2,903,478; 3,009,855

**Sulfur fineness: 93% through 325 mesh

***In copper oxide form

****Boron expressed as metallic ----3.34%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid prolonged or repeated contact with skin. Wash hands, and face before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Do not contaminate or store near feed or foodstuffs. NOTE TO PHYSICIAN: Carbaryl is a moderate,

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

USAGE CAUTIONS

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following dusting.

Do not use with lime, Bordeaux, lime sulfur, casein-lime spreaders or other alkaline materials.

Dispose of wastes by burying in non-crop lands away from water supplies. Dispose of bags by burying with wastes or by burning. Keep out of smoke.



LABEL

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGH-BORING CROPS OR NON-CROP AREAS OR USE IN A MAN-NER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UN-DESIRABLE RESULTS MAY OF CLEE

TERMS OF SALE OR USE. On purchase it ask of this product buyer and user agree to the full king. I tutons

WARRANTY Neagara warrant that these to to the associate to end to save by Nagara of the form of the large front statement on the labels of the first end to the form of the large front statement of the labels of the first end to the first end to

Transaction of the contract of the services of the contract of the damages of the property of the problem of the problem of the services of the services of the problem of

PROTOR FROM NON ONTROLLABOR CONFILIONS Be also to establish the state of established to add to the content of the brown of graphs of the brown and assert to a point of the brown and assert to a point of the brown as for a section of the brown and assert to a point of the protocology of the protoco

MANUFACH RED IN USA BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

