

CODE
3087

NET CONTENTS
GALS

ETHION 67 SUPERIOR 70 OIL

[279-2863]

INSECTICIDE - MITICIDE

DIRECTIONS FOR USE

Unless otherwise noted, begin applications when scale "crawlers" or mites first appear. Thorough coverage of foliage and branches is necessary for good control. If other spray materials are to be used with the oil in the same spray tank, these materials should be well-mixed in the water first, then the oil should be added last as the tank is being filled. Labels for such materials should be carefully examined to determine whether they are compatible with this material. Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milk-white spray mixture results, provided the spray mixture is being agitated. If the spray mixture, after adding the oil to the spray tank with the agitation running, is not milk-white, do not use the remainder of this product. Do not apply within 2 weeks of a sulfur spray or within 7 days of Captan.

APPLY LISTED AMOUNTS PER 100 GALLONS OF WATER

APPLES, PEARS: Brown Almond Mite, European Red Mite, Two-spotted Mite, Pear Psylla - Use 2 quarts per 100 gallons of water using a maximum of 750 gallons of spray per acre or use 3½ gallons of this product in sufficient water to treat one acre. Apply when pests first appear and repeat as necessary to maintain control. **Oystershell Scale:** Use 2 quarts per 100 gallons of water (using a maximum of 750 gallons of spray per acre) or 3 quarts per 100 gallons (using a maximum of 500 gallons of spray per acre) or use 2½ to 3½ gallons of this product in sufficient water to treat one acre. Apply to "crawlers" emerging from the overwintering eggs (about June 1) and repeat in 60 days if necessary to control second generation "crawlers."

DO NOT EXCEED 3 APPLICATIONS DURING THE FRUITING PERIOD AND DO NOT APPLY WITHIN 90 DAYS OF HARVEST

Do not use on varieties such as: Doona, Margaret Pratt, Melba, Wealthy, Williams, Bart (and Williams), Yellow Transparent or any varieties containing a high percentage of Intosh or on Grimes and N.E.S. (Red Mar) apples as they are highly susceptible.

CHERRY: European Red Mite, Two-spotted Mite, Pear Psylla - Use 2 quarts per 100 gallons of water using a maximum of 750 gallons of spray per acre or use 3½ gallons of this product in sufficient water to treat one acre. Apply when pests first appear and repeat as necessary to maintain control.

EVERGREENS, HARD TREES AND/OR ORNAMENTAL: Anthonomus, Citrus Red Mite, European Red Mite, Pear Psylla, Citrus Black Scale, Black Scale, Chaff Scale, Florida Red Scale, Glover Scale, Purple Scale, Snow Scale, Yellow Scale - Use 2 quarts per 100 gallons of water (using a maximum of 750 gallons of spray per acre) or 3 quarts per 100 gallons (using a maximum of 500 gallons of spray per acre) or use 2½ to 3½ gallons of this product in sufficient water to treat one acre. Apply to "crawlers" emerging from the overwintering eggs (about June 1) and repeat in 60 days if necessary to control second generation "crawlers" first appear. Do not apply within 90 days of harvest.

Do not use on varieties such as: Doona, Margaret Pratt, Melba, Wealthy, Williams, Bart (and Williams), Yellow Transparent or any varieties containing a high percentage of Intosh or on Grimes and N.E.S. (Red Mar) apples as they are highly susceptible.

EPA REG. NO. 105-105	
ACTIVE INGREDIENTS	
Ethion	70.00
Petroleum Oil	30.00
INERT INGREDIENTS	100.00

* 0.01% of total weight is S-methoxyethoxyethyl alcohol, U.S. Pat. No. 2,673,228. This product contains superior type oil. Contains 0.67 pounds ethion per gallon.

CAUTION
KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. May be absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near food or food products.

WARNING! FLAMMABLE

KEEP AWAY FROM HEAT AND OPEN FLAME

DO NOT STORE BELOW 10 F.

DO NOT USE IN UNDILUTED FORM

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Do not use on varieties such as: Doona, Margaret Pratt, Melba, Wealthy, Williams, Bart (and Williams), Yellow Transparent or any varieties containing a high percentage of Intosh or on Grimes and N.E.S. (Red Mar) apples as they are highly susceptible.

Do not use on varieties such as: Doona, Margaret Pratt, Melba, Wealthy, Williams, Bart (and Williams), Yellow Transparent or any varieties containing a high percentage of Intosh or on Grimes and N.E.S. (Red Mar) apples as they are highly susceptible.


Do not use on varieties such as: Doona, Margaret Pratt, Melba, Wealthy, Williams, Bart (and Williams), Yellow Transparent or any varieties containing a high percentage of Intosh or on Grimes and N.E.S. (Red Mar) apples as they are highly susceptible.

GRAPEFRUITS, LEMONS, LIMES, ORANGES, TANGZLOS, TANGERINES: Citrus Red Mite, Citrus Rust Mite, Texas Citrus Mite, Black Scale, Chaff Scale, Florida Red Scale, Glover Scale, Purple Scale, Snow Scale, Yellow Scale - Use 2 quarts per 100 gallons of water. Do not apply more than 2000 gallons of spray per acre. Apply when insects first appear and repeat as required. Where snow scale is a problem, it is important that trunks, limbs and twigs are thoroughly sprayed. Do not use on navel oranges.

On grapefruit, oranges and tangzlos do not apply within 90 days of preceding treatment. May be applied up to day of harvest.

On tangerines do not apply within 90 days of preceding treatment. Do not apply more than twice per season. May be applied up to day of harvest.

On lemons and limes do not apply more than once per season. Do not apply within 21 days of harvest.



Dealers Should Sell in Original Packages Only.

TERMS OF SALE OR USE - On the basis of the product label and user's agreement to the following conditions:

WARRANTY: FMC warrants that the product is of the type and quality as shown on the label and that it is reasonably fit for the purposes and uses indicated on the label. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: FMC is not liable for damages for breach of warranty or for consequential, special, or incidental or consequential damages. Notwithstanding to whom reported to FMC within the time limit, a user's liability for damages which he or she causes may exceed such damage.

EXCUSE FROM NON-CONTROL UNDER CONDITIONS: Because of certain natural conditions, the user may be unable to control or over-control the pest. FMC does not warrant or assume any liability arising from the use of this product under such conditions. FMC does not assume any liability for any damage or loss caused by such causes. No person shall be held liable in any way for a statement contained herein.

FMC