



Code 3087

Net Contents
Gallons

Ethion .67 Superior 70 Oil

Insecticide-Miticide

For Agricultural or Commercial Use Only.

EPA Reg. No. 279-2863 EPA Est., 279-

| | |
|--------------------------|--------|
| Active Ingredients: | By Wt. |
| *Ethion | 9.2% |
| Petroleum Oil | 77.0% |
| Inert Ingredients: | 13.8% |
| | <hr/> |
| | 100.0% |

*O,O,O',O'-tetraethyl S,S'-methylene bisphosphorodithioate
This product contains superior type oil. Contains 0.67 pounds ethion per gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

For Emergency Assistance Call 716-735-3765.
See Other Panels for Additional Precautionary Information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution
Harmful if swallowed. May be absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products.

Environmental Hazards

This product is toxic to fish, shrimp, crab and other wildlife. Keep out of lakes, ponds, streams, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from the areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Physical or Chemical Hazards

Flammable. Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not store below 10°F. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not reuse empty container. Do not cut or weld this container. To dispose of container, triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures.

Do not use in undiluted form.

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PA 66

ACCEPTED

JUL 31 1980

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 279-2863



EPA REG. NO. 101-101

Unless otherwise noted, begin applications when scale, crawlers, or mites first appear. Thorough coverage of foliage and branches is necessary for good control. If other spray materials are to be used with the oil in the same spray tank, these materials should be well mixed in the water first, then the oil should be added last as the tank is being filled. Labels for such materials should be carefully examined to determine whether they are compatible with this material. Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milk-white spray mixture results, provided the spray mixture is being agitated. If the spray mixture, after adding the oil to the spray tank with the agitator running, is not milk-white, do not use the remainder of this product. Do not apply within 2 weeks of a sulfur spray or within 7 days of Captan.

Apply listed amounts per 100 gallons of water.

Apples, Pears: Brown Almond Mite, European Red Mite, Twospotted Spider Mite, Pear Psylla—Use 2 quarts per 100 gallons of water using a maximum of 750 gallons of spray per acre or use 3½ gallons of this product in sufficient water to treat one acre. Apply when pests first appear and repeat as necessary to maintain control. Oystershell Scale—Use 2 quarts per 100 gallons of water (using a maximum of 750 gallons of spray per acre) or 3 quarts per 100 gallons (using a maximum of 500 gallons of spray per acre) or use 2¼ to 3½ gallons of this product in sufficient water to treat one acre. Apply to "crawlers" emerging from the overwintering eggs (about June 1) and repeat in 60 days if necessary to control second generation "crawlers." Do not exceed 3 applications during the fruiting period and do not apply within 30 days of harvest. Do not use on Astrachen, Davey, Duchess, Margaret Pratt, Melba, Wealthy, Williams Red (Red Williams), Yellow Transparent or any varieties maturing at or before early McIntosh or on Crimson and N.J. 63 (Brite Mac) apples as injury is likely to occur.

Blueberries: Blueberry Bud Mite—Use 1 to 2 gallons per 100 gallons of water. Apply 1 to 2 weeks after harvest and repeat 3 to 4 weeks later. Spray plants until thoroughly wet.

Evergreens, Shade Trees, Woody Ornamentals including Azaleas, Camellias, Gardenias, Ligustrum, Pittosporum, Sasangua, Viburnum, Boxwood, Cottoncaster, Dogwood, Lilac, Willow, Spruce, Pines, Juniper, Taxus (Yew), Camellia Scale, Euonymus Scale, Florida Red Scale, Florida Wax Scale, Oystershell Scale, Pine Needle Scale, Spruce Bud Scale, Tea Scale, Spider Mite, Taxus Mealy Bug—Use 2 to 3 quarts. Lecanium Scale—Use 3 quarts. For scale insect control, apply when the overwintering eggs start to hatch. If necessary for the control of the summer generation, apply when the "crawlers" first appear. For spider mite and taxus mealy bug control, apply when insects first appear and repeat as required to maintain control. Caution: Do not apply to American red bud trees as injury may occur. May cause slight needle drop on some varieties of Taxus.

Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines: Citrus Red Mite, Citrus Rust Mite, Texas Citrus Mite, Black Scale, Chaff Scale, Florida Red Scale, Glover Scale, Purple Scale, Snow Scale, Yellow Scale—Use 2 quarts per 100 gallons of water. Do not apply more than 2000 gallons of spray per acre. Apply when insects first appear and repeat as required. Where snow scale is a problem, it is important that trunks, limbs and twigs are thoroughly sprayed. Do not use on navel oranges.

On grapefruit, oranges and tangelos do not apply within 90 days of preceding treatment. May be applied up to day of harvest.

On tangerines, do not apply within 90 days of preceding treatment. Do not apply more than twice per season. May be applied up to day of harvest.

On lemons and limes, do not apply more than once per season. Do not apply within 21 days of harvest.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions.

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

(3087-PA 6/80)