

Code: 30375

10 Pounds Net Weight

[279-2872]

# KOLODUST<sup>®</sup> ETHION 50-4 DUST

## DIRECTIONS FOR USE

Apply when the pest or insect appears and repeat as necessary. For dust control, apply at the time of late stages of this product use a suitable fumigator. For coverage of most upper and lower leaf surfaces, use special hood control. Wash the applicator with detergent or similar application equipment. Observe days interval between last application and harvest indicated by number in following the crop.

**ALMONDS:** Brown Rot, Brown Almond Mite, European Red Mite, Two-Spotted Mite, Two-Spotted Mite. Use 40-60 pounds per acre. At a rate of 50 pounds per acre do not apply more than one application during the fruiting period. Do not apply within 30 days of harvest if used at rate of 51-60 pounds per acre do not apply after husk split.

**APPLES:** Powdery Mildew, Brown Almond Mite, European Red Mite, Apple Rust Mite, Two-Spotted Mite. Use 40-60 pounds per acre. Make no more than 4 applications during the fruiting period nor apply within 40 days of harvest or make no more than 3 applications during the fruiting period nor apply within 30 days of harvest or make no more than 2 applications during the fruiting period nor apply within 20 days of harvest. Sulfur may cause certain varieties to crack prior to harvest.

**PLUMS, PRUNES:** Brown Rot, Brown Almond Mite, European Red Mite, Two-Spotted Mite. Use 40-60 pounds per acre. Apply when first appear and repeat as required. At a rate not exceeding 43 pounds per acre do not make more than 3 applications during the fruiting period nor apply within 21 days of harvest. At a rate of 44-60 pounds per acre do not make more than 2 applications during the fruiting period nor apply within 40 days of harvest.

**CITRUS (Oranges (0), Grapefruit (0), Tangerines (0), Lemons (21), Limes (21), Tangelos (0),** California and Arizona Only. Citrus Thrips. Use 35-100 pounds per acre. Citrus Thrips, Citrus Red Mite, Texas Citrus Mite, Citrus Rust Mite. Use 70-100 pounds per acre. On Grapefruit, Oranges, Tangelos and Tangerines, if a single application is required do not apply within 90 days of the previous treatment. On Tangerines do not apply more than 2 applications per year. On Limes and Lemons do not apply more than one per season.

**STRAWBERRIES (2):** Two Spotted Spider Mite, Powdery Mildew. Use 30 pounds per acre. Treat only just both leaf surfaces and grow into the crown of the plants.

**PEACHES (30), NECTARINES (30):** Brown Rot, Powdery Mildew, European Red Mite, Two Spotted Mite, Brown Almond Mite. Use 40-60 pounds per acre. Do not apply more than 2 applications during the fruiting period.

**GRAPES:** Grape Leafhopper, Two Spotted Mite, Willow Mite, Powdery Mildew. Use 20-30 pounds per acre. Make no more than 2 applications nor apply within 30 days of harvest. Do not make more than 3 applications nor apply within 30 days of harvest. Do not apply Sulfur Mite at any time. Spray varieties as indicated on the label.

## FUNGICIDE-INSECTICIDE-MITICIDE

EPA REGISTRATION NO. 279-2872-AA

ACTIVE INGREDIENTS	BY WT.
Ethion	4.00%
Sulfur	42.00%
INERT INGREDIENTS	54.00%
	100.00%


000-0 Temporary SS Dimethyl bisphosphorothioate U.S. Patent No. 2,873,228.

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed storage, animal feed and foodstuffs.

This product contains bentonite. Bentonite Sulfur is made by the absorption of fused sulfur into Bentonite in the proportions of thirty parts sulfur to twenty parts Bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, response to spray tank and safety to host plants.



**BEANS:** Bean Leaf Beetle, Two Spotted Spider Mite, Thrips. Use 20 pounds per acre. Cover the lower leaf surfaces. Do not apply within 2 days of harvest. Do not feed treated beans to livestock. Eggus Bugs, Bean Leaf Beetle, Rust Thrips, Two-Spotted Mite, Powdery Mildew. Use 30-40 pounds per acre. Do not apply within 4 days of harvest. Do not feed treated beans to livestock. Sulfur may injure some varieties. Check sulfur tolerance prior to use.

**TOMATOES (2):** Tropical Mite, Two Spotted Mite, Russet Mite, Leafminer. Use 20 pounds per acre.

**PEPPERS (21):** Serpentine Leafminer, Two Spotted Mite. Use 25 pounds per acre. Do not make more than 3 applications during the fruiting period.

## USAGE CAUTIONS

This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Applications of materials containing sulfur should not be made just prior to or during periods of excessive heat, temperatures or within 10 days to 2 weeks of a hard spray.

Do not allow this material to drift into sulfur sensitive crops.

Do not apply when weather conditions favor drift from the areas treated. Do not apply when drift is likely to occur.

DISPOSE OF WASTES by burying in non-crop land away from water sources. Empty tanks should be cleaned with water or burned. Keep out of smoke.

For Emergency Assistance Call  
716-735-3765

