

Code: 30375

50 Po

[279-2872]

KOLODUST ETHION 50-4 DUST

DIRECTIONS FOR USE

Apply when the disease, mites or insects appear and repeat as necessary. For disease control beyond the immediate vicinity of this product use a suitable fungicide. Full coverage of both upper and lower leaf surfaces are essential to good control. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in column to lower the crop.

ALMONDS: Brown Rot, Brown Almond Mite, European Red Mite, Two Spotted Mite. Use 40-60 pounds per acre. At a rate of 50 pounds per acre or less do not apply more than one application during the fruiting period. Do not apply within 30 days of harvest. If used at a rate of 51-60 pounds per acre, do not apply after creek split.

APPLES: Powdery Mildew, Brown Blight, Mite, European Red Mite, Apple Rust Mite, Two Spotted Mite. Use 40-60 pounds per acre. Make no more than 4 applications during the fruiting period nor apply within 40 days of harvest or make no more than 3 applications during the fruiting period nor apply within 30 days of harvest or make no more than 2 applications during the fruiting period nor apply within 20 days of harvest. Sulfur may injure certain varieties. Check tolerance prior to use.

PLUMS, PRUNES: Brown Rot, Brown Almond Mite, European Red Mite, Two Spotted Mite. Use 40-60 pounds per acre. Apply when mites first appear and repeat as required. At a rate not exceeding 43 pounds per acre, do not make more than 3 applications during the fruiting period nor apply within 21 days of harvest. At a rate of 44-60 pounds per acre do not make more than 2 applications during the fruiting period nor apply within 45 days of harvest.

CITRUS (Oranges (0), Grapefruit (0), Tangerines (0), Lemons (21), Limes (21), Tangelos (0), (California and Arizona Only) Citrus Thrips. Use 35-100 pounds per acre. Citrus Thrips, Citrus Red Mite, Texas Citrus Mite, Citrus Rust Mite. Use 50-100 pounds per acre. On Grapefruit, Oranges, Tangelos and Tangerines, if additional applications are required, do not apply within 90 days of the preceding treatment. On Tangerines do not apply more than 2 applications per crop. On Lemons and Limes do not apply more than once per season.

STRAWBERRIES (2) Two Spotted Spider Mite, Powdery Mildew. Use 30 pounds per acre. Thoroughly dust both leaf surfaces and down into the crown of the plants.

PEACHES (30), NECTARINES (30) Brown Rot, Powdery Mildew, European Red Mite, Two Spotted Mite, Brown Almond Mite. Use 40-60 pounds per acre. Do not apply more than 2 applications during the fruiting period.

GRAPES: Grape Leafhopper, Two Spotted Mite, Williamsite Mite, Powdery Mildew. Use 20-30 pounds per acre. Make no more than 2 applications nor apply within 15 days of harvest or make no more than 3 applications nor apply within 30 days of harvest. Do not use on Salvador, Muscat and Fehr-Szajzo varieties as injury may occur. Ground application preferred.

FUNGICIDE-INSECTICIDE-MITICIDE

EPA REG. STRATH. N. N. 279-2872 AA

ACTIVE INGREDIENTS

Ethion
Sulfur

BY WT

4.00
42.00
54.00
100.00

INERT INGREDIENTS


0.00 0.00 Inert ingredients, phosphorus
EPA REG. STRATH. N. N. 279-2872 AA

CAUTION

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of or storage near feed and foodstuffs.

This product contains Bentonite Sulfur. Bentonite Sulfur is made by the absorption of sulfur by bentonite. Bentonite in the proportions of thirty parts sulfur to seventy parts Bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the nature of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, susceptibility to spray tank and safety to host plants.



BEANS: Bean Leaf Beetle, Two Spotted Spider. Cover the lower leaf surfaces. Do not apply treated foliage to livestock. Lygus Bugs, Bean Mite, Powdery Mildew. Use 30-40 pounds of product. Do not feed treated foliage to livestock for tolerance prior to use.

TOMATOES (2) Tropical Mite, Two Spotted Mite. Use 30-40 pounds per acre.

PEPPERS (21) Spiny Leaf Miner, Two Spotted Mite. Do not make more than 3 applications during the fruiting period.

USAGE CAUTIONS

This product is poisonous to fish and wildlife. Do not contaminate water by cleaning of equipment or disposal of this product only as specified on the label.

This product is toxic to bees and should not be used near flowering plants.

Applications of materials containing sulfur should be avoided during periods of excessively high temperatures or winds.

Do not allow this material to drift into sulfur-sensitive crops.

Do not apply when weather conditions favor runoff when runoff is likely to occur.

DISPOSE OF WASTES by burying in non-crop areas should be buried with wastes or burned.

For Emergency Assistance
716-735-27

50 Pounds Net Weight

KOLODUST[®] ETHION 50-4 DUST

USES FOR USE

... appear and repeat as necessary. For disease... product use a suitable fungicide. Full... faces are essential to good control. May... cation equipment. Observe days interval... ted by number in following the crop.

... European Red Mite, Two Spotted Mite... pounds per acre or less. Do not apply more... Do not apply within 30 days of harvest... do not apply after husk split.

... Mite, European Red Mite, Apple Rust Mite... acre. Make no more than 4 applications... 40 days of harvest, or make no more than... not apply within 30 days of harvest, or... the fruiting period nor apply within 20 days... check tolerance prior to use.

... Mite, European Red Mite, Two Spotted Mite... mites first appear and repeat as required... re do not make more than 3 applications... in 21 days of harvest. At a rate of 34.60... 2 applications during the fruiting period nor...

... Citrus (0), Lemons (21), Limes (21), Tangelos (0)... Use 35-100 pounds per acre. Citrus Thrips... Rust Mite. Use 50-100 pounds per acre. On... es if additional applications are required. Oc... ing treatment. On Tangerines do not apply... ons and Limes do not apply more than once.

... Mite, Powdery Mildew. Use 30 pounds per... nd down into the crown of the plants.

... Rot, Powdery Mildew, European Red Mite... Use 40-60 pounds per acre. Do not apply... ing period.

... Mite, Willamette Mite, Powdery Mildew... re than 2 applications nor apply within 15... s applications nor apply within 30 days of... nd Fehr Szajo varieties as injury may occur.

FUNGICIDE-INSECTICIDE-MITICIDE

EPA REGISTRATION NO. 27-2872-AA

ACTIVE INGREDIENTS

Ethion

Sulfur

INERT INGREDIENTS

000 0 tetraethyl ammonium disphosphoric
thionate U.S. Patent No. 2,873,228

BY WT

4.00

15.00


100.00

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. May be absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of or storage near feed and foodstuffs.

This product contains bentonite Sulfur. Bentonite Sulfur is made by the absorption of fused sulfur into Bentonite in the proportion of thirty parts sulfur to seventy parts Bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, insects and plants susceptible to spray tank and safety, to host plants.



BEANS (Do not use on Two Spotted Mite, Thrips). Use 20 pounds per acre. Control the lower leaf nodes. Do not apply within 2 days of harvest. Do not feed treated beans to livestock. Lygus Bugs, Bean Leaf Beetle, Rust, Thrips, Two Spotted Mite, Powdery Mildew. Use 30-40 pounds per acre. Do not apply within 4 days of harvest. Do not feed treated foliage to livestock. Sulfur may injure some varieties, check tolerance prior to use.

TOMATOES (2) Tropic Mite, Two Spotted Mite, Russet Mite, Leafminer. Use 20 pounds per acre.

PEPPERS (21) Spider Mite, Leafminer, Two Spotted Mite. Use 25 pounds per acre. Do not make more than 3 applications during the fruiting period.

USAGE CAUTIONS

This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Applications of materials containing sulfur should not be made just prior to or during periods of excessively high temperatures or within 10 days to 2 weeks of an oil spray.

Do not allow this material to drift into sulfur sensitive crops.

Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty cans should be buried with waste or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765