

Code 3214

RESTRICTED USE PESTICIDE

Net Weight

Due To Avian Toxicity

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

Furadan® 5 G

Insecticide-Nematicide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-2874

EPA Est. 279-

Active Ingredient:

*Carbofuran 5.0%
 Inert Ingredients: 95.0%
 100.0%

*2,3-Dihydro-2,2-dimethyl-7-benzofuranylmethylcarbamate

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in eyes: Flush with plenty of water for at least 15 minutes. Get medical attention.

If on skin: Wash immediately with soap and water. Remove contaminated clothing and wash before reuse.

Antidote

Note to Physician: Carbofuran is a reversible cholinesterase inhibitor. Do not use oximes such as 2-PAM. Start by giving 2 mg. atropine intramuscularly. According to clinical response, continue until signs of atropinization occur (dry mouth or dilated pupils). If in eye, instill one drop of homatropine.

For Emergency Assistance Call (800)-331-3148.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

Poisonous if swallowed. May be fatal or harmful as a result of skin or eye contact or by breathing dust. Causes cholinesterase inhibition. Warning symptoms of poisoning include weakness headache, sweating, nausea vomiting, diarrhea, tightness in chest, blurred vision, pinpoint eye pupils, abnormal flow of saliva, abdominal cramps, and unconsciousness. Atropine sulfate is antidotal.

Wear long-sleeved clothing and protective gloves when handling. Wash hands and face before eating or smoking. Bathe at the end of the work day. Change clothing daily and wash before reuse.

See other panels for additional precautionary information.

Environmental Hazards

This pesticide is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Birds killed by carbofuran pose a hazard to hawks and other birds-of-prey; contact your state wildlife agency or the U.S. Fish and Wildlife Service upon finding birds or other animals that may have been poisoned. Cover or incorporate granules in spill areas. Runoff from treated areas may be hazardous to fish in neighboring areas. Do not apply directly to water. Do not contaminate wells, wetlands or any body of water by cleaning of equipment or disposal of waste.

Carbofuran is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Carbofuran has been found in ground water as a result of agricultural use. Users are advised not to apply carbofuran where the water table (ground water) is close to the surface and where the soils are very permeable, i.e., well-drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not re-enter treated fields less than 24 hours following application.

Do not use this product on Long Island, N.Y.

Do not use this product on rice after 8/31/95.

STORAGE AND DISPOSAL**Pesticide Storage**

Do not store in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.



Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



FMC Corporation
 Agricultural Chemical Group
 Philadelphia PA 19103

3/93



Do not plant with any crop other than alfalfa, artichokes, bananas, barley, coffee, corn (field, pop or sweet), cotton, cranberries, cucurbits (cucumbers, melons, pumpkins, squash), flax, grapes, non-bearing fruit, oats, ornamentals, peanuts, peppers, potatoes, rice, seed crops (Bermuda-grass, spinach), sorghum, strawberries, soybeans, sugar beets, sugarcane, sunflowers, tobacco, or wheat for at least 10 months following use of this product.

Rice: To control Rice Water Weevil and Dark Rice Field Mosquito in Arkansas, Louisiana, Mississippi and Texas apply 10 pounds of Furadan 5 G insecticide per acre one day before or within 21 days after flooding. Do not make more than one application per season. The subsequent use of propanil herbicide may result in crop injury.

For effective control of the dark rice field mosquito, application must be made within 2 to 4 days after flooding.

To control Rice Water Weevil in California apply 10 pounds of Furadan 5 G per acre prior to flooding and planting. To reduce risk to waterfowl all preplant applications of Furadan 5 G must be incorporated prior to initial flood. Method of incorporation must insure that Furadan 5 G is mixed no deeper than 2 inches for effective weevil control. Do not make more than one application per season. The subsequent use of propanil herbicide may result in crop injury.

Note: FMC will sell this product in conformity to volume limitations agreed to with EPA.

FMC will not sell, nor release for shipment, this product for use on rice after 8/31/94

Product in channels of trade may be sold and used until 8/31/95.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Furadan and  —FMC Trademarks

(3669-10/2/92-A)

