


Code 2837

1 Gallon Net Contents

Furadan® 4 Flowable



Insecticide

EPA Reg No 279-2876ZA

EPA Est., 279-NY-1

Active Ingredient:

*Carbofuran 40.64%

Inert Ingredients: 59.36%

100.00%

*2,3-dihydro-2, 2-dimethyl-7-benzofuranyl methylcarbamate
This product contains 4 lbs. of carbofuran per gallon.

Furadan -- FMC Trademark

U.S. Patent No. 3,474,171

KEEP OUT OF REACH OF CHILDREN



For Emergency Assistance Call 716-735-3765.

First Aid

If **swallowed**—Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If **inhaled**, remove to fresh air. Call a physician immediately.

If in **eyes**, flush with plenty of water for at least 15 minutes. Get medical attention.

If on **skin**, in case of contact, remove contaminated clothing and immediately wash skin with soap and water.

Antidote

To **Physician** Furadan is a reversible cholinesterase inhibitor. Do not use oximes such as 2-PAM. Give atropine 2 mg. intravenously or subcutaneous. If in eye instill one drop of Homatropine.

See side panel for additional precautionary statements.

ACCEPTED
SEP 18 1978
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under the pesticide Act. EPA Reg. No. 279-2876

FMC
FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia PA 19103

PA-2 6/78

Precautionary Statements

Hazards To Humans (& Domestic Animals)

Danger

Poisonous by swallowing or inhalation. Do not breathe spray mist. Do not get in eyes. Avoid contact with skin. Wear a pesticide respirator jointly approved by the Mining Enforcement and Safety Administration (formerly the U.S. Bureau of Mines) and by the National Institute for Occupational Safety and Health under the provisions of 30 CFR Part 11.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not reuse the container. Completely empty the contents and bury the unused chemical at least 18 inches deep in an isolated location away from water supplies. Bury the container in the same manner.

Not for use or storage in or around the house.

Do not plant crops other than alfalfa, corn, peanuts, peppers, potatoes, rice, sorghum, strawberries, sugar beets, sugarcane and tobacco within 18 months of last application. Soybeans and oats may be planted the following season. Any other crop may be planted if it is not harvested or grazed.

Alfalfa Weevil Larvae, Egyptian Alfalfa Weevil Larvae, Pea Aphid, and in N.Y. State the Snout Beetle control — Apply the amount of Furadan 4 Flowable, indicated in the chart, when feeding is noticed or when insects appear. **Alfalfa Weevil Adult** — Apply 1 to 2 pints per acre when insects appear. **Alfalfa Blotch Leafminer (Northeastern States only)** and **Potato Leafhopper** — Apply 2 pints per acre when insects appear. **Lygus Bugs** — Apply 2 pints per acre prior to bloom. **Grasshoppers** — Apply ¼ to ½ pint per acre when grasshopper feeding is noticed. For control of **Blue Alfalfa Aphid (nymphs and wingless adults)** in California and Arizona — Apply Furadan 4 Flowable at ½ to 1 pint per acre when insect feeding is noticed or when insects appear. Observe the indicated number of days after application before cutting or grazing. Do not apply more than once per season. Apply only to fields planted to pure stands of alfalfa. Do not move bees to Alfalfa fields within 7 days of application.

Pints of
Furadan 4 Flowable
Per Acre

¼
1
2

Do not cut or
graze within:

7 days
14 days
28 days

Minimum gallonage requirements: Ten gallons of finished spray per acre with ground equipment, two gallons per acre with aircraft, except Blue Alfalfa Aphid use 30 gallons of finished spray per acre with ground equipment, 5 gallons per acre with aircraft.

For water fowl protection do not apply on fields in proximity of water fowl nesting areas and/or on fields where water fowl are known to repeatedly feed.

Field Corn Corn Rootworms — Use 1½ to 2 pints of Furadan 4 Flowable per 13,000 linear feet (1 acre with 40 inch spacing). Apply, at planting, as a 7 inch band over the row or inject on each side of the row by mixing with water or liquid fertilizers. When Furadan 4 Flowable is used with liquid fertilizers, mix in the following way making certain that the mixture is physically compatible: Premix 1 part of Furadan 4 Flowable with 2 parts of water. Add this premix to the tank of fertilizer along with rinsings from the premixing container. Maintain agitation in the tank after mixing and during application. Do not mix until ready to use.

Pine Seedlings Pines Weevil and Pitch-eating Weevil in pine plantations — Apply a 1% (W/W) active Furadan clay slurry (see following for preparation) to the roots of pine seedlings prior to transplanting. Treat seedlings by dipping roots or use any other suitable means which allows thorough coating. Keep roots moist until transplanted. Prepare the slurry as follows: Add 1.8 ounces (2½ tablespoons) of Furadan 4 Flowable to ¼ gallon of water. Mix thoroughly. Add 2 pounds of pulverized kaolin clay (pH 4.5) to this suspension. Mix thoroughly. This is sufficient to treat the roots of 150 to 200 seedlings. Adequate ventilation is required for indoor treatment.

Potatoes (For use in the Northeast & Northcentral States including Colorado) Colorado Potato Beetle, Potato Flea Beetle, Potato Leafhopper — Use 1 to 2 pints of Furadan 4 Flowable in sufficient water to treat one acre. Apply when insects first appear and repeat, as necessary to maintain control. Do not make more than 8 foliar applications per season. Do not apply more than 3 quarts to the foliage if Furadan 10 Granules were used at planting. Do not apply more than 1 quart per application. Do not apply within 14 days of harvest. Apply by ground equipment only.

For water fowl protection, do not apply on furrow irrigated fields and/or on fields where water fowl are known to repeatedly feed.

Strawberry (For use in Washington & Oregon) Root Weevils — Use Furadan 4 Flowable at 2 to 4 pints per acre (2.6 to 5.1 fluid ounces per 1000 linear feet of row for 42 inch row spacing). Apply as a 10 to 12 inch band over the row after last harvest but before October 1st. Do not make more than one application per season. Do not apply if berries are present.

Sugarcane Sugarcane Borer — Apply 1 to 1½ pints Furadan 4 Flowable per acre using ground or aerial equipment. Check sugarcane fields weekly, beginning in early June and continuing through August. Make first application only after visible joints form and 5% or more of the plants are infested with young larvae feeding in or under the leaf sheath and which have not bored into the stalks. Repeat whenever field checks indicate the infestation exceeds 5 percent. Do not apply within 17 days of harvest. Do not use in Hawaii.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

279-2876

9-18-78

10A1