

Code: 30878

Net Weight

50 Pounds

SULPHUR 25 KELTHANE* 4 THIODAN[®] 4 DUST

DIRECTIONS FOR USE

Apply when pests or disease first appear and repeat as necessary to maintain control. For control beyond the use limitations of this product use a suitable material. May be applied by air or ground. Observe days interval between last application and harvest indicated by number in () following crop.

APPLES, PEARS (7): Clover Mite, European Red Mite, European Red Mite, Twospotted Spider Mite and Apple Aphid — Use 50 to 60 pounds per acre. On Pears do not apply more than twice during fruiting season. On Apples do not apply within 21 days of harvest or more than 3 times during the fruiting season; or do not apply within 30 days of harvest with no limit on the number of applications. Do not feed pomace from treated apples to livestock. Sulphur may injure certain varieties, check tolerance prior to use.

GRAPES (7): Powdery Mildew, Grape Leafhopper, Grape Leafroller (Calif. Only), Twospotted Spider Mite, Pacific and Willamette Mites — Use 30 to 35 pounds per acre. For Leafroller control repeat applications may be necessary. Do not use on Concord variety of grapes. Ground application preferred.

PEACHES (30), NECTARINES (30): Green Peach Aphid, Peach Twig Borer (Calif. Only), Peach Silver Mite (Calif. Only) and Twospotted Spider Mite — Use 50 to 60 pounds per acre. Do not make more than 2 applications during the fruiting season.

PLUMS (7), PRUNES (7): Peach Twig Borer (Calif. Only) and Twospotted Spider Mite — Use 50 pounds per acre. Do not reapply within 30 days.

FILBERTS (14): Filbert Aphid, Filbert Leafroller, Clover Mite, Twospotted Spider Mite and European Red Mite — Use 50 pounds per acre. Do not graze livestock in treated groves.

WALNUTS: Walnut Aphid, Clover Mite, European Red Mite and Twospotted Spider Mite — Use 50 pounds per acre. Do not apply after husks split. Do not graze livestock in treated groves.

TOMATOES (2): Aphid, Flea Beetle, Hornworm, Colorado Potato Beetle, Tomato Psyllid, Whitefly, Stinkbug, Tomato Fruitworm, Cabbage Looper, Yellowstriped Armyworm, Tomato Russet Mite, Pacific and Tropical Spider Mites — Use 25 pounds per acre.

ORNAMENTALS (Greenhouse or Out-of-doors): Aphids, Whitefly and Mite. Dust for complete coverage lower and upper leaf surfaces. Do not use on Birch trees. In greenhouse applications wear mask or respirator passed by U.S. Bureau of Mines for Endosulfan (Thiodan) protection.

DRIED BEANS (7): Black Bean Aphid, Bean Leafskeltonizer, Mexican Bean Beetle, Twospotted Spider and Pacific Spider Mites — Use 25 pounds per acre. Do not apply more than 3 times to bean pods. Do not feed treated vines to livestock or allow to graze in treated fields. Sulphur may injure certain varieties, check tolerance prior to use. Do not use on Lima Beans that are to be removed from the field for processing.

FUNGICIDE-INSECTICIDE-MITICIDE

F.P.A. REG NO. 279-2885 AA

ACTIVE INGREDIENTS:	BY WT.
Endosulfan (Hexachloroexahydromethano-2,4,3-benzodioxathiepin oxide)	4.00%
1,1-Bis (p-chlorophenyl)-2,2,2-trichloroethanol	4.00%
Sulphur	25.00%
INERT INGREDIENTS:	67.00%
	100.00%

Thiodan is a registered trademark of Canadian Hoechst Ltd.
U.S. Patent No. 2,799,685.
*Trademark Rohm & Haas Company.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of or storage near feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

NOTE TO PHYSICIANS: Thiodan is a central nervous system stimulant. Treat symptomatically. There is no specific antidote.

USAGE CAUTIONS

Workers entering treated fields within 24 hours of treatment should wear protective clothing.

Do not plant root crops other than carrots, potatoes, sweet potatoes, or sugar beets as follow-up crops.

Do not allow this material to drift onto sulphur sensitive crops.

Applications of material containing sulphur should not be made prior to or during periods of excessively high temperatures.

For Emergency Assistance Call: 716—735-3765

SUCCULENT BEANS (7): Black Bean Aphid, Bean Leafskeltonizer, Cucumber Beetles, Green Stink Bug, Mexican Bean Beetle, Twospotted Spider and Pacific Spider Mites — Use 25 pounds per acre. Do not apply more than 3 applications per season. Do not use on Lima Beans that are to be removed from the field for processing. Do not feed treated threshing to livestock or allow to graze in treated fields. Sulphur may injure certain varieties, check tolerance prior to use.

COTTON: Cotton Stink Bug, Lygus Bugs, Bollworm, Cabbage Looper, Cotton Leafperforator, Twospotted Spider Mite, Pacific Spider, Atlantic and Desert Mite — Use 35 pounds per acre. Do not apply after bolls open. Do not graze treated fields by meat or dairy animals.

PEPPERS (2): Twospotted Spider, Pacific Spider and Tropical Spider Mites, Russet Mite, Flea Beetle, Green Peach Aphid, Pepper Maggot and Hornworm — Use 25 pounds per acre.

STRAWBERRIES (4): Powdery Mildew, Meadow Spittlebug, Strawberry Aphid, and Twospotted Spider Mite — Use 30 pounds per acre. Do not reapply within 35 days when fruit is present.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep Out Of Smoke.

ACCEPTED

AUG 22 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED



277-2887

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured by FMC
277-2887
Approved for sale by MDA
MCA 105

FMC
Chemicals