

CODE

991

NET WEIGHT

50 LBS.

TOXAKIL 20 DUST

INSECTICIDE

DIRECTIONS FOR USE

Begin applications when insects first appear and repeat as required to maintain control.

Best results will be obtained if applications are made when the air is calm. Applications which are washed off by rain within 24 hours should be repeated.

BEANS: Thrips, bean beetle, cowpea curculio, climbing cutworms, armyworms - Use at the rate of 15 to 20 pounds per acre. Do not apply on green or snap beans within 7 days of harvest. Do not feed treated vines to dairy animals or animals being finished for slaughter.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER: Imported cabbage worm, cross-striped cabbage worm, diamond-back caterpillar - Use at the rate of 15 to 20 pounds per acre. Do not apply after edible parts begin to form or within 30 days of harvest.

CABBAGE: Imported cabbage worm, cross-striped cabbage worm, diamond-back caterpillar - Use at the rate of 15 to 20 pounds per acre. Do not apply after heads start to form.

LETTUCE: Cucumber beetles, climbing cutworms, flea chinch bug - Use at the rate of 15 to 20 pounds per acre. Do not apply on leaf lettuce after seedling stage or on head lettuce after heads start to form.

E. P. A. Reg. No. 279-2888

ACTIVE INGREDIENTS:

*Toxaphene -----	20.0%
INERT INGREDIENTS: -----	80.0%
	<u>100.0%</u>

*Technical chlorinated camphene, 67-69% chlorine.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Avoid storage near feed and food products.

Toxaphene is a central nervous system stimulant. There is no specific antidote. Barbituric acid derivatives may be used for convulsions. Treat symptomatically.

For Emergency Assistance Call
716-735-3765

ACCEPTED

PEPPERS: Climbing cutworms, Flea beetle, pepper weevils, cucumber beetles - Use at the rate of 10 pounds per acre. Do not apply within 5 days of harvest.

TOMATOES: Colorado potato beetle, flea beetle, leafhoppers, thrips - Use at the rate of 10 pounds per acre. Do not apply within 1 day of harvest.

COTTON: Boll weevil, bollworm, leafworm, lygus bugs, fleahoppers, grasshoppers, armyworms - Use at the rate of 15 to 20 pounds per acre. If applied late in season, do not graze or feed to dairy animals being finished for slaughter.

USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding in treated areas may be killed. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are active in visiting the area.

Do not apply under weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur.

Dispose of waste in a manner non-crop lands away from water supplies. Bags should be disposed of in a safe manner with wastes or by burning in a safe manner.

Printed in U. S. A.