

Code 388

50 Lbs. Net Weight

Chlor Kil 10 Dust

Insecticide

EPA Reg. No. 270-204 AA EPA Est., 2704 A-2

Active ingredient	
Technical Chlor丹	10.00%
Inert ingredients	90.00%

*Equivalent to 0.04% (chlorinated, p-dithienyl)-hydroquinone and 4.1% of related compounds.

CAUTION
KEEP OUT OF REACH OF CHILDREN.

Avoid breathing dust. Avoid contamination of food or feedstuffs. Avoid contact with skin and eyes.


For Emergency Assistance Call First National

Environmental and Disposal Solutions

This product is toxic to fish and birds. Keep out of reach of children and animals. Avoid breathing dust. Avoid contact with skin and eyes. Avoid contamination of food or feedstuffs. Avoid contact with skin and eyes. Avoid contamination of food or feedstuffs.



1000 West 10th Street
 Fargo, ND 58103



Directions for Use

Beans, Beets, Broccoli, Cabbage, Crucifer Sprouts, Cauliflower, Peppers, Potatoes, Rucifera, White Yams (Michigan only). Broadcast 10 pounds of dust per acre as a pre-plant soil application or at time of planting or transplanting. Mix the dust into the top 4 to 6 inches of soil.

Ornamental Sod, Turf, and Nursery Stock. Japanese Beetle Larvae (Federal and State Quarantine Programs). Use 20 pounds per acre (10 ounces per 1000 sq. ft.) on established turf areas immediately upon installation. On new sodded areas immediately cultivate into the soil to a depth of 4 to 6 inches. Apply when larvae feeding damage first appears and repeat as required. For 4 to 6 year control, increase dosage to 40 to 60 pounds per acre (15 to 20 ounces per 1000 square feet). Keep children and domestic pets off turf until dust has been thoroughly watered in and is completely dry.

Trade: should call in shipping packages only.

Keep out of reach of children and animals. Avoid breathing dust. Avoid contact with skin and eyes. Avoid contamination of food or feedstuffs. Avoid contact with skin and eyes. Avoid contamination of food or feedstuffs.

