## DIRECTIONS FOR USE PANEL 1

PYRENONE 25.5 M A.G. Concentrate is effective. the control of the indicated insects if the applicator follows directions for use as enumerated below.

A	Eara in	Misquiloes
A	5. · · · · · · · · · · · · · · · · · · ·	Mushroom Nes
Bush tert tular	F - 3., 4 2hain	Red flour beetles
Cade et	anals	Rice Aeevils
***P\$1 ***	Grain milles	Saw routhed grain
5 1-12 SF- SH	Granary APPN 5	heeties
a grant far i ven i sen	H, IP AS	Smail living moths
a feet of terk	HIMES	Sc der beetles
f ni sert, rigele-	Hrise Hes	Stable flies
a'r mea ar ms	House thes	Yobacco muths
· · · · · · · · ·	In tan meal moths	Wascs
HAT IS I TRAFPS	- FF	A6hCM areafMorun2
r, store teet wy	Mediterranear frunk ths	

#### INDOOR USE AS A SPACE SPRAY, UNDILUTED

The use of this product in food processing or food handling establishments (bakeries, beverage plants, canneries, food processing plants, hospitals, hotels, industrial installations, meat packing plants, restaurants, schools, supermarkets, and warehouses) should be confined to time periods when the plants or facilities are not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Close windows and doors and shut off ventilating systems. Use a togger or vaporizor adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at a rate of 1/2 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill annessible, exposed stages of files, (including

Ties) mosquitoes, small flying moths, gnats,

.s, hornets, clover mites, cheese mites, cheese skippers, boxelder bugs, earwigs, ants, and the following common stored product pests: granary weevils, rice weevils, confused flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, tobacco moths, yellow mealworms, dark mealworms, grain mites, cadelles, and red flour beetles.

DRIED FRUIT PRODUCTS in storage or being processed for control of confused flour beetles. dried fruit beetles, Indian meal moths, sawtoothed grain beetles. Remove and destroy infested products. As noted above, disperse this product as a fine mist in the air above the trays. and shelves. Do not apply to fruit directly. Use one nunne per 1000 cubic feet of spane. Repeat application at this dosage in any given month-Leale the room closed for one hour. Do not remain in treated areas and ventilate area before. re-entry. This application will kill accessible. exposed stages of dried fruit beetles as well as those pests named above.

STORED FOOD (Stored in multi-walled bags or in most hags) in warehouses, storage rooms, flour mills, grain elevators, granaries, for control of Angoumois grain moths, cadelles, confused flour

Hes, Indian meal moths, Mediterranean flour s, rice weevils, saw-toothed grain beetles.

uw mealworms, cheese mites, cheese skippers, cigarette beetles, grain mites, granary weevils, dark mealworms, red flour beetles Remove and destroy infested products. As noted above, disperse this product

#### Code 980.00

# **PYRENONE**® 25-5 M.A.G. Concentrate

EPA Est., 279-NY-1

#### FOR PROFESSIONAL USE ONLY:

To be applied only by or under the supervision of Pest Control Operators, Mosquito Abatement Districts, Public Health Organizations and other Trained Personnel responsible for Insect Control Programs.

For indoor or outdoor application as a space, area or contact spray. Dependent upon pests to be controlled and the area to be treated, may be applied through Mechanical Aerosol Generators (ULV) as well as conventional fogging or spraying equipment.

#### SPACE AND/OR CONTACT USE AREAS:

Bakeries	Homes
Beverage Plants	Horse Barns
Cannerles	Hospitals
Cattle Barns	Hotels
Dried Food Products	Industrial Instal
in storage or processing	Meat Packing P
lour Mills	Motels
Food Processing Plants	Office Building
Grain Elevators	Zoos
Granaries	<b>Poultry Houses</b>

#### OUTDOOR USE AREAS:

Corrals Drive-in Restaurants Drive-in Theaters Feedlots

Golf Courses **Municipalities** Parks Playgrounds

#### Active Ingredients

Pyrethrins					
*Piperonyl Butoxide, Technical		•			
Petroleum Oil				•	

\*Equivalent to 20% (butylcarbityl) (6-propylpiperonyl) ether and to 5% related compounds

**PYRENONE** — FMC Trademark

## CAUTION: **KEEP OUT OF REACH OF CHILDREN**

See back panel for use and disposal cautions

**FMC** Corporation Agricultural Chemical Division Middleport New York 14105 105

### **Carton Insert**




EPA Reg. No. 279-2918

4	Railroad Cars
•	Restaurants
	Schools
	Stored Food in mult
llations	wall paper or
Plants	cloth bags
	Supermarkets
15	Swine Houses
	Truck Trailers
S	Warehouses

**Recreational Areas** Residences Swine Yards Zoos

										_			_								5%
																					25%
•	•	-	•	•	٠	-		-	•	•	٠	·	٠		•	•	•	-	•	•	20.00
-	-	•					•			-	•		-	-						•	70%
																					100%



FMC 

#### PANEL 2

first by directing spray into corners under paliets, and around stacks of packaged toods hitting as many insects as possible and then treat as a space spray. Infested stored products should be fumigated or treated by other effective methods using an approved product intended for this purpose. Use a dosage of 1 fluid ounce per 1000 cubic feet of room space. Repeat applications as necessary for insect control. Leave the room closed for 1 hour. Do not remain in treated areas and ventilate area before re-entry. This application will k-l accessible, exposed stages of those stored product pests listed above.

#### INDOOR USE AS A SPACE SPRAY, DILUTED

For use in conventional mechanical fogging equipment, to kill flies, fruit flies, mosquitoes, small flying moths, and gnats: Cover or remove exposed food and food handling surfaces. Close room and shut off all air conditioning or ventilating equipment. Dilute 1 part of Pyrenone 25-5 M.A.G. Concentrate plus 49 parts of oil (refined kerosene) and mix well. Apply at the rate of 1-2 fluid ounces per 1000 cubic feet filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

SURFACE SPRAY: To insure maximum control of crawling insects (ants, boxelder bugs, clover mites, drugstore beetles, earwigs, lice, spider beetles) dilute 1 part of Pyrenone 25-5 M.A.G. Concentrate plus 49 parts of oil (refined kerosene) and mix well and apply as a coarse droplet spray using approximately 1 gallon to 750 square feet of surface. Spray walls, floors, and other surfaces of bins and storage and handling areas thoroughly. being especially careful to treat cracks, crevices. and similar hiding places. Treat unloading, handling and processing areas, and inside conveying, processing and handling equipment Spray thoroughly around the base of heavy machinery and equipment. Spray insects directly when possible.

For in transit protection, clean and spray truck beds, boxcars, and ship's hold before loading Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked

To control **Indian meal moth** and **Angoumois meal moth** in stored grain, apply to the surface of binned grain at a rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continuing until early fall. In case of severe infestation, break up webbing with rake before spraying and make a second application in two weeks

As an aid in the control of **mosquitoes, gnats,** wasps, and skippers: treat walls, ceilings, mouldings, screens, door and window frames, beams, light cords, and similar resting places

ANIMAL QUARTER USE (cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray diluted for use in conventional mechanical fogging equipment to kill flies, mosquitoes, small flying moths, and grats. Dilute 1 part of Pyrenone 25-5 M A G. Concentrate plus 49 parts of oil (refined kerosene) and mix well Apply at the rate of 2 fluid ounces per 1000 cubic feet of space above the animals. Direct spray towards the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Vacate treated area and ventilate before reoccupying. Repeat treatment when reinfestation occurs.

IN MUSHROOM PRODUCTION AND PROCESSING: To control mushroom flies or fungus gnats in growing houses, dilute 1 part of Pyrenone 25-5 M A G. Concentrate plus 49 parts

#### PANEL 3

of oil (refined kerosene) and mix well and apply at a rate of 1 ounce per 1000 cubic feet of space Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

OUTDOOR GROUND APPLICATION UNDILUTED To control adult mosquitoes, gnats and small flying moths. For best results, apply when there is a light breeze at approximately 5 miles per hour. Apply from the windward side so the fog will be carried across the area to be treated. Treat shrubbery and vegetation where mosquitoes may rest. Shrubbery and vegetation around stagnant pools, marshy areas and ponds may be treated. Application of this product to any body of water is prohibited. Repeat treatment as required.

Using truck-mounted equipment, the objective is to treat the area with .002 to .0025 pounds of pyrethrins and .010 to .0125 pounds of piperony! butoxide per acre.

PYRENONE 25-5 M.A.G. Concentrate should be applied without further dilution. Dispense fog at the following flow rates depending upon the speed at which the truck is moving:

Truck Speed Miles Per Hour	Flow Rate of PYRENONE 25-5 M.A.G. Concentrate Fluid Ounces Per Minute
5	2 to 2.25 fluid ounces
10	4 to 4.5 fluid ounces

Delivery rate and traveling speeds can be varied slightly so long as the basic objective of .002 to .0025 pounds actual pyrethrins per acre is maintained. Use a swath width of 300 feet for acreage calculations.

To control mosquito larvae, dilute at the rate of 9 ounces per 50 gallons of oil and apply at the rate of 20 to 25 gallons per acre. Apply as a coarse wetting spray to the border or on shallow water of wet lands, swamps, and marshes where mosquito larvae are known to breed

TEMPORARY REDUCTION OF ANNOYANCE from flies, mosquitoes, and small flying moths outdoors. The aforementioned application for outdoor ground application will afford temporary reduction of annoyance from these pests in public gathering places (drive-in restaurants, drive-in theaters, golf courses, municipalities, parks, playgrounds, and recreational area). Direct application into tall grass, shrubbery and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. Application should be made prior to attendance. Repeat as necessary.

In additional outdoor areas (corrals, feed lots, swine lots, and zoos), cover water, drinking fountains, and animal feed before use. Fill area with mist directing application into tall grass, shrubbery, and around lawns where these pests may hover or rest, apply while air is still. Avoid wetting foliage. In zoos, avoid exposure of reptiles to the product. Repeat as necessary.

FOR USE ON ANIMALS: To protect beef and dairy cattle, horses, and dogs from horn files, house files, mosquitoes, and gnats, dilute 1 part of Pyrenone 25-5 M.A.G. Concentrate plus 49 parts oil (refined kerosene) and mix well and apply a light mist sufficient to wet the tips of the hair. To control stable files, horse files, and deer files on beef and dairy cattle and horses, apply 2 ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals to give continued protection

To control the motile stages of blood sucking lice

#### PANEL 4

on beef and dairy cattle, apply the spray to thoroughly wet the hair. Use 2 ounces or less per animal and repeat every 2 to 3 weeks as required.

#### ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment of disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

#### USE CAUTIONS

Harmful if swallowed.

Do not pour or spill near open flame. Remove pets, birds and cover fish aquaria before spraying.

Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing. Wash exposed skin surfaces with soap and water after handling and before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not apply directly to food Do not apply space spray while food processing is underway. Foods should be removed or covered before treatment. Thoroughly wash all food processing surfaces before reuse. After spraying in meat packing plants, bakeries, and the food processing plants, wash with potable water all equipment, benches, shelving, etc. where exposed food will be handled. In case washing is impractical, do not allow food to come in contact with treated surfaces for at least 48 hours. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions given above.

When using in animal quarters, do not apply directly to food, water or food supplements. Wash teats of dairy animals before milking.

#### DISPOSAL CAUTION'S:

Do not reuse empty container. Destroy by perforatine; or crushing and burying in a safe place or return to drum reconditioner.

Terms of sale or use On purchase or use of this product buyer and user agree to the following conditions:

Warranty FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. Except as so warranted the product is sold as is. FMC makes no other warranty express or implied.

Damages Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by huyer or user, failing which buyer or user waives any claim for such damage.

Excuse from non-controllable conditions Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein. Code 980.00

# PYRENONE® 25-5 M.A.G. Concentrate

EPA Est., 279-NY-1

EPA Reg No. 279-2918

## FOR PROFESSIONAL USE ONLY:

To be applied only by or under the supervise rand Pest Control Operators, Mosquito Abatement Districts, Public Health Organizations and other Trained Personnel responsible for Insect Control Programs.

For indoor or outdoor application as a space, area or contact spray Dependent upon pests to be controlled and the area to be treated, may be applied through Mechanical Aerosol Generators (ULV) as well as conventional logging or spraying equipment

#### Active Ingredients

Pyrethrins	5.5
*Piperonyl Butoxide, Technical	25 °.
Petroleum Oil	70° c
	100

\*Equivalent to 20% (butylcarbity!) (6-propylpiperonyl) ether and to 5% related compounds

### **PYRENONE** — FMC Trademark

# CAUTION: KEEP OUT OF REACH OF CHILDREN

See back parently use and disployal captions.



FMC Corporation Agricultural Chemical Division Middleport New York 14105 016

#### ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes. streams, ponds, tidal marshes, and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment of disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

#### USE CAUTIONS

Harmful if swallowed

Do not pour or spiil near open flame Remove pets, birds and cover fish aduaria before spraying

Do not breathe spray mist. Wear mask or respirator of a type recommended by the U.S. Bureau of Mines to give adequate protection against this material. Wear protective clothing Wash exposed skin surfaces with so-ip and water after handling and before eating or smoking Wash all contaminated clothing with suap and hot water before rause.

Do not apply directly to lood. Do not apply space spray while food processing is underway. Foods should be removed or covered before treatment. Thornughly wash all food processing surfaces before reuse. After spraying in meat packing plants, bakeries, and the food processing plants wash with potable water all equipment, benches shelving, etc. where exposed food will be handled. In case washing is impractical, do not allow food to come in contact with treated surfaces for at least 48 hours. Food processing operations do not have to be stopped while applying a wet spray with nare and in accordance with the directions and cautions given above.

When using in animal quarters, do not apply directly to food, water or food supplements. Wash teats of dairy animals before milking.

#### DISPOSAL CAUTIONS

Do not reuse erepty container. Destroy by perforating or crushing and horving in a safe parle or return to drum reconditioner.

# 1 Quart Net Contents

Terms of sale or use On purchase or use of this product buyer and user agree to the following conditions

Warranty FMC warrants that this product as it the time of sale by FMC. (I) conforms to the ingredient statement on the label, and its is reasonably fit for the purposes set forth in the Directions for Use. Except as so warranted the product is sold as is. FMC makes no other warranty express or implied.

Damages Buyer's criuser's exolusive named, for damages for breach of warranty or nog-mentshall be limited to direct damages not - energies the purchase processed and shall not collude incidental or consequential damages. Note: of any claim must be reported to FMC within 10 baye of discovery by boyer or user failing which byger or user waives any crain for such damage.

Excuse from non-controllable conditions Because of certain critical conditions preated or incurred by buyer or user or over which FMC has no control, buyer and user las a condition of purchase or use lassime responsibility for and release FMC from trability arising out of the handling or use of the product altributable to such causes. No person is authorized to vary or waive only statement contained herein.

Read entire Carton insert before using. Use only as directed.

See attached label for instructions for use.

