

CODE
3209



Net Contents
53 GALLONS

KEEP-R-CORN

FUNGICIDE

ACCEPTED
OCT 9, 1972
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 279-2920

DIRECTIONS FOR USE

This product is intended for use only on high-moisture feed grains (barley, shelled corn, oats, sorghum and wheat) to be used in animal feeds. Do not use on grain which might be used for seed, malting or human consumption.

Apply 1.44 to 4.2 U.S. gallons of Keep-R-Corn per ton of grain before storage at harvest time. Dosage rate is dependent upon the moisture content of the grain. Use a moisture tester to determine the moisture content of the grain to be stored. Apply rates as given in the dosage table shown on the label. Thorough coverage of the grain is essential to assure preservation of the grain during the storage period. Apply only by means of a carefully calibrated and metered type of applicator. Follow manufacturers instructions for grain treating procedures. Allow at least one-half hour for Keep-R-Corn to penetrate and be absorbed by the grain before blowing into storage. No waiting is necessary if auger or conveyor transport to storage is used. Do not underdose grain or exceed the maximum dosage rate of 4.2 gallons Keep-R-Corn per ton of grain.

DOSAGE TABLE

Application Rates Per Ton of Feed Grain

% By Weight Moisture In Grain	Gallons Keep-R-Corn
15	1.44
16	1.56
17	1.68
18	1.92
19	2.04
20	2.16
21	2.28
22	2.40
23	2.64
24	2.76
25	2.88
26	3.00
27	3.12
28	3.36
29	3.48
30	3.60
31	3.72
32	3.84
33	3.96
34	4.08
35	4.20

EPA Reg. No. 279-2920

For Use on High Moisture Feed Grains (barley, shelled corn, oats, sorghum and wheat) to Prevent Bacterial and Mold Growth During Storage.

ACTIVE INGREDIENTS:

Acetic Acid 59.90%
Propionic Acid 39.80%

INERT INGREDIENTS: 0.30%
100.00%

CAUTION
KEEP OUT OF REACH OF CHILDREN.

For Emergency Assistance Call
716-735-3765

Harmful if swallowed. Material causes irritation of skin and eyes. Avoid contact with eyes, skin and clothing. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention; Wash skin promptly with soap and water; Remove contaminated clothing. Provide good ventilation during treating, and avoid prolonged exposure in the treated grain storage until the pungent odor has dissipated.

STORAGE AREA PREPARATION

Prepare the storage area by removing dirt, straw and remnants of previously stored grain. Wood Structures (bins, sheds) - Place clean, new polyethylene plastic sheeting on floor. Cover all wooden surfaces (walls, posts, etc.) that will be in contact with the grain with plastic sheeting. Concrete Silos - Line the base with plastic sheeting to a height about 10 feet above ground level. Steel and Galvanized Bins - Line the inner surfaces with plastic sheeting to avoid corrosion of the metal by the treated grain.

Protect treated grain from rain and snow during the storage period. Do not mix treated and untreated grain before or during the storage period.

CARE OF EQUIPMENT

At the end of treating day, flush with water and drain application equipment, conveyors and augers used to transport treated grain to storage area.

CAUTION: Do not contaminate water by cleaning of equipment, or disposal of wastes.

STORED GRAIN INSPECTION

If rain or untreated high moisture grain gets into the treated stored grain it may cause "hot spots" and result in spoilage. Check the stored grain periodically with a temperature probe. If hot spots are located, remove the affected grain and spread to dry. If the grain is to be returned to storage, retreat it with Niagara Keep-R-Corn.

Dealers Should Sell in Original Packages Only.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, N. Y. 14105

CODE
3209



Net Contents
53 GALLONS

KEEP-R-CORN

FUNGICIDE

EPA Reg. No. 179-2990

DIRECTIONS FOR USE
This fungicide is intended for use on corn, sorghum, and grain sorghum. It is effective against the following diseases: ...

ACTIVE INGREDIENTS:
Azinphos Methyl 20.00
Propiconazole 20.00
 INERT INGREDIENTS: 60.00

For Use on High Moisture Food Grains (Corn, Sorghum, Grain Sorghum) and Wheat. For Use on Field and Mill Growth During Storage.

ACTIVE INGREDIENTS:
Azinphos Methyl 20.00
Propiconazole 20.00
 INERT INGREDIENTS: 60.00

CAUTION
 KEEP OUT OF REACH OF CHILDREN.

Emergency Assistance
1-800-4-A-SHIELD

Read the entire label carefully before using this product. It contains important information about safe handling, use, disposal, and first aid. ...

TOXICITY INFORMATION

This fungicide is highly toxic to fish and aquatic life. It is also toxic to birds and mammals. ...

Keep this product out of the reach of children. Do not use it near water. ...

Dealers should sell in Original Packages Only. ...

MANUFACTURED IN U. S. A. BY

