

Code: 31247

Net Weight
50 Pounds

ZINEB 6 DUST

DIRECTIONS FOR USE

Apply at first sign of, or prior to appearance of disease. Repeat at 1 to 2 week intervals as necessary unless otherwise noted. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in () following the crop.

BEANS (7): Downy Mildew, Rust, Anthracnose — Use 35 to 40 pounds per acre. Begin when first blossoms appear.

BEETS (Roots - 0, Tops - 7), TURNIPS (Roots - 0, Tops 7): Cercospora Leaf Spot, Downy Mildew — Use 35 to 40 pounds per acre.

CABBAGE (7), BROCCOLI (7), CAULIFLOWER (7), COLLARDS (10): Downy Mildew, Alternaria — Use 35 to 40 pounds per acre.

CARROTS (Roots - 0, Tops - 7): Alternaria Leaf Blight, Cercospora Leaf Spot — Use 35 to 40 pounds per acre.

CELERY (14): Early and Late Blight — Use 35 to 40 pounds per acre. Begin spraying in the plant bed at 3 to 5 day intervals until the plants are set in field. Continue field application on regular weekly schedule. Remove residue remaining at harvest by stripping, washing or trimming.

EGGPLANT (0): Alternaria, Anthracnose, Phomopsis — Use 35 to 40 pounds per acre.

HOPS (14): Downy Mildew — Use 50 pounds per acre. Apply after training vines and repeat at 7 to 10 day intervals as long as needed. It is necessary that plants be protected when wet or when humid weather prevails.

KALE (10), MUSTARD GREENS (10), SWISS CHARD (7): Cercospora Leaf Spot, Downy Mildew — Use 35 to 40 pounds per acre.

LETTUCE (10): Downy Mildew — Use 35 to 40 pounds per acre. Apply at weekly intervals.

MELONS (CANTALOUPE, WATERMELONS, MUSKMELONS) (5), CUCUMBERS (5), SQUASH (5), PUMPKINS (0): Downy Mildew, Alternaria Blight, Anthracnose, Gummy Stem Blight — Use 40 pounds per acre. Dust underside of leaves.

MUSHROOMS: Verticillium (Brown Spot, Dry Bubble), Dactylium (Mildew, Soft Decay, Cobwebs, Mycogone Bubbles) and Trichoderma (Green Mold) — Use 2 1/2 pounds per 4000 square feet of bed. Apply at 3 to 7 day intervals until buttons begin to form; also between "breaks" as required. Begin applications soon after casing soil is applied.

ONIONS (0), GREEN ONIONS (7): Downy Mildew, Botrytis Leaf Blight (blast), Purple Blotch — Use 40 to 50 pounds per acre.

PEPPERS (0): Phytophthora, Anthracnose, Fire Rot — Use 35 to 40 pounds per acre. Apply at 10 day intervals when disease is evident. Repeat as necessary throughout the season.

FUNGICIDE

EPA REGISTRATION NO. 279-2921

ACTIVE INGREDIENT:	BY WT.
†Zineb	6.00%
INERT INGREDIENTS:	94.00%
	100.00%

†Zinc ethylene bisdithiocarbamate

Zinc as metallic1.43%

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause irritation of Nose and Throat. Do not breathe dust.

SPINACH (10): Downy Mildew (Blue Mold), White Rust — Use 35 to 40 pounds per acre.

TOMATOES (5): Early and Late Blights, Anthracnose, Stemphylium and Septoria Leaf Spots, Cladosporium Gray Mold — Use 35 to 40 pounds per acre. Begin application when disease is in the area and repeat applications at 7 to 10 day intervals or more often if blight conditions are severe.

WHEAT (3 to 4 weeks): Stem and Leaf Rust — Use 30 pounds per acre. Begin applications when plants are in boot to early heading stage, or when rust is first reported in area. Repeat at 7 to 10 day intervals until milk stage.

ORNAMENTALS: Use 35 to 40 pounds per acre. Repeat at 7 to 10 day intervals or more frequently during periods of prolonged rainfall or high humidity. Apply on Roses when foliage first appears. Begin applications when the following diseases threaten or before diseases appear. **ANTHRACNOSE, LEAF SPOT** on Carnations, Chrysanthemums, Hollyhocks, Pansies, and Snapdragons. **BOTRYTIS BLIGHT** on Carnations, Chrysanthemums, Cyclamen, Dahlias, Delphinium and Snapdragons. **BOTRYTIS GRAY MOLD** on Roses and Geraniums. **DOWNY and POWDERY MILDEW** on Asters, Roses, and Snapdragons. For Downy Mildew on Snapdragons, repeat applications at 2 to 4 day intervals. **POWDERY MILDEW** on Carnations, Dahlias, Delphiniums, Hydrangea, and Geraniums. **RUST** on Asters, Chrysanthemums, Delphiniums, Hollyhocks, Hydrangea, Iris, Roses, and Snapdragons.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765

Handwritten:
Zineb 6 Dust
EPA Reg. No. 279-2921

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



79-2921

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:
WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED
DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.
EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals