

Code: 31247




Net Weight  
50 Pounds

# ZINEB 6 DUST

FUNGICIDE

EPA REGISTRATION NO. 279-2921

EPA EST. 279-



ACTIVE INGREDIENT:	BY WT.
Zineb	6.00%
INERT INGREDIENTS:	94.00%
	100.00%

Zinc ethylene bisdithiocarbamate  
Zinc as metallic 1.43%

KEEP OUT OF REACH OF CHILDREN

## CAUTION

For Emergency Assistance Call  
716-735-3765

See Other Panels for Additional Precautionary Information

# FMC

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia PA 19103

EPA REG. No. 279-2921  
PAGE 1 OF 3

(PA-12/82)

PRECAUTIONARY STATEMENTS

Hazards To Humans (and Domestic Animals)

Caution

May cause irritation of Nose and Throat. Do not breathe dust.

During the mixing and loading process, wear protective clothing (long pants, long sleeve shirt, impermeable gloves, hat and boots).

Environmental Hazards

Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

EPA REG. No. 279-2921  
PAGE 2 OF 3

(PA - 10/82)

Apply at first sign of, or prior to appearance of disease. Repeat at 1 to 2 week intervals as necessary unless otherwise noted. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in ( ) following the crop.

**BEANS (7):** Downy Mildew, Rust, Anthracnose — Use 35 to 40 pounds per acre. Begin when first blossoms appear.

**BEETS (Roots - 0, Tops - 7), TURNIPS (Roots - 0, Tops 7):** Cercospora Leaf Spot, Downy Mildew — Use 35 to 40 pounds per acre.

**CABBAGE (7), BROCCOLI (7), CAULIFLOWER (7), COLLARDS (10):** Downy Mildew, Alternaria — Use 35 to 40 pounds per acre.

**CARROTS (Roots - 0, Tops - 7):** Alternaria Leaf Blight, Cercospora Leaf Spot — Use 35 to 40 pounds per acre.

**CELERY (14):** Early and Late Blight — Use 35 to 40 pounds per acre. Begin spraying in the plant bed at 3 to 5 day intervals until the plants are set in field. Continue field application on regular weekly schedule. Remove residue remaining at harvest by stripping, washing or trimming.

**EGGPLANT (0):** Alternaria, Anthracnose, Phomopsis — Use 35 to 40 pounds per acre.

**HOPS (14):** Downy Mildew — Use 50 pounds per acre. Apply after training vines and repeat at 7 to 10 day intervals as long as needed. It is necessary that plants be protected when wet or when humid weather prevails.

**KALE (10), MUSTARD GREENS (10), SWISS CHARD (7):** Cercospora Leaf Spot, Downy Mildew — Use 35 to 40 pounds per acre.

**LETTUCE (10):** Downy Mildew — Use 35 to 40 pounds per acre. Apply at weekly intervals

**MELONS (CANTALOUPE, WATERMELONS, MUSKMELONS) (5), CUCUMBERS (5), SQUASH (5), PUMPKINS (0):** Downy Mildew, Alternaria Blight, Anthracnose, Gummy Stem Blight — Use 40 pounds per acre. Dust underside of leaves.

**MUSHROOMS:** Verticillium (Brown Spot, Dry Bubble), Dactylium (Mildew, Soft Decay, Cobweb), Mycogone (Bubbles) and Trichoderma (Green Mold) — Use 2 1/2 pounds per 4000 square feet of bed. Apply at 3 to 7 day intervals until buttons begin to form; also between "breaks" as required. Begin applications soon after casing soil is applied.

**ONIONS (0), GREEN ONIONS (7):** Downy Mildew, Botrytis Leaf Blight (blast), Purple Blotch — Use 40 to 50 pounds per acre.

**PEPPERS (0):** Phytophthora, Anthracnose, Ripe Rot — Use 35 to 40 pounds per acre. Apply at 10 day intervals when disease is evident. Repeat as necessary throughout the season.

**SPINACH (10):** Downy Mildew (Blue Mold), White Rust — Use 35 to 40 pounds per acre.

**TOMATOES (5):** Early and Late Blights, Anthracnose, Stemphylium and Septoria Leaf Spots, Cladosporium Gray Mold — Use 35 to 40 pounds per acre. Begin application when disease is in the area and repeat applications at 7 to 10 day intervals or more often if Blight conditions are severe.

**WHEAT (3 to 4 weeks):** Stem and Leaf Rust — Use 30 pounds per acre. Begin applications when plants are in boot to early heading stage, or when rust is first reported in area. Repeat at 7 to 10 day intervals until milk stage.

**ORNAMENTALS:** Use 35 to 40 pounds per acre. Repeat at 7 to 10 day intervals or more frequently during periods of prolonged rainfall or high humidity. Apply on Roses when foliage first appears. Begin applications when the following diseases threaten or before diseases appear. **ANTHRACNOSE, LEAF SPOT** on Carnations, Chrysanthemums, Hollyhocks, Pansies, and Snapdragons. **BOTRYTIS BLIGHT** on Carnations, Chrysanthemums, Cyclamen, Dahlias, Delphinium and Snapdragons. **BOTRYTIS GRAY MOLD** on Roses and Geraniums. **DOWNY and POWDERY MILDEW** on Aster, Roses, and Snapdragons. For Downy Mildew on Snapdragons, repeat applications at 7 to 7 day intervals. **POWDERY MILDEW** on Asters, Dahlias, Delphiniums, Hydrangea, and Geraniums. **RUST** on Asters, Chrysanthemums, Delphiniums, Hollyhocks, Hydrangea, Iris, Roses, and Snapdragons.

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use** On purchase of this product buyer and user agree to the following conditions.

**Warranty** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risks of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

EPA REG. No. 277-292!

PAGE 3 OF 3

(PA 12/82)

BEST