Page 1 of 1 EPA REG. No. 279-2922

Code 3061

Net Weight Pounds



Warning

Insecticide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-2922 E

EPA Est., 279-

Active ingredients:

*Carbofuran	 •	•	•	•			•	•	•	• •	 •	•		٠			• •	• •	•	• •	•	5.0%
Inert ingredients:	•	•	•	•	• •	•	•	•	•	•	 •	•	••	•	•	•	• •	••	•		• •	95.0%
																						100.0%

*2,3-Dihydro-2,2-dimethyl-7-benzofuranyl methylcarbamate Furadan® — FMC Trademark U.S. Patent No. 3,474,171

KEEP OUT OF REACH OF CHILDREN

WARNING

First Ald

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or blunt object. Do not induce vomiting or give anything by mouth to an unconscieus person. Get medical attention.

If in eyes, flush with plenty of water for at least 15 minutes. Get medical attention.

Nc: Physician:

Furadan is a reversible cholinesterase inhibitor. Do not use oximes such as 2-PAM Start by giving 2 mg. atropine intramuscularly. According to clinical

response, continue until signs of at opinization occur (dry mouth or dilated pupils).

If in eye, instill one drop of homatropine.

For Emergency Assistance Call 716—735-3765. See Other Panels for Additional Precautionary Information.



FMC Corporation Agricultural Chemical Group 2000 Market Street Philadelphia Pennsylvania 19103

£4 •7 _3

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

May be fatal if swallowed or absorbed through eyes. Do not get in eyes or on clothing. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating or smoking. Hemove contaminated clothing and wash before reuse.

Environmental Hazards

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chamically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not reuse empty container. Dispose of container in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult Federal, State, or local disposal authorities for approved alternative procedures.

Bananas and Plantains-Nematodes, Weevils.

At Planting 1 Ime: Use 25 to 30 grams of Furadan 5% Granules per unit of production (cepa). Apply 16 to 20 grams to the planting hole and 9 to 10 grams to the soil surface (after the hole has been covered). Retreat soil surface with 40 grams at 4 months and again at 8 months after planting.

Established Plantings: Use 40 to 50 grams of Furadan 5% Granules per unit of production (cepa). Make two (2) applications per year. Apply 8 to 10 grams of Furadan 5% Granules around the base of the mother, daughter, and granddaughter plants and 32 to 40 grams of Furadan 5% Granules distributed over an area of 50 cms around the producing writemother, daughter and granddaughter plants).

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty, FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this productare beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.