



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

446634

Ms. Nancy Hilton
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JUL 15 1993

Dear Ms. Hilton:

Subject: Amendment - Supplemental Label for use
in California only

Thiodan 50 WP
EPA Reg. No. 279-1380
Thiodan WSB
EPA Reg. No. 279-3129
Thiodan 3 EC
EPA Reg. No. 279-2924
Your label submissions dated May 24, 1993

The amendments referred to above, submitted in connection with the above registrations under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, are acceptable. It should be understood that acceptance of the supplemental labeling for use only in the State of California to address CAL-EPA's criteria for issuing permits by a County Agriculture Commissioner for the subject product does not constitute the Agency's determination in regard to the reregistration of endosulfan, nor does it represent the Agency's conclusions based upon review of any data, nor should it be construed in any way as signifying the Agency's agreement or determination that the instructions and restrictions appearing on the supplemental labeling for use in California only adequately address the issues of mitigating off-target movement and aquatic exposure. The Agency reserves these decisions until all data submitted in support of reregistration for endosulfan have been reviewed and evaluated. A stamped copy of the label is enclosed for your records.

Sincerely,

A handwritten signature in black ink, appearing to read "G. LaRocca".

George T. LaRocca
Product Manager (13)
Insecticide Rodenticide Branch
Registration Division (H7505C)



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber.

P.S. Note that this acceptance of your label does not relieve you of your obligation to comply with the Worker Protection Standard. If any of your products are covered by this Standard, you are required to submit, and receive the Agency's approval by April 21, 1994, a revised label reflecting the required label statements of 40 CFR 156, published in the FEDERAL REGISTER on August 21, 1992 (57 FR 38102). Please refer to PR Notice 93-7 for further guidance. According to 40 CFR 156, subpart K, "No product to which this subpart applies shall be distributed or sold without amended labeling by any registrant after April 21, 1994. Please call 1-800-777-2185 if you have any labeling questions regarding the Worker Protection Standard.

Code 3212

Net Contents

Thiodan® 3 E.C. Insecticide

For Agricultural or Commercial Use Only.

EPA Reg. No. 279-2924

EPA Est. 279-

Active Ingredients:	By Wt.
*Endosulfan: (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin 3-oxide)	33.7%
Inert Ingredients:**	66.3%
	100.0%

*Thiodan
**Contains Xylene Range Aromatic Solvents.
Thiodan 3 E.C. contains 3 pounds of endosulfan per gallon.

KEEP OUT OF REACH OF CHILDREN



PELIGRO

See Other Panels for Additional Precautionary Information.

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by giving one (1) ounce of syrup of ipecac, if available, or by touching the back of the throat with a finger. Do not induce vomiting or give anything by mouth to a person who is unconscious or convulsing.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin: Remove contaminated clothing and wash skin with soap and water.

If in eyes: Flush eyes with plenty of water. Call a physician immediately.

Note to Physicians: Endosulfan is a central nervous system stimulant absorbable by mouth, inhalation or through contact with skin. It may cause convulsions. There is no specific antidote. Diazepam I.V. is the drug of choice. Barbituric acid derivatives such as Phenobarbital may be used additionally. A neuromuscular blocking agent may be used if convulsions persist. This type of drug may be used *only* if complete control of respiration can be maintained. Epinephrine derivatives are absolutely contraindicated.

For Emergency Assistance Call (800) 331-3148.

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUL 15 1993

FMC

FMC Corporation
Agricultural Chemical Group
Philadelphia PA 19103

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
Recommended, for the pesticide
registered under EPA Reg. No.

279-2924

4/92

PRECAUTIONARY STATEMENTS Hazards to Humans (& Domestic Animals)

Danger

Fatal if swallowed. May be fatal if inhaled or absorbed through the skin. Do not breathe vapors or spray mist. Causes eye damage. Do not get in eyes, on skin, or on clothing. Applicators must wear a long sleeve shirt, trousers, unlined waterproof gloves made of natural rubber or neoprene, full foot covering (such as shoes or boots), head covering (such as a hat), and a pesticide mask or respirator jointly approved by the Mine Safety and Health Administration (MSHA) and the National Institute for Occupational Safety and Health (NIOSH) under the provisions of 30 CFR Part 11 for protection against endosulfan. Mixers and/or loaders must wear a chemical resistant apron and goggles or face shield in addition to all of the protective equipment specified for applicators. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Wash thoroughly with soap and water after handling and before smoking or eating. Remove contaminated clothing and wash with soap and hot water before reuse.

Environmental Hazards

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Due to the risk of runoff and drift, do not apply within a distance of 300 feet of lakes, ponds, streams, and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label.

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

Physical/Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing for 24 hours.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, such as the 48 hour reentry interval established by California, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language primarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

DANGER

Area treated with endosulfan on (insert date). Do not enter without appropriate protective clothing for 24 hours (i.e. 1 State Department of Agriculture's reentry interval, if more restrictive).

In case of accidental exposure to pesticide spray, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

STORAGE AND DISPOSAL

Pesticide Storage

Do not store in or around the home.

Do not store below 20°F, (-7°C).

Do not use or store near heat, open flame or hot surfaces.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC (800-331-3148).

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

Returnable/Refillable Sealed Containers

Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

See supplemental label for directions for use in California

GENERAL INSTRUCTIONS

Not For Use Or Storage In Or Around The Home.

Do not use in undiluted form.

Apply the listed amount per acre when insects first appear and repeat as required, unless otherwise noted, to maintain effective control. Use in sufficient water for thorough coverage of listed crops, unless otherwise noted. Coverage of upper and lower leaf surfaces is essential for good control. For ground application, apply recommended amount of pesticide in a minimum of 10 gallons of water per acre on vegetable or row crops. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides.

When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 1 gallon of finished spray per acre on vegetable and field crops, unless otherwise noted, and a minimum of 20 gallons of finished spray per acre on fruit and nut trees and on vines. Where more than 2 quarts of this material are recommended, mix with sufficient water to provide a minimum of finished spray equal to twice the amount of Thiodan 3 E.C. insecticide used.

Do not plant root crops other than carrots, potatoes, sweet potatoes, and sugar beets as follow-up crops. Observe days interval between last application and harvest indicated by number in () following the crop.

Alfalfa (Grown for Forage) (21)

Insects Controlled	Rate of Application	Method of Application
Meadow Spittlebug	2/3 pint per acre	Apply one week after eggs begin to hatch or when first small masses of froth are produced in the crowns of the plant which is usually about the time the alfalfa is 4 to 6 inches tall. Apply in a minimum of 10 gallons of water per acre by ground or 2 gallons of water per acre by air.

Do not graze or cut forage within 21 days after treatment.
 Do not apply more than once per cutting.
 Do not make more than 3 applications per year.
 Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.

Apples (21)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Apple Aphid, Rosy Apple Aphid, Woolly Apple Aphid) Apple Rust Mite Green Fruitworm Tarnished Plant Bug Tentiform Leafminers White Apple Leathopper (first generation)	2/3 quart per 100 gallons or a maximum of 3 1/2 quarts per acre	Applications made at pink and/or petal fall provide best control of Tarnished Plant Bug and Green Fruitworm. For best control of first generation White Apple Leathopper, apply when nymphs first appear on leaves. If necessary, prior to petal fall, use 1 quart per 100 gallons to control Apple Aphids and Rosy Apple Aphids. For control of Tentiform Leafminers, make first application as soon as moth flight begins. A second application should be made 10 days later.

Do not feed pomace from treated apples to livestock.
 Do not feed cull fruits to animals or allow livestock to graze in treated orchards.
 Do not make more than 2 applications during the fruiting period.
 Do not make more than 3 applications per year.
 Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Apricots (21), Nectarines (21), Peaches (21)

Insects Controlled	Rate of Application	Method of Application
Peachtree Borer Lesser Peachtree Borer	1 quart per 100 gallons; West Coast—2/3 to 1 quart per 100 gallons; South-eastern States—2 to 3 1/2 quarts per 100 gals.	Best control is obtained with a single application post-harvest during the first week of September. Spray all bark areas from ground level to lower scaffold limbs.

Do not feed cull fruits to animals or allow livestock to graze in treated orchards.
 Do not make more than 2 applications per year.
 Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Apricots (30), Nectarines (30), Peaches (30)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Black Cherry Aphid, Black Peach Aphid, Green Peach Aphid, Rusty Plum Aphid) Catfacing insects (Stink Bug type) Green Fruitworm Peach Silver Mite Peach Twig Borer	2/3 quart per 100 gallons or a maximum of 3 1/2 quarts per acre; West Coast—do not use more than 4 quarts per acre	Make applications when insects appear or feeding is noticed.

Do not feed cull fruits to animals nor allow livestock to graze in treated orchards.
 Do not make more than 2 applications per year.
 Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Artichokes (7)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Artichoke Aphid, Green Peach Aphid) Artichoke Plume Moth Whitely	1 to 1 1/3 quarts per acre 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed. Apply in a minimum of 20 gallons of water per acre by ground and in a minimum of 10 gallons of water per acre by air.

Do not make more than 2 applications per year.
 Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2 2/3 quarts) per acre per year.

Barley, Oats, Rye, Wheat

Insects Controlled	Rate of Application	Method of Application
Army Cutworm	2/3 quart per acre	For aerial application, apply in 2 gallons of diesel fuel oil per acre.
Cereal Leaf Beetle (Illinois, Indiana, Michigan and Ohio only)	1/3 to 2/3 quart per acre	Apply when small larvae are readily found in the field. For aerial application, use a minimum of 1 to 2 gallons of water per acre.
Aphids	2/3 to 1 quart per acre	Make applications when insects appear or feeding is noticed.

Do not apply after heads begin to form.
Do not feed treated forage to livestock.
Do not make more than 2 applications per year.
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.

Beans, Succulent and Dry (except Lima Beans) (3) and Southern Field Peas (Succulent type, including Black-eyed Peas, Crowder Peas and Southern Peas) (3)

Insects Controlled	Rate of Application	Method of Application
Black Bean Aphid Bean Leaf Skeletonizer Cowpea Curculio Cucumber Beetles Flea Beetles Green Stink Bug Leafhoppers Mexican Bean Beetle	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed. For control of Cowpea Curculio, make 3 applications at 5-day intervals starting when the pods are 1/2 inch long.
Aphids Armyworms Western Bean Cutworm Whitely	1 1/2 quarts per acre	

Do not feed treated threshings or allow livestock to graze in treated fields.
Do not use on Lima Beans that are to be removed from the field for processing.
Do not make more than 3 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Blueberries

Insects Controlled	Rate of Application	Method of Application
Blueberry Bud Mite	2 quarts per 300 gallons	Apply immediately after harvest and repeat 6 to 8 weeks later.

Do not apply after buds are well formed.
Do not make more than 2 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Broccoli (7), Brussel Sprouts (14), Cabbage (7), Cauliflower (14)

Insects Controlled	Rate of Application	Method of Application
Cabbage Aphid Cabbage Looper Cross-striped Cabbage-worm Diamondback Moth larvae Flea Beetles Harlequin Bug Imported Cabbageworm Leafhoppers Stink Bugs	1 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Armyworms Cutworms Whitely	1 1/2 quarts per acre	

Do not make more than 4 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Carrots (7)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Leafhoppers	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Armyworms Flea Beetles Whitely	1 1/2 quarts per acre	

Do not use tops for food or feed.
Do not make more than one application per year.
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.

Celery (4)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Cabbage Looper Leafhoppers	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Armyworms Flea Beetles Whitely	1 1/2 quarts per acre	

Do not make more than one application per year.
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.

Celery (7)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid	2/3 quart per acre	Make applications when insects appear or feeding is noticed.

Do not make more than 2 applications per year.
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.

Cherries (21)

Peachtree Borer Lesser Peachtree Borer	1 quart per 100 gallons; <i>West Coast:</i> 2/3 to 1 quart per 100 gallons	Thoroughly wet all bark from ground to scaffold limbs.
Black Cherry Aphid Green Fruitworm Plum Rust (Nursery) Mite	2/3 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
<i>Pacific Northwest only:</i> Eyespotted Bud Moth Fruitree Leafroller		For Bud Moth control, apply at "popcorn" stage. For Leafroller control, apply during pre-pink stage of growth.
<i>Michigan only:</i> Mineola Moth	1 1/2 quarts per 100 gallons	Apply in the delayed dormant period.

Do not feed cull fruits to animals or allow livestock to graze in treated orchards.
Do not make more than 2 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Cherry, Peach, Plum Nursery Stock Dip

Insects Controlled	Rate of Application	Method of Application
Peachtree Borer	2 2/3 quarts per 40 gallons	Mix thoroughly. Immerse trees so that the roots and crowns are covered well above the grafting bud scar.

Wear rubber gloves during the dipping operation.
Plant immediately or dry before returning stock to storage.

Citrus (Non-Bearing Trees and Nursery Stock)

Insects Controlled	Rate of Application	Method of Application
Citrus Aphid	2/3 quart per 100 gallons or a maximum of 3 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Do not apply to bearing trees or trees that will bear fruit within twelve months. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Collards (21)

Insects Controlled	Rate of Application	Method of Application
Aphids Cabbage Looper Diamondback Moth larvae Fall Armyworm Flea Beetles Harlequin Bug Imported Cabbageworm Leafhoppers	1 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1 1/2 quarts per acre	
Do not make more than one application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.		

Cotton

Insects Controlled	Rate of Application	Method of Application
Aphids	1/2 to 1 quart per acre	Make applications when insects appear or feeding is noticed. For control of aphids, thorough coverage is important.
Boll Weevil	2/3 to 2 quarts per acre	
Bollworm Cabbage Looper Cotton Leafperforator Cotton Leafworm Flea-hoppers Lygus Bugs Stink Bugs Tobacco Budworm	1 1/2 to 2 quarts per acre	
Thrips	2 quarts per acre	
Applications may be made using ground or aerial application equipment. A higher rate should be used under heavy pest pressure. Do not apply after bolls open. Do not graze dairy or meat animals in treated fields. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Cucumbers (2), Melons (2), Pumpkins (2), Summer and Winter Squash (2)

Insects Controlled	Rate of Application	Method of Application
Aphids Cucumber Beetles Melonworm Pickleworm Rindworm (on watermelons) Squash Beetle Squash Bug Squash Vine Borer Striped Flea Beetle	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed. For Squash Vine Borer control, apply weekly to flower buds, stems, and vines beginning when moths first appear.
Cabbage Looper Omnivorous Leafroller Whitefly	1 1/2 quarts per acre	
Do not make more than 6 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Eggplant (1)

Insects Controlled	Rate of Application	Method of Application
Blister Beetle Colorado Potato Beetle Flea Beetles Green Peach Aphid Green Stink Bug	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1 1/2 quarts per acre	
Do not make more than 2 applications per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.		

Grapes (7)

Insects Controlled	Rate of Application	Method of Application
Grape Leafhopper Grape Phylloxera (leaf form) Rose Chafer	2/3 quart per 100 gallons c: 1 1/2 to 2 quarts per acre	Make applications when insects appear or feeding is noticed.
Do not use on Concord variety as severe injury is likely to occur. Ground application is preferred. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Kale (21)

Insects Controlled	Rate of Application	Method of Application
Cabbage Flea Beetle Harlequin Bug Imported Cabbageworm	1 quart per acre	Make applications when insects appear or feeding is noticed.
Do not make more than one application per year. Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.		

Lettuce (14)

Insects Controlled	Rate of Application	Method of Application
Cabbage Looper Diamondback Mnth larvae Green Peach Aphid Imported Cabbageworm Leafhoppers	1 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed
Armyworms Whitefly	1 1/2 quarts per acre	
On Head Lettuce: Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. On Leaf Lettuce: Do not make more than 2 applications per year. Do not feed crop refuse to livestock. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Mustard Greens (21)

Insects Controlled	Rate of Application	Method of Application
Aphids Cabbage Looper Diamondback Moth larvae Fall Armyworm Flea Beetles Harlequin Bug Imported Cabbageworm Leafhoppers	1 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1 1/2 quarts per acre	
Do not make more than one application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.		

Pears (7)

Insects Controlled	Rate of Application	Method of Application
Green Fruitworm Tarnished Plant Bug	2/3 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre	Make applications at white bud or petal fall when insects appear or feeding is noticed.
Pear Psylla	3 1/2 quarts per acre	Apply in a minimum of 10 gallons of water for aerial application or in 300 gallons of water for dilute application. Apply when adults are first observed or nymphs are small and repeat to maintain control.
Pear Rust Mite Conspere Stink Bug (foliar treatment)	2/3 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre	Make applications when insects appear or feeding is noticed. Stink Bugs must be wet by spray to obtain control.
Conspere Stink Bug (soil treatment)	2/3 quart per 100 gallons; 200 to 400 gallons per acre	Apply to orchard floor and around trees prior to bloom.
Pear Leaf Blister Mite	1/2 to 2/3 quart per 100 gallons	Apply as a post harvest or dormant treatment.

Do not feed cull fruits to animals or allow livestock to graze in treated orchards.
Do not make more than 2 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.
Note: Aerial application may not result in satisfactory control and should only be employed if impossible to apply by ground.

Peas (Seed Crop Only) (1)

Insects Controlled	Rate of Application	Method of Application
Pea Aphid Pea Weevil	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.

Use only on peas to be harvested by combine.
Do not feed treated vines or threshings to livestock or allow livestock to graze in treated fields.
Do not make more than 2 applications per year.
Do not exceed a maximum of 1.5 lbs. active ingredient (i.e., 2 quarts) per acre per year.

Pecans

Insects Controlled	Rate of Application	Method of Application
Black Pecan Aphid Pecan Nut Casebearer Spittlebug	1 quart per 100 gallons	Make applications when insects appear or feeding is noticed. For Casebearer, apply when eggs of first generation appear on the tips of the young nuts. Another application may be required after the second generation of eggs is deposited. For Spittlebug, apply when first leaves are half grown and repeat as required.
Pecan Leaf Phylloxera	2/3 to 1 quart per 100 gallons	Apply when nymphs appear and before they are enclosed in plant tissue. For high populations, use the higher rate.

Do not apply after shuck split.
Do not graze livestock on orchard crops or grasses in treated areas.
Do not make more than 2 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Peppers (1)

Insects Controlled	Rate of Application	Method of Application
Flea Beetles Green Peach Aphid Hornworms Pepper Maggot	2/3 quart per acre	Make applications when insects appear or feeding is noticed.

Do not make more than 2 applications per year.
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.

Peppers (4)

Insects Controlled	Rate of Application	Method of Application
Armyworms Flea Beetles Green Peach Aphid Hornworms Leafhoppers Pepper Maggot	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1 1/2 quarts per acre	

Do not make more than 2 applications per year.
Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2 2/3 quarts) per acre per year.

Pineapple (For Fresh Market Only) (7)

Insects Controlled	Rate of Application	Method of Application
Pineapple Fruit Mite	2 to 2 2/3 quarts per acre	Make applications when insects appear or feeding is noticed. Apply at intervals of 7 to 10 days, if necessary, particularly during the 40 day period of blooming.

Do not feed forage or pineapple by-products to livestock.
Do not make more than 2 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Plums (7), Prunes (7)

Insects Controlled	Rate of Application	Method of Application
Fruitree Leafroller (Pacific Northwest Only)	2/3 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre	Apply during pre-pink stage of growth when insects appear or feeding is noticed.
Aphids (including Hop Aphid, Leafcurl Plum Aphid, Thistle Aphid) Plum Rust (Nursery) Mite	2 2/3 to 3 1/2 quarts per acre	For control of aphids, apply when eggs hatch during pre-bloom or petal fall. Summer applications should be made before leaves curl.
Peach Twig Borer	2/3 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Lesser Peachtree Borer	1 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre	Apply spray to thoroughly wet the trunk and main branches.
Peachtree Borer	1 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre. West Coast--2/3 to 1 quart per 100 gallons or 2 2/3 to 3 1/2 quarts per acre	Spray thoroughly to cover all bark areas from ground to scaffold limbs.

Do not allow livestock to graze on orchard crops or grasses in treated areas.
Do not make more than 2 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Potatoes (1)

Insects Controlled	Rate of Application	Method of Application
Aphids Armyworms Colorado Potato Beetle Green Stink Bug Leaf-footed Bug Plant Bugs Potato Flea Beetles Potato Leafhopper Potato Tuberworm Threelined Potato Beetle	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
European Corn Borer Potato Psyllid	1 to 1 1/2 quarts per acre	
False Chinch Bug Whitefly	1 1/2 quarts per acre	
Do not make more than 6 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Safflower

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Whitefly	1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Do not apply after flower heads open. Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2 2/3 quarts) per acre per year.		

Spinach (21)

Insects Controlled	Rate of Application	Method of Application
Armyworms Crown Mite Flea Beetles Green Peach Aphid Leafhoppers	1 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1 1/2 quarts per acre	
Do not make more than one application per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1 1/2 quarts) per acre per year.		

Strawberries (4)

Insects Controlled	Rate of Application	Method of Application
Meadow Spittlebug Strawberry Aphid Winged Plant Bug Whitefly	1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed. Do not reapply within 15 days or more than twice during a 35 day period when fruit is present.
Cyclamen Mite	2 2/3 quarts per acre in 400 gallons	Make applications when insects appear or feeding is noticed. Thoroughly wet the foliage, stem and crown of the plant. For multiple applications, do not apply at intervals less than 35 days when fruit is present.
Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Strawberries—Northwest Use Only

Insects Controlled	Rate of Application	Method of Application
Garden Symphylan (aids in reducing damage)	1 1/2 quarts per 100 gallons	Mix thoroughly. Dip entire plant. When immersing bundles of plants, make certain any trapped air is forced out to assure thorough wetting of entire plant.
Wear rubber gloves during the dipping operation. Drain and allow plants to dry before setting them out in the field.		

Sugar Beets (30)

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Meadow Spittlebug	2/3 to 1 1/2 quarts per acre	Make application(s) when insects appear or feeding is noticed. Make one or two applications per year as needed for control.
Sugar Beet Webworm	1 to 1 1/2 quarts per acre	
Whitefly	1 1/2 quarts per acre	
Do not feed treated tops to livestock. Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2 2/3 quarts) per acre per year.		

Sunflowers (1)

Insects Controlled	Rate of Application	Method of Application
Sunflower Moth	1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed. Make the first application at the onset of bloom. Allow 7 days between repeat applications.
Do not feed treated forage to livestock. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Sweet Corn (Fresh Vegetable Use Only) (1)

Insects Controlled	Rate of Application	Method of Application
Corn Leaf Aphid Whitefly	1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Corn Earworm	2 quarts per acre	Apply when silks first appear and continue until they begin to dry. Allow 5 days between applications.
Do not apply to Sweet Corn to be processed. Do not feed treated forage or ensilage to livestock or allow livestock to graze in treated fields. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Sweet Potatoes (1)

Insects Controlled	Rate of Application	Method of Application
Sweet Potato Flea Beetle Sweet Potato Weevil	2/3 quart per acre	Make applications when insects appear or feeding is noticed. For flea beetle control, begin applications shortly after transplanting or as soon as flea beetles appear. For Sweet Potato Weevil control, apply in sufficient water for complete coverage. Applications may be made to transplants in the nursery and/or in the field. Repeat applications may be made as necessary. For Sweet Potato Weevil control in a clean up program, apply at a rate of 2 2/3 quarts per acre to the soil under the sweet potato nursery beds and to the area immediately surrounding the nursery beds. Do not place the seed tubers directly on the ground treated with Thiodan®.
Whitefly	1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Banded Cucumber Beetle larvae (aids in control) (South Central States and Puerto Rico Only)	1 1/2 to 2 1/2 quarts per acre broadcast or 1/2 to 1 quart per acre on a 16 inch band over the row (48 inch row spacing)	Work into the soil to a depth of approximately 3 inches. Treatment should be made just prior to plant set.

Do not feed cull potatoes to livestock or allow livestock to graze in treated fields.
Do not make more than 3 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Tobacco (5)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Green Peach Aphid, Tobacco Aphid) Budworm Cabbage Looper Flea Beetles Hornworms	Seed Bed: 2/3 quart per 100 gallons	Make applications when insects appear or feeding is noticed. Apply about 6 gallons of finished spray per 100 square yards.
Green June Bug larvae	Plant Bed: 2/3 pint per 100 gallons	Make applications when insects appear or feeding is noticed. Drench at a rate of 1 gallon per square yard.
Aphids (including Green Peach Aphid, Tobacco Aphid) Budworm Cabbage Looper Flea Beetles Green June Bug larvae Hornworms	Field: 2/3 to 1 1/2 quarts per acre	Make applications when insect activity or feeding is noticed.
Stink Bugs	Field: 2/3 quart per 100 gallons	

Do not make more than 6 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Tomatoes (Field and Greenhouse) (2)

Insects Controlled	Rate of Application	Method of Application
Aphids Blister Beetles Colorado Potato Beetle Flea Beetles Tomato Hornworm Webworm (California Only)	2/3 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Cabbage Looper Stink Bugs	1 to 1 1/2 quarts per acre	
Tomato Fruitworm Tomato Russet Mite Whitefly Yellowstriped Armyworm	1 1/2 quarts per acre	
Whitefly	2/3 quart per 100 to 200 gallons of water	

Do not make more than 6 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Walnuts

Insects Controlled	Rate of Application	Method of Application
Walnut Aphid	2 to 3 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.

Do not apply after husk split.
Do not graze livestock on orchard crops of grasses in treated areas.
Do not make more than 2 applications per year.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

COMMERCIALLY GROWN ORNAMENTALS

Ornamental Plants

Leatherleaf Fern (Leather Holly Fern)

Insects Controlled	Rate of Application	Method of Application
Leatherleaf Fern Borer	2/3 quart per 100 gallons	Begin treatment when first larval feeding is observed in the mid-vein area at the base of the leaflets. Repeat at intervals of 2 to 3 weeks as necessary.

Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Ornamentals (Greenhouse, and Out-of-Doors)

Insects Controlled	Rate of Application	Method of Application
Aphids Cyclamen Mite Rose Chafer Whitefly	2/3 quart per 100 gallons	Make applications when insects appear or feeding is noticed.

On chrysanthemums, best results will be obtained if applied before plants flower.
Do not apply to "Bonafon Deluxe", "Fred Shoemith", and "White Knight" chrysanthemums as injury may result.
Do not use on Birch trees.
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Ornamental Trees and Shrubs

Dogwood, Lilac

Insects Controlled	Rate of Application	Method of Application
Dogwood Borer Lilac Borer	$\frac{2}{3}$ to $1\frac{1}{3}$ quarts per 100 gallons ($1\frac{1}{3}$ – $2\frac{2}{3}$ teaspoons per gallon)	Apply in early June and repeat in 10 to 14 days. Drench all bark areas down to the ground level.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Douglas Fir (Grown for Ornamentals, Nurser; Stock or Christmas trees)—Pacific Northwest Only

Insects Controlled	Rate of Application	Method of Application
Aphids Cooley Spruce Gall Adelgid Douglas Fir Needle Midge	$\frac{2}{3}$ quart per 100 gallons ($1\frac{1}{3}$ teaspoons per gallon)	Make applications when insects appear or feeding is noticed. For control of gall adelgid, apply when white cottony tufts appear. For control of needle midge, apply in late April or early May just before buds open.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Pines (Austrian, Jack, Red, Scotch, White)

Insects Controlled	Rate of Application	Method of Application
Zimmerman Pine Moth	1 quart per 100 gallons (2 teaspoons per gallon)	Apply in mid-April and again, if necessary, in late fall. Thoroughly wet bark and main stem, especially where branches join main stem.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Shade Trees (except Birch), Shrubs

Insects Controlled	Rate of Application	Method of Application
Aphids	$\frac{2}{3}$ quart per 100 gallons ($1\frac{1}{3}$ teaspoons per gallon)	Make applications when insects appear or feeding is noticed.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Spruce (New England Area Only)

Insects Controlled	Rate of Application	Method of Application
Spruce Gall Aphid	$\frac{2}{3}$ quart per 100 gallons ($1\frac{1}{3}$ teaspoons per gallon)	Apply in late April or early May when aphids are present but before galls are formed.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Taxus

Insects Controlled	Rate of Application	Method of Application
Taxus Bud Mite	$\frac{2}{3}$ quart per 100 gallons ($1\frac{1}{3}$ teaspoons per gallon)	Make 3 to 5 applications beginning in mid-May. Thoroughly spray foliage, twigs, and bark.
Black Vine Weevil	$1\frac{1}{3}$ quarts per 100 gallons ($2\frac{2}{3}$ teaspoons per gallon)	Spray thoroughly and drench the soil under the plants. Apply when the weevils first appear and repeat in 14 days.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

ACCEPTED
with COMMENTS
in EPA Letter Dated

CALIFORNIA SUPPLEMENTAL LABEL

JUL 15 1993

PROCEDURES FOR REDUCING RISK OF CONTAMINATION OF
AQUATIC ECOSYSTEMS

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
279-2924

FOR CROP USES LISTED ON THE PRODUCT LABEL

Endosulfan may be a source of contamination in aquatic ecosystems as a result of run-off from agricultural land due to discharge of irrigation water or during periods of heavy rainfall.

To minimize the chance of aquatic impact, the following conditions must be met to allow endosulfan applications:

DIRECTIONS FOR USE (General)

IN ADDITION TO SPECIFIC INSTRUCTIONS UNDER DIRECTIONS FOR USE ON THE PRINCIPAL LABELS, THE FOLLOWING CONDITIONS MUST ALSO BE FOLLOWED FOR USE IN CALIFORNIA:

1. Do not apply this product to fields during irrigation or for 72 hours following an irrigation.
2. Do not apply this product through any type of irrigation system.
3. Do not apply more than 1.5 lbs. a.i./acre/application or more than 3.0 lbs. a.i./acre/season. (Orchard crops and strawberries exempted on rate/acre/application restriction.)
4. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.
5. Do not apply by ground or air within a distance of 300 feet to natural bodies of water including lakes, ponds, streams and estuaries.

FOR DRIFT CONTROL

1. All application equipment should be properly maintained and calibrated using appropriate carriers.
2. Do not make aerial or ground applications during temperature inversions.
3. Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 10 mph. Do not discharge at a height of more than 10 feet above

the crop or target.

4. For aerial application a positive shutoff system shall be in place to control flow of liquid to spray nozzles as follows: each individual nozzle shall be equipped with a check valve and the flow controlled by a suckback device or a boom pressure release device; or each nozzle shall be equipped with a positive action valve.
5. For aerial applications, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor vortices. The working boom length must not exceed 75 percent of wing span or rotor diameter.
6. Aircraft boom pressure shall not exceed 40 pounds per square inch.
7. Aircraft nozzles shall be equipped with orifices directed backward parallel to the horizontal axis of the aircraft in flight.
8. Aircraft nozzles shall not be equipped with any device or mechanism which would cause a sheet, cone, fan, or similar type dispersion of the discharged material except as otherwise provided.
9. Fixed wing aircraft and helicopters operating in excess of 60 mph shall be equipped with jet nozzles having an orifice not less than 1/16 inch in diameter (D-4 or larger).
10. Helicopters operating at 60 mph or less shall be equipped with nozzles having an orifice not less than 1/16 inch in diameter. A number 46 or equivalent whirlplate may be used.
11. For ground applications, equipment other than handguns shall be equipped with nozzles having an orifice not less than 1/16 inch in diameter or larger. Boom nozzles shall be operated at a boom pressure not to exceed 30 pounds per square inch.

ACCEPTABLE SITES FOR ENDOSULFAN APPLICATION

1. FIELDS, WITH NO DRAINS, HAVING CONTAINMENT THROUGH BERMS (able to hold irrigation water and normal rainwater), LEVEES AND NON-DRAINING CIRCULATION SYSTEMS.
2. FIELDS WITH DRAINS, BUT DRAINAGE FLOWS TO SUMP, FARM POND, EVAPORATION POND OR RECIRCULATION SYSTEM WITHOUT DRAINAGE (uncontrollable overflow acceptable, e.g., 10 or 100 year flood). Only grower maintained recirculation systems without drainage are acceptable.

3. FIELDS WITH SURFACE DRAINS THAT NATURALLY FLOW ONTO PASTURELAND OR FALLOW FIELDS THAT HAVE NO CHANNELED DRAINAGE AND ARE AT LEAST 300 FEET LONG.

UNACCEPTABLE SITES FOR ENDOSULFAN APPLICATION

1. FIELDS DRAINING DIRECTLY INTO SURFACE WATERS SUCH AS STREAMS, RIVERS, LAKES, PONDS, MARSHES, BAYS, ESTUARIES, OR THE OCEAN.
2. FIELDS DRAINING INTO SURFACE DRAINAGE DITCHES OR CANALS THAT FLOW DIRECTLY INTO SURFACE WATERS SUCH AS STREAMS, RIVERS, LAKES, PONDS, MARSHES, BAYS, ESTUARIES, OR THE OCEAN.
3. ANY AND ALL FIELDS WITH SURFACE DRAINAGE PATTERNS IN THE MOSS LANDING DRAINAGE SYSTEM HAVING MAIN ARTERIES THAT DRAIN INTO THE ELKHORN SLOUGH (northern portion of Monterey County).
4. ANY AND ALL FIELDS WITH DRAINAGE PATTERNS IN THE SAN DIEGO CREEK DRAINAGE SYSTEM HAVING MAIN ARTERIES THAT DRAIN INTO NEWPORT BAY.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.