


Code 1593

Supplemental Label for Use in California



# 3 EC Insecticide

For Agricultural or Commercial Use Only.

EPA Reg. No. 279-2924

EPA Est. 279-FL-1<sup>®</sup>; 279-IL-1<sup>®</sup>; 279-IA-1<sup>®</sup>

Superscript used is first letter of lot number stamped on container.

Active Ingredient:	By Wt.
*Endosulfan: (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin 3-oxide).....	33.7%
Inert Ingredients:** .....	66.3%
	<hr/> 100.0%

\*Thiodan  
 \*\*Contains Xylene Range Aromatic Solvent.  
 Thiodan 3 EC contains 3 pounds of endosulfan per gallon.

KEEP OUT OF REACH OF CHILDREN


 **DANGER-POISON**   
**PELIGRO**

See Other Panels for Additional Precautionary Information.

PRECAUCION AL USUARIO: Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)



FMC Corporation  
 Agricultural Products Group  
 Philadelphia PA 19103



### STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center immediately. If possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce vomiting by giving one (1) ounce of syrup of ipecac, if available, or by touching the back of the throat with a finger. Do not induce vomiting or give anything by mouth to a person who is unconscious or convulsing.

If inhaled: Remove victim to fresh air. Apply artificial respiration if indicated. Get medical attention.

If on skin: Remove contaminated clothing and wash skin with soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician immediately.

Note to Physicians: Endosulfan is a central nervous system stimulant absorbable by mouth, inhalation or through contact with skin. It may cause convulsions. There is no specific antidote. Diazepam I.V. is the drug of choice. Barbituric acid derivatives such as Phenobarbital may be used additionally. A neuromuscular blocking agent may be used if convulsions persist. This type of drug may be used *only* if complete control of respiration can be maintained. Epinephrine derivatives are absolutely contraindicated. This formula contains petroleum hydrocarbons (Xylene range aromatic solvent). Care should be taken to prevent aspiration because of the possibility of chemical pneumonia or pulmonary edema due to the organic solvent in the formulation.

For Emergency Assistance Call (800) 331-3148.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans (and Domestic Animals)

**Danger**  
 Fatal if swallowed. Corrosive. Causes irreversible eye damage. Do not get in eyes, on skin, or on clothing. May be fatal if inhaled or absorbed through skin. Do not breathe vapors or spray mist. Do not contaminate food or feed. Keep out of reach of domestic animals. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides.

#### Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Coveralls over long-sleeved shirt and long pants; Chemical-resistant gloves, such as Barrier Laminar or Viton ≥ 14 mils; Chemical-resistant footwear plus socks; Protective eyewear; Chemical-resistant headgear for overhead exposure; a Respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G); and add a Chemical-resistant apron when cleaning equipment, mixing or loading.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statement

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d) (4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Due to the risk of runoff and drift, do not apply within a distance of 300 feet of lakes, ponds, streams, and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic are important resources. Do not contaminate water by cleaning equipment or disposal of equipment washwaters. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label.

This pesticide is toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

### Physical/Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants;
- Chemical-resistant gloves, such as Barrier Laminate or Viton  $\geq$  14 mils;
- Chemical-resistant footwear plus socks;
- Protective eyewear;
- Chemical-resistant headgear for overhead exposure.

### STORAGE AND DISPOSAL

#### Pesticide Storage

Do not store in or around the home.

Do not store below 20°F, (-7°C).

Do not use or store near heat, open flame or hot surfaces.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lid and close tightly. Do not put concentrate or dilute materials into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: 1 (800) 331-3148.

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

#### Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

#### Returnable/Reillable Sealed Containers

Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

This Product is intended for Use in California under this supplemental label only. Be sure to read and follow the required procedures for reducing the risk of contamination of aquatic ecosystems in California.

### GENERAL INSTRUCTIONS

Not For Use Or Storage In Or Around The Home.

Do not use in undiluted form.

Apply the listed amount per acre when insects first appear and repeat as required, unless otherwise noted, to maintain effective control. Use in sufficient water for thorough coverage of listed crops, unless otherwise noted. Coverage of upper and lower leaf surfaces is essential for good control. For ground application, apply recommended amount of pesticide in a minimum of 10 gallons of water per acre on vegetable or row crops unless otherwise noted. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides.

When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 1 gallon of finished spray per acre on vegetable and field crops unless otherwise noted, and a minimum of 20 gallons of finished spray per acre on fruit and nut trees and on vines. Where more than 2 quarts of this material are recommended, mix with sufficient water to provide a minimum of finished spray equal to twice the amount of Thiodan 3 E.C. insecticide used.

Do not plant root crops other than carrots, potatoes sweet potatoes, and sugar beets as follow-up crops. Observe days interval between last application and harvest indicated by number in ( ) following the crop.

**PROCEDURES FOR REDUCING RISK OF CONTAMINATION OF AQUATIC ECOSYSTEMS**

**FOR CROP USES LISTED ON THE PRODUCT LABEL**

Endosulfan may be a source of contamination in aquatic ecosystems as a result of run-off from agricultural land due to discharge of irrigation water or during periods of heavy rainfall.

To minimize the chance of aquatic impact, the following conditions must be met to allow endosulfan applications:

**DIRECTIONS FOR USE (General)**

IN ADDITION TO SPECIFIC INSTRUCTIONS UNDER DIRECTIONS FOR USE ON THE PRINCIPAL LABELS, THE FOLLOWING CONDITIONS MUST ALSO BE FOLLOWED FOR USE IN CALIFORNIA:

1. Do not apply this product to fields during irrigation or irrigate for 72 hours following an application.
2. Do not apply this product through any type of irrigation system.
3. Do not apply more than 1.5 lbs. a.i./acre/application or more than 3.0 lbs. a.i./acre/season. (Orchard crops and strawberries exempted on rate/acre/application restriction).
4. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark.
5. Do not apply by ground or air within a distance of 300 feet to natural bodies of water including lakes, ponds, streams and estuaries.

**FOR DRIFT CONTROL**

1. All application equipment should be properly maintained and calibrated using appropriate carriers.
2. Do not make aerial or ground applications during temperature inversions.
3. Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 10 mph. Do not discharge at a height of more than 10 feet above the crop or target.
4. For aerial applications a positive shutoff system shall be in place to control flow of liquid to spray nozzles as follows: each individual nozzle shall be equipped with a check valve and the flow controlled by a suckback device or a boom pressure release device; or each nozzle shall be equipped with a positive action valve.
5. For aerial applications, the spray boom must be mounted on the aircraft so as to minimize drift caused by wing tip or rotor vortices. The working boom length must not exceed 75 percent of wing span or rotor diameter.
6. Aircraft boom pressure shall not exceed 40 pounds per square inch.
7. Aircraft nozzles shall be equipped with orifices directed backward parallel to the horizontal axis of the aircraft in flight.
8. Aircraft nozzles shall not be equipped with any device or mechanism which would cause a sheet, cone, fan, or similar type dispersion of the discharged material except as otherwise provided.
9. Fixed wing aircraft and helicopters operating in excess of 60 mph shall be equipped with jet nozzles having an orifice not less than 1/16 inch in diameter (D-4 or larger).
10. Helicopters operating at 60 mph or less shall be equipped with nozzles having an orifice not less than 1/16 inch in diameter. A number 46 or equivalent whirlplate may be used.
11. For ground applications, equipment other than handguns shall be equipped with nozzles having an orifice not less than 1/16 inch in diameter or larger. Boom nozzles shall be operated at a boom pressure not to exceed 30 pounds per square inch.

**READ CONTAINER LABEL FOR COMPLETE USE INSTRUCTIONS**

**ACCEPTABLE SITES FOR ENDOSULFAN APPLICATION**

1. FIELDS, WITH NO DRAINS, HAVING CONTAINMENT THROUGH BERMS (able to hold irrigation water and normal rainwater), LEVEES AND NON-DRAINING CIRCULATIONS SYSTEMS.
2. FIELDS WITH DRAINS, BUT DRAINAGE FLOWS TO SUMP, FARM POND, EVAPORATION POND OR RECIRCULATION SYSTEM WITHOUT DRAINAGE (uncontrollable overflow acceptable, e.g., 10 or 100 year flood). Only grower maintained recirculation systems without drainage are acceptable.
3. FIELDS WITH SURFACE DRAINS THAT NATURALLY FLOW ONTO PASTURELAND OR FALLOW FIELDS THAT HAVE NO CHANNLED DRAINAGE AND ARE AT LEAST 300 FEET LONG.

**UNACCEPTABLE SITES FOR ENDOSULFAN APPLICATION**

1. FIELDS DRAINING DIRECTLY INTO SURFACE WATERS SUCH AS STREAMS, RIVERS, LAKES, PONDS, MARSHES, BAYS, ESTUARIES, OR THE OCEAN.
2. FIELDS DRAINING INTO SURFACE DRAINAGE DITCHES OR CANALS THAT FLOW DIRECTLY INTO SURFACE WATERS SUCH AS STREAMS, RIVERS, LAKES, PONDS, MARSHES, BAYS, ESTUARIES, OR THE OCEAN.
3. ANY AND ALL FIELDS WITH SURFACE DRAINAGE PATTERNS IN THE MOSS LANDING DRAINAGE SYSTEM HAVING MAIN ARTERIES THAT DRAIN INTO THE ELKHORN SLOUGH (northern portion of Monterey County).
4. ANY AND ALL FIELDS WITH DRAINAGE PATTERNS IN THE SAN DIEGO CREEK DRAINAGE SYSTEM HAVING MAIN ARTERIES THAT DRAIN INTO NEWPORT BAY.

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

**THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.**

**Apples (30)**

Insects Controlled	Rate of Application	Method of Application
Aphids (including Apple Aphid, Rosy Apple Aphid, Woolly Apple Aphid) Apple Rust Mite White Apple Leafhopper (first generation)	$\frac{3}{4}$ quart per 100 gallons or a maximum of $3\frac{1}{2}$ quarts per acre	For best control of first generation White Apple Leafhopper, apply when nymphs first appear on leaves. If necessary, prior to petal fall, use 1 quart per 100 gallons to control Apple Aphids and Rosy Apple Aphids.
Do not feed pomace from treated apples to livestock. Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

**Apricots (30), Nectarines (30), Peaches (30)**

Insects Controlled	Rate of Application	Method of Application
Aphids (including Black Cherry Aphid, Black Peach Aphid, Green Peach Aphid, Rusty Plum Aphid) Cattacing insects (Stink Bug type) Peach Silver Mite Peach Twig Borer	$\frac{3}{4}$ quart per 100 gallons or a maximum of $3\frac{1}{2}$ quarts per acre.	Make applications when insects appear or feeding is noticed.
Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

**Barley, Oats, Rye, Wheat**

Insects Controlled	Rate of Application	Method of Application
Army Cutworm	$\frac{3}{4}$ quart per acre	Apply when small larvae are readily found in the field.
Aphids (including Russian Wheat Aphid)	$\frac{3}{4}$ to 1 quart per acre	Make applications when insects appear or feeding is noticed.
Do not apply after heads begin to form. Do not feed treated forage to livestock. Do not make more than 2 applications per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., $1\frac{1}{2}$ quarts) per acre per year.		

**Beans, Succulent and Dry (3) and Southern Field Peas (Succulent type, including Black-eyed Peas, Crowder Peas and Southern Peas) (3)**

Insects Controlled	Rate of Application	Method of Application
Black Bean Aphid Bean Leaf Skeletonizer Cucumber Beetles Flea Beetles Green Stink Bug Leafhoppers Mexican Bean Beetle	$\frac{3}{4}$ to $1\frac{1}{2}$ quarts per acre	Make applications when insects appear or feeding is noticed.
Aphids Western Bean Cutworm Whitefly	$1\frac{1}{2}$ quarts per acre	
Do not feed treated threshings or allow livestock to graze in treated fields. Do not use on Lima Beans that are to be removed from the field for processing. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

**Broccoli (7), Brussels Sprouts (14), Cabbage (7), Cauliflower (14)**

Insects Controlled	Rate of Application	Method of Application
Cabbage Aphid Cabbage Looper Cross-striped Cabbage-worm Diamondback Moth larvae Flea Beetles Harlequin Bug Imported Cabbageworm Stink Bugs	1 to $1\frac{1}{2}$ quarts per acre	Make applications when insects appear or feeding is noticed.
Armyworms Cutworms Whitefly	$1\frac{1}{2}$ quarts per acre	
Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., $2\frac{3}{4}$ quarts) per acre per year.		

**Carrots (7)**

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid	¾ to 1½ quarts per acre	Make application when insects appear or feeding is noticed.
Flea Beetles Whitefly	1½ quarts per acre	

Do not use tops for food or feed.  
Do not make more than one application per year.  
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1½ quarts) per acre per year.

**Celery (4)**

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid Cabbage Looper	¾ to 1½ quarts per acre	Make application when insects appear or feeding is noticed.
Flea Beetles Whitefly	1½ quarts per acre	

Do not make more than one application per year.  
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1½ quarts) per acre per year.

**ly (7)**

Insects Controlled	Rate of Application	Method of Application
Green Peach Aphid	¾ quart per acre	Make applications when insects appear or feeding is noticed.

Do not make more than 2 applications per year.  
Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1½ quarts) per acre per year.

**Cherries (21)**

Insects Controlled	Rate of Application	Method of Application
Black Cherry Aphid Plum Rust (Nursery) Mite	¾ quart per 100 gallons or 2¾ to 3½ quarts per acre	Make applications when insects appear or feeding is noticed.

Do not feed cull fruits to animals or allow livestock to graze in treated yards.  
Do not make more than 2 applications per year.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**Cherry, Peach, Plum Nursery Stock Dip**

Insects Controlled	Rate of Application	Method of Application
Peachtree Borer	2¾ quarts per 40 gallons	Mix thoroughly. Immerse trees so that the roots and crowns are covered well above the grafting bud scar.

Full Personal Protective Equipment (PPE) requirements for applicators also apply to this dipping operation.  
Plant immediately or dry before returning stock to storage.

**Citrus (Non-Bearing Trees and Nursery Stock)**

Insects Controlled	Rate of Application	Method of Application
Citrus Aphid	¾ quart per 100 gallons or a maximum of 3½ quarts per acre	Make applications when insects appear or feeding is noticed.

Do not apply to bearing trees or trees that will bear fruit within twelve months.  
Do not make more than 2 applications per year.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**Collards (21)**

Insects Controlled	Rate of Application	Method of Application
Aphids Cabbage Looper Diamondback Moth larvae Flea Beetles Harlequin Bug Imported Cabbageworm Whitefly	1 to 1½ quarts per acre	Make application when insects appear or feeding is noticed.

Do not make more than one application per year.  
Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.

**Cotton**

Insects Controlled	Rate of Application	Method of Application
Aphids	½ to 1 quart per acre	Make applications when insects appear or feeding is noticed. For control of aphids, thorough coverage is important.
Bollworm Cabbage Looper Cotton Leafperforator Flea-hoppers Lygus Bugs Stink Bugs	1½ to 2 quarts per acre	
Thrips	2 quarts per acre	
Whitefly	1½ quarts per acre	

Applications may be made using ground or aerial application equipment. The higher rate should be used under heavy pest pressure.  
Do not apply after bolls open.  
Do not graze dairy or meat animals in treated fields.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**Cucumbers (2), Melons (2), Pumpkins (2), Summer and Winter Squash (2)**

Insects Controlled	Rate of Application	Method of Application
Aphids Cucumber Beetles Melonworm Pickleworm Rindworm (on watermelons) Squash Beetle Squash Bug Squash Vine Borer Striped Flea Beetle	¾ to 1½ quarts per acre	Make applications when insects appear or feeding is noticed. For Squash Vine Borer control, apply weekly to flower buds, stems, and vines beginning when moths first appear.
Cabbage Looper Omnivorous Leafroller Whitefly	1½ quarts per acre	

Do not make more than 3 applications per year.  
Do not exceed a maximum of 3.0 lb. active ingredient (i.e., 4 quarts) per acre per year.

6098

**Eggplant (1)**

Insects Controlled	Rate of Application	Method of Application
Blister Beetle Colorado Potato Beetle Flea Beetles Green Peach Aphid Green Stink Bug Whitefly	1½ pints per acre	Make application when insects appear or feeding is noticed.
Do not make more than 1 application per year. Do not exceed a maximum of 0.5 lb. active ingredient (i.e., 1½ pints) per acre per year.		

**Grapes (7)**

Insects Controlled	Rate of Application	Method of Application
Grape Leafhopper Grape Phylloxera (leaf form) Rose Chafer	½ quart per 100 gallons or 1½ to 2 quarts per acre	Make applications when insects appear or feeding is noticed.
Do not use on Concord variety as severe plant injury is likely to occur. Ground application is preferred. Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

**Kale (21)**

Insects Controlled	Rate of Application	Method of Application
Cabbage Flea Beetle Harlequin Bug Imported Cabbageworm	1 quart per acre	Make application when insects appear or feeding is noticed.
Do not make more than one application per year. Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.		

**Lettuce (14)**

Insects Controlled	Rate of Application	Method of Application
Cabbage Looper Diamondback Moth larvae Green Peach Aphid Imported Cabbageworm Leafhoppers	1 to 1½ quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1½ quarts per acre	
On Head Lettuce: Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. On Leaf Lettuce: Do not make more than 2 applications per year. Do not feed crop refuse to livestock. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

**Mustard Greens (21)**

Insects Controlled	Rate of Application	Method of Application
Aphids Cabbage Looper Diamondback Moth larvae Fall Armyworm Flea Beetles Harlequin Bug Imported Cabbageworm Whitefly	1 quart per acre	Make application when insects appear or feeding is noticed.
Do not make more than one application per year. Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.		

**Pears (7)**

Insects Controlled	Rate of Application	Method of Application
Pear Psylla	3½ quarts per acre	Apply in a minimum of 10 gallons of water for aerial application or in 300 gallons of water for dilute application. Apply when adults are first observed or nymphs are small and repeat to maintain control.
Pear Rust Mite Conspire Stink Bug (foliar treatment)	½ quart per 100 gallons or 2½ to 3½ quarts per acre	Make applications when insects appear or feeding is noticed. Stink Bugs must be wet by spray to obtain control.
Conspire Stink Bug (soil treatment)	½ quart per 100 gallons; 200 to 400 gallons per acre	Apply to orchard floor and around trees prior to bloom.
Pear Leaf Blister Mite	½ to ¾ quart per 100 gallons	Apply to trees as a post harvest or dormant treatment.
Do not feed cull fruits to animals or allow livestock to graze in treated orchards. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year. Note: Aerial application may not result in satisfactory control and should only be employed if impossible to apply by ground.		

**Pecans**

Insects Controlled	Rate of Application	Method of Application
Black Pecan Aphid Pecan Nut Casebearer Spittlebug	1 quart per 100 gallons	Make applications when insects appear or feeding is noticed. For Casebearer, apply when eggs of first generation appear on the tips of the young nuts. Another application may be required after the second generation of eggs is deposited. For Spittlebug, apply when first leaves are half grown and repeat as required.
Do not apply after shuck split. Do not graze livestock on orchard crops or grasses in treated areas. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Peppers (1)

Insects Controlled	Rate of Application	Method of Application
Flea Beetles Green Peach Aphid Homworms Pepper Maggot	¾ quart per acre	Make applications when insects appear or feeding is noticed.
Do not make more than 2 applications per year. Do not exceed a maximum of 1.0 lb. active ingredient (i.e., 1½ quarts) per acre per year.		

Peppers (4)

Insects Controlled	Rate of Application	Method of Application
Flea Beetles Green Peach Aphid Homworms Pepper Maggot	¾ to 1½ quarts per acre	Make applications when insects appear or feeding is noticed.
Whitefly	1½ quarts per acre	
Do not make more than 2 applications per year. Do not exceed a maximum of 2.0 lbs. active ingredient (i.e., 2¾ quarts) per acre per year.		

Apple (For Fresh Market Only) (7)

Insects Controlled	Rate of Application	Method of Application
Pineapple Fruit Mite	2 to 2¾ quarts per acre	Make applications when insects appear or feeding is noticed. Apply at intervals of 7 to 10 days, if necessary, particularly during the 40 day period of blooming.
Do not feed forage or pineapple by-products to livestock. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Plums (7), Prunes (7)

Insects Controlled	Rate of Application	Method of Application
Aphids (including Hop Aphid, Leafcurl Plum Aphid, Thistle Aphid) Plum Rust (Nursery) Mite	¾ quart per 100 gallons or 2¾ to 3½ quarts per acre	For control of aphids, apply when eggs hatch during pre-bloom or petal fall. Summer applications should be made before leaves curl.
Peach Twig Borer	¾ quart per 100 gallons or 2¾ to 3½ quarts per acre	Make applications when insects appear or feeding is noticed.
Lesser Peachtree Borer Peachtree Borer	¾ to 1 quart per 100 gallons or 2¾ to 3½ quarts per acre.	Best control is obtained with a single application post-harvest after the leaves have dropped. Spray all bark areas from ground level to lower scaffold limbs.
Do not allow livestock to graze on treated orchard crops or grasses in treated areas. Do not make more than 2 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Potatoes (1)

Insects Controlled	Rate of Application	Method of Application
Aphids Armyworms Colorado Potato Beetle Green Stink Bug Leaffooted Bug Plant Bugs Potato Flea Beetles Potato Leafhopper Potato Tuberworm Threelined Potato Beetle	¾ to 1½ quarts per acre	Make applications when insects appear or feeding is noticed.
European Corn Borer Potato Psyllid	1 to 1½ quarts per acre	
False Chinch Bug Whitefly	1½ quarts per acre	
Do not make more than 6 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

Spinach (21)

Insects Controlled	Rate of Application	Method of Application
Crown Mite Flea Beetles Green Peach Aphid Whitefly	1 quart per acre	Make application when insects appear or feeding is noticed.
Do not make more than one application per year. Do not exceed a maximum of 0.75 lb. active ingredient (i.e., 1 quart) per acre per year.		

Strawberries (4)

Insects Controlled	Rate of Application	Method of Application
Meadow Spittlebug Strawberry Aphid Tarnished Plant Bug Whitefly	1½ quarts per acre	Make applications when insects appear or feeding is noticed. Do not reapply within 15 days or more than twice during a 35 day period when fruit is present.
Cyclamen Mite	2¾ quarts per acre in 400 gallons of water	Make applications when insects appear or feeding is noticed. Thoroughly wet the foliage, stem and crown of the plant. For multiple applications, do not apply at intervals less than 35 days when fruit is present.
Do not make more than 3 applications per year. Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.		

8048

**Sweet Corn (Fresh Vegetable Use Only) (1)**

Insects Controlled	Rate of Application	Method of Application
Com Leaf Aphid Whitfly	1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Com Earworm	2 quarts per acre	Apply when silks first appear and continue until they begin to dry. Allow 5 days between applications.

Do not apply to Sweet Corn to be processed.  
Do not feed treated forage or ensilage to livestock or allow livestock to graze in treated fields.  
Do not make more than 3 applications per year.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**Sweet Potatoes (1)**

Insects Controlled	Rate of Application	Method of Application
Sweet Potato Flea Beetle Sweet Potato Weevil	3/4 quart per acre	Make applications when insects appear or feeding is noticed. For flea beetle control, begin applications shortly after transplanting or as soon as flea beetles appear. For Sweet Potato Weevil control, apply in sufficient water for complete coverage. Applications may be made to transplants in the nursery and/or in the field. Repeat applications may be made as necessary.

Do not feed cull potatoes to livestock or allow livestock to graze in treated fields.  
Do not make more than 3 applications per year.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**Tomatoes (Field and Greenhouse) (2)**

Insects Controlled	Rate of Application	Method of Application
Aphids Bristle Beetles Colorado Potato Beetle Flea Beetles Tomato Hornworms Yobworm	5/8 to 1 1/2 quarts per acre	Make applications when insects appear or feeding is noticed.
Cabbage Looper Sink Bugs	1 to 1 1/2 quarts per acre	
Tomato Fruitworm Tomato Russet Mite Whitfly Yellowstriped Armyworm	1 1/2 quarts per acre	
Whitfly	3/4 quart per 100 gallons of water. Use 100 to 200 gallons of spray per acre.	Make applications when insects appear or feeding is noticed. Higher spray volumes may be necessary for thorough coverage when high Whitfly populations exist.

Do not make more than 3 applications per year.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**COMMERCIALLY GROWN ORNAMENTALS**

**Ornamental Plants**

**Ornamentals (Greenhouse, and Out-of-Doors)**

Insects Controlled	Rate of Application	Method of Application
Aphids Cyclamen Mite Rose Chafer Whitfly	3/4 quart per 100 gallons (1 1/2 teaspoons per gallon)	Make applications when insects appear or feeding is noticed.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides.  
On chrysanthemums, best results will be obtained if applied before plants flower.  
Do not apply to "Bonnie Delux", "Fred Shoemith", and "White Knight" chrysanthemums as injury may result.  
Do not use on Birch trees.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**Shade Trees (except Birch), Shrubs**

Insects Controlled	Rate of Application	Method of Application
Aphid	3/4 quart per 100 gallons (1 1/2 teaspoons per gallon)	Make applications when insects appear or feeding is noticed.

Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides.  
Do not exceed a maximum of 3.0 lbs. active ingredient (i.e., 4 quarts) per acre per year.

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Thiodan® is a registered trademark of AgrEvo USA Company  
 FMC —FMC Trademark (3212-11/13/97draft A)