

CODE
20027

SULFUR 50 ZINEB 4 THIODAN 3 DUST

NET WEIGHT
50 lbs.

FUNGICIDE INSECTICIDE


DIRECTIONS FOR USE

This material is designed for use as an undiluted dust for control of diseases and insects indicated on the crops below using a machine capable of thoroughly covering all plant surfaces.

BEANS (String and Snap): Bean rust, Powdery mildew, Leafspot, Anthracnose, Downy mildew, Black bean aphid, Bean leaf skeletonizer, Cucumber beetles, Mexican bean beetle, Green stink bugs - Apply at the rate of 33 pounds per acre when insects or diseases first appear, repeating as necessary. Under severe disease or insect conditions applications should begin earlier. Do not apply to beans within 7 days of harvest. Do not exceed 3 applications per season. Do not feed treated threshings to livestock or allow to graze in treated fields.

PEPPERS: Anthracnose, Powdery mildew, Ripe rot, Cercospora leaf spot, Green peach aphid, Pepper maggot, Flea beetles, Hornworms - Apply at the rate of 35 to 50 lbs. per acre when insects or diseases first appear, repeating as necessary. Do not apply to peppers within 1 day of harvest. Remove residues at time of harvest by washing or by brushing.

SQUASH: Anthracnose, Downy mildew, Powdery mildew, Scab, Aphids, Cucumber beetles, Squash vine borer, Striped flea beetle. Apply at the rate of 30 to 33 pounds per acre when insects or diseases first appear, repeating as necessary. For Squash vine borer, apply weekly to flower buds, stems and vines, beginning when moths first appear. Do not use sulfur on cucumbers, melons or squash unless local experience has shown sulfur to be safe on such plants. Do not apply within 5 days of harvest.



013

E. P. A. Reg. No. 279-2927

ACTIVE INGREDIENTS:

*Sulfur	50.0%
**Zineb	4.0%
***Endosulfan (Hexachlorohexahydro- methano-2,4,3-benzodioxathiepin oxide)...	3.0%
INERT INGREDIENTS:.....	43.0%
	<u>100.0%</u>

- *Sulfur fineness: 93% through 325 mesh
- **Zinc ethylene bisdithiocarbamate
Zinc expressed as metallic ----- 0.94%
- ***Thiodan, U. S. Pat. No. 2,799,685

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

Workers entering treated fields within 24 hours should wear protective clothing.

NOTE TO PHYSICIANS: Endosulfan is a central nervous system stimulant. There is no specific antidote. Barbituric acid derivatives may be used for convulsions. Treat symptomatically.

For Emergency Assistance Call 716-735-3765

Thiodan is a registered trademark of Canadian Hoechst Limited

EGGPLANT: Powdery mildew, Blister beetle, Colorado potato beetle, Green peach aphid, green stink bugs, Whitefly, Flea beetles - Apply at the rate of 20 pounds per acre when insects or diseases first appear, repeating as necessary. Do not apply to eggplant within 1 day of harvest. Remove residues at time of harvest by brushing or washing.

TOMATOES: Tomato fruitworm, Aphids, Colorado potato beetles, Flea beetles, Whitefly, Early blight, Late blight - Use 33 pounds of dust per acre. Do not apply within 5 days of picking fruit. Do not apply on tender young plants as injury may occur.

USAGE CAUTIONS

Applications of dusting or spraying materials containing sulfur should not be made just prior to or during periods of excessively high temperatures.


This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Dispose of wastes by burying in non-crop lands away from water supplies. Bags should be disposed of by burying with wastes or by burning. Keep out of smoke.

Printed in U. S. A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE OF THIS PRODUCT SHALL BE AS SHOWN ON THE ORIGINAL LABEL AND PACKAGING CONTAINING THIS PRODUCT.

WARRANTY: Niagara Chemical Division warrants that the product is as described on the label and that it conforms to the specifications stated on the label. It is sold "AS IS" and is not warranted to perform in any particular manner. EXCEPT AS SO WARRANTED, THE PRODUCT IS NOT A TRADEMARK OF NIAGARA. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGE: Niagara Chemical Division is not liable for damage to property or persons caused by the use of this product in accordance with the label directions. The user shall be responsible for any damage to property or persons caused by the use of this product in any manner other than as directed on the label. Niagara Chemical Division is not liable for any damage to property or persons caused by the use of this product in any manner other than as directed on the label.

EXCUSE FROM NON-CONFORMANCE: NIAGARA CHEMICAL DIVISION is not liable for any damage to property or persons caused by the use of this product in any manner other than as directed on the label. Niagara Chemical Division is not liable for any damage to property or persons caused by the use of this product in any manner other than as directed on the label.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

