

CODE: 3226


NET WEIGHT
51 POUNDS

DIAZINON® 40 WP

DIRECTIONS FOR USE

DIAZINON® 40 WP

INSECTICIDE	
E.P.A. REG. NO. 279-2932 AA	
ACTIVE INGREDIENT:	BY WT.
*O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	40.00%
INERT INGREDIENTS:	60.00%
	100.00%



WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. May be absorbed through skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or household pets. Do not permit children or pets to go on to sprayed grass until spray has completely dried.

USAGE CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.


Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765

*Diazinon is a registered trademark of CIBA-GEIGY Corporation.



Mix with sufficient water (unless otherwise noted) to obtain thorough coverage. Apply when pests first appear and repeat as necessary to maintain control. May be applied by ground or air. Observe re-entry interval between last application and harvest. See label for number in () to cover crops.

ALMONDS (0): Mites, Egg Borers - Use 1 1/2 to 2 1/2 pounds per 100 gallons of water. Parlatoria and San Jose Scales - Mix 1 1/2 pounds with 1 1/2 to 3 gallons of a dormant oil per 100 gallons of water and apply as a dormant spray. May be used to control scales.

APPLES (14), PEARS (14): San Jose Scale, Brown Apricot Scale, Mealybugs, European Red Mite Eggs, Brown Mite Eggs, Pear Leafhopper, Pear Aphid Eggs - Mix 1 1/2 pounds with 2 1/2 gallons of a dormant oil per 100 gallons of water. Apply as a dormant or a dormant spray. Eye-Spotted Bud Moths, Fruit Tree Leafhoppers, Tentative Leafminers, Mistlebees, Codling Moths, Pear (Russet) Apple, and Pear and Walnut Mites - Use 1 1/2 pounds per 100 gallons of water. Make 3 to 5 applications beginning at petal fall to control Pear and Walnut Mites, and Mealybugs, San Jose Scale Crawlers, and Pear Leafhoppers. Use 1 1/2 pounds per 100 gallons of water. Apply early in the season when crawlers first appear. If necessary, make a second application 7 to 10 days later. To control Leafhoppers, apply only, apply 1 1/2 pounds per 100 gallons of water. Make 3 to 5 applications beginning at petal fall and continuing through the growing season.

APRICOTS (10), NECTARINES (20), PEACHES (20): Scales (San Jose, Brown Apricot, and Walnut), and Apricot Mealybugs - Mix 1 1/2 pounds with 1 to 2 gallons of a dormant oil per 100 gallons of water. Apply as a dormant spray. Aphids, Brown Mites, Cover Mites, Spotted Spider Mites, Green Scale Crawlers, and San Jose Scale Crawlers - Mix 1 1/2 pounds with 1 1/2 gallons of foliage spray oil per 100 gallons of water.

APRICOTS (only): Mealybugs - Use 1 1/2 pounds per 100 gallons of water. Apply as a cover spray from petal fall to June.

NECTARINES and PEACHES (only): Peach Twig Borers, and Parlatoria Scales - Use 1 1/2 pounds plus 1 gallon of foliage spray oil per 100 gallons of water. Apply at petal fall. Oriental Fruit Moths - Use 1 1/2 pounds per 100 gallons of water and apply at complete petal fall. Make 2 additional applications at 10 to 15 day intervals. Time the last application to peak adult Oriental Fruit Moths activity, if possible.

PEACHES (only): Peach Twig Borers - Mix 1 1/2 pounds per 100 gallons of water and apply in May and June. Leafhoppers - Apply 1 1/2 pounds per 100 gallons of water in 3 to 5 applications beginning at petal fall and continuing through the growing season. Apply at infestations develop. Write Peach Scale - As a post-harvest spray use 1 1/2 to 2 1/2 pounds per 100 gallons of water timed to coincide with peak crawler and immature scale activity. Do not apply more than twice.

BUSHBERRIES (BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES) (7): Leafhoppers, Thrips, Aphids, Two-Spotted Spider Mites, Raspberry Swainson and Raspberry Fruitworm - Use 1 1/2 pounds per 100 gallons of water applying no more than 2 1/2 gallons of spray per acre, or apply 2 1/2 pounds per acre in sufficient water to cover. To control Raspberry Fruitworm make first application when blossom buds separate and repeat immediately before blossoms open. Dewberry Mites - Mix 2 1/2 pounds in 100 gallons of water. Use as a dormant spray in early spring. Make no more than 2 applications before buds open. Raspberry Crown Borer (Raspberry Root Borer) - Use 5 pounds per 100 to 200 gallons of water and apply as a drench to the crown area and to the lower vines. Apply either in the spring before berries begin to form, or as a post-harvest spray.

CHERRIES (10): Brown Apricot Scales, San Jose Scales, Black Scales, Black Cherry Aphid Eggs, and Brown Mite Eggs - Mix 1 1/2 pounds with 1 to 3 gallons of a dormant oil per 100 gallons of water. Apply as a dormant spray. Eye-Spotted Bud Moths, Fruit Tree Leafhoppers and Leafhoppers - Use 1 1/2 pounds per 100 gallons of water. Begin treatment at petal fall. Do not make more than 5 applications. Black Cherry Aphids, Cherry Fruit Flies - Use 2, 3 to 1 1/2 pounds per 100 gallons of water. Make 3 to 4 applications beginning at petal fall. adult flies begin to emerge and repeat at 10 days intervals. San Jose Scale Crawlers - Use 1 1/2 pounds per 100 gallons of water. Apply during the pre-pink stage or when crawlers first appear. Cherry Post Mites - Use 2, 3 to 1 1/2 pound per 100 gallons of water. Apply as a supplemental post harvest spray.

CITRUS (GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGELOS, TANGELINES, CITRON, KUMQUATS) (21): Citrus Aphids, Cottony Citrus Scale Crawlers, Soft Scale Crawlers, and Fruit Tree Leafhoppers - Use 2, 3 to 1 1/2 pounds in 100 gallons of water (maximum 1000 gallons per acre). Citrus Thrips - Use 1 1/2 pounds in 100 gallons of water (maximum 1000 gallons per acre). Citrus Snow Scales - Use 1 1/2 to 2 1/2 pounds per 100 gallons of water (maximum 1000 gallons per acre) as a post-harvest spray and again as a summer spray. Use the higher dosage rate for heavy infestations. Thorough coverage of trunk, limbs, and twigs is essential for good control. Do not apply during bloom.

FIGS (5): Vinegar Flies and Dried Fruit Beetles - Use 1 1/4 pounds in 100 gallons of water.

FILBERTS (0): Filbert Lestroller, and Aphids - Use 2 1/2 to 5 pounds in 250 to 400 gallons of water per acre.

GRAPES: Pacific Spider Mites, Leafhoppers, Grape Berry Moths, and Grape Leafhoppers - Use 1 1/2 pounds per 100 gallons of water, or apply 2 1/2 pounds per acre in sufficient water for thorough coverage. Drosophila spp. - Use 1/2 pound per 100 gallons of water, or apply a maximum of 1 1/2 pounds per acre in sufficient water for thorough coverage when insects first appear. Make additional applications at weekly intervals as necessary. Do not use more than 100 gallons per acre. Do not apply within 10 days of harvest. If a spreader-sticker is used, do not apply within 18 days of harvest.

OLIVES (75): Olive Scales - Mix 1 to 1 1/4 pounds with 1 1/2 gallons of foliage spray oil per 100 gallons of water. Apply as a cover spray in June or July. Do not apply during bloom to avoid injury to pollinating insects.

PLUMS (10), PRUNES (10): Peach Twig Borers, Mealy Plum Aphid Eggs, (San Jose, and Parlatoria), and Brown Almond Mites - Use 1 1/2 pounds plus 1 to 2 1/2 gallons of a dormant oil per 100 gallons of water. Apply as a dormant spray. For Pear and Walnut Mites, make applications mix 1 1/2 pounds plus 1 gallon of foliage spray oil per 100 gallons of water. Pear and Walnut Mites (Cover Mites), European Red Mites, Aphids, Leaf Curl Plum, Mealy Plum, and Thrips - Use 2, 3 to 1 1/2 pounds per 100 gallons of water.

PECAN (0): Aphids, Mites, Pecan Nut Casebearers, Walnut Caterpillars - Apply 2 1/2 to 7 1/2 pounds per acre as a foliage spray. Do not apply after husks crack.

WALNUTS (0): Aphids, Mites - Use 2 1/2 pounds per acre as a foliage spray. Codling Moths, Scale Crawlers, Walnut Caterpillars - Use 7 1/2 pounds per acre. Do not apply after the husks open.

STRAWBERRIES (5): Aphids, Two-Spotted Spider Mites - Use 1 1/2 pounds in 100 gallons of water per acre. Cyclamen Mites - Use 2 1/2 pounds in 100 gallons of water per acre, directing spray to plant crowns. Plants should be sufficiently irrigated to assure thorough coverage of the foliage. Make additional post-harvest applications if necessary. Strawberry Leafhoppers - Use 1 to 1 1/2 pounds per 100 gallons of water. Apply 100 gallons of spray per acre when leaves show color.

ORNAMENTAL INSECTS: In Ashberry tree, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Dogwood, Elm, Gladiolus, Hawthorn, Holly, Juniper, Lilac, Lorain Apple, Oak, Pear, Plum, Poplar, Rose, Spirea, and Willow - Apply the following recommended rates to control: Aphids, bagworms, Carnation Bud Mites, Carnation Root Mites, Cover Mites, Cotton Mites, European Red Mites, European Fine Spotted Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Citrus Root Weevils, Citrus Vicious Leafhoppers, Privet Mites, Scale Crawlers (Cottony-Cushion, Lecanium, Pine Needle, San Jose Scale, Citrus Thrips), Two-Spotted Mites, Web worms, and Whiteflies - Use 2, 3 ounce per 3 gallons of water or 1 1/2 pounds per 100 gallons of water. Apple and Pear Skeletonizers, Cotoneaster Webworms, Fall Webworms, Hemlock Caterpillars, Oak Loopers, Cabbage Looper, Leafhoppers, Pear Scales, and Tent Caterpillars - Use 2 ounces per 3 gallons of water or 3 1/2 pounds per 100 gallons of water. Ringneck Webworms - Mix 2 ounces in 3 gallons of water for 1 1/2 pounds in 100 gallons of water and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4 to 5 week intervals. Do not use on certain Ferns, Poinsettia, Hibiscus, Papaya, Ficus, and Gardenia, since injury to the plants may occur.

LAWN INSECTS: Lawn Chinch Bugs - Use 2 1/2 to 3 1/2 ounces, or 3 1/2 to 7 1/2 ounces in 25 gallons of water per 1000 square feet. Use the higher rate for target residual control, and in lawns of dense growth with Ants, Armyworms, Clover Mites, Columbia Weevil, Crickets, Cutworms, Orange Wasps, Earwigs, Fruit Flies, Lawn Billbugs, Sod Webworms, Lawn Moths, and Sowbugs.

ACTIVE INGREDIENT:
*0.0...
4...
INERT INGREDIENTS

BY WT.
40.00%
10.00%

WATERBURY

KEEP OUT OF REACH OF CHILDREN

May be used if used with the... spray mist
Avoid...
nation of...
household...
completely dried.

USAGE CAUTIONS

This product is toxic to fish, birds, and...
feeding...
or...
Do not apply when weather conditions...
when...
with...
able...
This product is highly toxic to...
tive...
DANGER: ...
should...

For Emergency Assistance Call
710-730-0765

*Diazinon is registered with the EPA...




Made in U.S.A. by
FMC Corporation
Agricultural Division
P.O. Box 1000
Philadelphia, PA 19106

Waterbury
Diazinon

APRICOTS 10...
PEACHES 10...
PLUMS 10...
PRUNES 10...
WALNUTS 10...
STRAWBERRIES 5...
LAWN INSECTS

PRODUCT BULLETIN



CHLORINOL 4.5% WP

EPA Reg. No. 273-2332

GENERAL DIRECTIONS: Mix with sufficient water (unless otherwise noted) to obtain thorough coverage. Apply when pests first appear and repeat as necessary to maintain control. May be applied by ground or air. Observe days interval between last application and harvest as indicated by number in () following crop.

CRANBERRIES (7): Black-Headed Fireworms, and Sparganothis sulfureana - Use 5 pounds in 400 gallons of water per acre. Cranberry Fruitworms - Use 7½ pounds in 400 gallons of water per acre.

MELONS, CANTALOUPE, CASABAS, CRENSHAW, HONEYDEW, MUSKMELONS, PERSIAN MELONS and hybrids of these; WATERMELONS and hybrids (3); SQUASH (Summer - 7, Winter - 3): Aphids, Leafhoppers, and Mites - Apply 1¼ to 1¾ pounds per acre. Serpentine Leafminers, Thrips - Use 2½ to 3¼ pounds per acre.

PINEAPPLE (0): Pineapple Scales - Use 12½ pounds with 2 gallons of foliage spray oil in 500 gallons of water per acre. Apply as a post-harvest spray to the vegetative slips on the plant. Pineapple Mealybugs - Use 7 pounds per acre. Do not apply within 7 days if forage and refuse is to be fed to livestock.

SUGAR BEETS: Beet Leafhoppers, Dipterous Leafminers, and Aphids - Apply 1 to 1¼ pounds per acre. Grasshoppers - Use 1¼ pounds per acre. Wireworms - broadcast ½ to ¾ pounds per acre just prior to planting, and work immediately 4 to 6 inches into the soil. Tops may be fed to beef and dairy animals.

SUGARCANE (1): Aphids - Use 2/3 to 1¼ pounds per acre. Maggots and grubs - Use 2 to 3 pounds per acre. Wireworms - Apply 10 pounds per acre in a 12-inch band centered over the furrow in sufficient water to obtain thorough coverage. Apply directly to feed pieces in open furrow at covering time.

IMPORTANT

READ AND OBEY ALL DIRECTIONS ON THIS LABEL CAREFULLY. Do not use this product on crops or in a manner not recommended on this label. Do not use this product in a manner not recommended on this label.

For more information, contact your local FMC representative or write to FMC Chemicals, Dept. 100, P.O. Box 100, Salt Lake City, Utah 84110.

FMC Chemicals

PRODUCT BULLETIN

U.S. DEPARTMENT OF AGRICULTURE
BUREAU OF ENTOMOLOGY AND PLANT QUARANTINE

BEANS (POLE, SNAP, LIMA) (17): Black bear, Aphids, Mexican bean beetles, Leafminers - Apply $1\frac{1}{4}$ to $1\frac{3}{4}$ pounds per acre. Cucumber Beetles - Use 10 to 15 ounces in 100 gallons of water per acre. **LIMA BEANS:** Dipterous Leafminers, Leafhoppers, and Mites - Use $1\frac{1}{4}$ pounds per acre. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay.

PEPPERS (5): Aphids, Serpentine Leafminers - Use 2.5 pounds per acre.

POTACES (35): Aphids, Flea beetles, Colorado potato beetles, Dipterous Leafminers - Use 10 to 15 ounces per acre. Banded Cucumber beetles, Leafhoppers, and Southern Armyworms - Use 1 to $1\frac{1}{4}$ pounds per acre.

RADISHES (10), PARSNIPS (10), TURNIPS (17): Aphids, Flea beetles, Dipterous Leafminers - Use $\frac{1}{2}$ to $1\frac{1}{4}$ pounds per acre.

SPINACH (10), BEETS (4): Aphids, Dipterous Leafminers - Apply $\frac{1}{2}$ to $1\frac{1}{4}$ pounds per acre.

SWEET CORN (17): Corn earwigs - Use $\frac{1}{2}$ to 1 ounce per acre when signs first appear. Two or three applications at 7 to 10 day intervals will usually be necessary. Corn forage may be fed to dairy and beef cattle and sheep.

MAYONNAISE (17): Aphids, Dipterous Leafminers - Use 2.5 pounds per acre. Flea beetles, weekly applications are usually required. Banded Cucumber beetles, Fall Armyworm, and Southern Armyworm - Use $1\frac{1}{4}$ to 2 pounds per acre. Mites and Mites are usually required.

IMPORTANT

U.S. DEPARTMENT OF AGRICULTURE

PRODUCT BULLETIN

12A.21.1.1.1
EPA REG. NO. 100-10000



RANGE, PASTURE, AND GRASSLAND INSECTS: BERMO DAGRASS:rips, whiteflies, and Grasshoppers - Use 1 1/4 pounds per acre.

RANGELAND, PASTURE, GRASS FORAGE, DITCH BANKS, ROADSIDES, WASTELAND, NONCROP AREAS, AND BARRIER STRIPS: Grasshoppers - Apply 1 to 1 1/4 pounds per acre in sufficient water or oil to obtain thorough coverage when insects first appear, preferably in the nymphal stage. Dairy and beef cattle and sheep may be grazed immediately following application or may be fed green forage immediately following cutting. If the spray is a water solution do not cut grass for hay for 21 days. If the spray is an oil solution wait 30 days before cutting for hay. Do not spray livestock directly except as directed under Livestock Insects. Do not repeat application for 30 days.

LIVESTOCK INSECTS: Sheep "Ticks" (Keds), and Lice - Using a high gallonage, high pressure spray, mix 2 1/2 pounds in 100 gallons of water and apply at a rate of 1 gallon per animal. As a low gallonage, low pressure spray, mix 1 1/4 pounds in 100 gallons of water. Apply at a rate of 1 quart per animal. In a sprinkler can, mix 1 1/4 ounces in 10 gallons of water. Dip sheep into a small pen and sprinkle 1 quart of mixture over the head, neck, top, and side of each sheep. Agitate mixture frequently. Do not slaughter sheep for 14 days after treatment. Repeat applications.

IMPORTANT


FMC Corporation
P.O. Box 1000
Pittsburgh, Pa. 15224

FMC CORPORATION

PRODUCT BULLETIN

DIAZINON 4-4E

EPA Reg. No. 499-2932



PEANUTS (21): Cutworms - Use 5 pounds per acre in a 2-inch band over the row. Use sufficient water to achieve thorough coverage. Begin application when small larvae appear. Two or three additional applications at 3 week intervals may be necessary. Livestock may be fed peanut hulls and forage 7 days following treatment. Do not feed peanut hay within 21 days following treatment.

SORGHUM (7): Sorghum Midge - Apply $2/3$ to $1\frac{1}{4}$ pounds per acre when 90% of the heads have emerged from the boot. Repeat application within 4 days. Crop may be fed to livestock immediately following treatment. Cutworms (Surface and Subterranean) - Broadcast 5 to 10 pounds per acre just prior to planting. Work into the soil immediately, 2 to 3 inches for Surface Cutworms, and 3 to 6 inches for Subterranean Cutworms.

ALFALFA AND CLOVER: Spotted Alfalfa Aphids, Yellow Clover Aphids, Pea Aphids, Leafhoppers, Plant Bug Nymphs, and Mites - Use 1 to $1\frac{1}{4}$ pounds per acre. Grasshoppers - Apply $1\frac{1}{4}$ pounds per acre. Cutworms (Surface and Subterranean) - Broadcast 5 to 10 pounds per acre just prior to planting. Immediately work 2 to 3 inches into the soil for Surface Cutworms, and 3 to 6 inches for Subterranean Cutworms. Dairy and beef cattle and sheep may be grazed immediately following application, or may be fed green alfalfa or clover immediately following cutting. If treated alfalfa or clover is to be cut for hay, do not cut for 7 days following treatment. Alfalfa Weevil Larvae - For spring control use $2\frac{1}{2}$ to $3\frac{3}{4}$ pounds per acre when feeding damage to growing tips become evident. Repeat application if necessary. If damage is already extensive, cut the hay and make application to the stubble. Do not graze dairy and beef animals or sheep for 2 days. Do not cut for hay for 10 days following treatment. Do not spray livestock directly except as directed under Livestock Insects. Do not treat during bloom to avoid injury to pollinating insects.

IMPORTANT

Read the entire label carefully before using this product. Do not use if the label is missing or if the product is not in the original container.


This product is a restricted use pesticide. Only persons who are licensed under the Federal Insecticide, Fungicide, and Rodenticide Act may apply this product. For more information, see the label.

FMC Corporation

Agricultural Chemical Division
Fresno, California 93717

FMC Chemicals


PRODUCT BULLETIN



The following information is provided for your information and is not intended to constitute an offer or a recommendation of any investment product. The information is provided for your information only and should not be relied upon as a basis for investment decisions. The information is provided for your information only and should not be relied upon as a basis for investment decisions. The information is provided for your information only and should not be relied upon as a basis for investment decisions.

The following information is provided for your information and is not intended to constitute an offer or a recommendation of any investment product. The information is provided for your information only and should not be relied upon as a basis for investment decisions. The information is provided for your information only and should not be relied upon as a basis for investment decisions.

PRODUCT BULLETIN



WASHINGTON, DC
December 10, 1964

CARROT RUST FLIES: Carrots - Apply 5 pounds per acre in a trench to the seed furrow at planting time.

CUTWORMS (SURFACE AND SUBTERRANEAN): Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Cauliflower, Celery, Collards, Cucumbers, Endive (Escarole), Kale, Lettuce, Lima Beans, Melons, Parsley, Peas, Peppers, Potatoes, Snap Beans, Spinach, Summer Squash - Broadcast 5 to 10 pounds per acre just prior to planting. Work into the soil immediately, to 3 inches for Surface Cutworms, and 3 to 6 inches for Subterranean Cutworms.

GARDEN SYMPHYLANS (CENTIPEDES): Cabbage, Carrots, Corn, Lettuce, Peas, Pole Beans, Radishes, Red Beets, Snap Beans, Tomatoes, Turnips - Broadcast 25 pounds per acre just prior to planting. Work into soil 4 to 6 inches immediately. This pre-plant treatment will also control Cutworms (Surface and Subterranean), Mole Crickets, Root Maggots, and Wireworms.


MOLE CRICKETS: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Endive (Escarole), Kale, Lettuce, Melons, Peppers, Radishes, Tomatoes, Turnips - Broadcast 4 pounds per acre when insects are present. Apply 1 month before planting or 1 to 2 days prior to sowing of seed or transplanting, and immediately work into the soil.

WIREWORMS: Broadcast 10 pounds per acre in 5 gallons of water per acre. Spray on top and inside walls of garden beds, and in cracks and crevices, after compost in the garden has been applied. Do not spray on plants. Treat the sides of beds, paths, ditches, and borders after planting, and in cracks and crevices with a trowel. Do not treat plants.

IMPORTANT:

RECEIVED
12/10/64

PRODUCT BULLETIN



REGISTRATION NO. 101-101-101
EPA Reg. No. 101-101-101

ROOT MAGGOTS: Broccoli, brussels Sprouts, Cabbage, and Cauliflower - Broadcast 5 to 7½ pounds per acre and immediately work into the soil 3 to 4 inches before seeding, either in the seedbed or in the field before transplanting. As transplant water (trench) application - Mix 1/3 to 2/3 pound in 10 gallons of transplant water, when 200 to 300 gallons of water are used per acre. May be applied at the rate of 2/3 to 1¼ cupful per plant, either by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Transplant water treatments may result in stand reduction due to plant stress at time of transplanting. Radishes - Mix 2½ pounds in 10 gallons of water. Apply 60 to 100 gallons of spray per acre (1 to 1½ gallons per 1000 linear feet of row) in the seed furrow at planting time.

WIREWORMS: Broccoli, brussels sprouts, Cabbage, cauliflower, Carrots, Cauliflower, Celery, Collards, Cucumbers, Eggplant (egg-plant), Kale, Lettuce, Lima Beans, Muskmelons, Onions, Parsley, Peas, Peppers, Snap beans, Spinach, Summer squash, Sweet corn, Sweet Potatoes, Swiss Chard, Tomatoes, Watermelon, Winter squash - As a broadcast apply 1 to 2 pounds per acre prior to planting and immediately work 4 to 6 inches into the soil. Potatoes - broadcast 1½ to 2 pounds per acre in the seed furrow. Immediately rake or harrow into the soil.

GENERAL WIREWORMS: Potatoes, sweet corn, winter squash, winter squash, broadcast 1 to 2 pounds per acre in the seed furrow and immediately work into the soil.

WIREWORMS: As a broadcast apply 1 to 2 pounds per acre prior to planting and immediately work 4 to 6 inches into the soil.

PRODUCT BULLETIN



APR 10 1973
Pesticides Regulation Division, EPA

BROCCOLI (5), CABBAGE (7), CAULIFLOWER (5): Aphids, Diamondback Moth Larvae, Imported Cabbageworms, Dipterous Leafminers - Use 2 3/4 to 1 1/2 pounds per acre. Applications at 7 to 10 day intervals will usually be required.

BRUSSELS SPROUTS (7): Aphids, Diamondback Moth Larvae, Imported Cabbageworms - Use 1 1/2 pounds per acre. Applications at 7 to 10 day intervals will usually be required.

CARROTS (10): Aphids - Apply 1 1/2 pounds per acre.

CALERY (10): Aphids, Flea Beetles - Use 1 1/2 pounds in 100 gallons of water per acre. Dipterous Leafminers - Apply 2/3 to 1 1/2 pounds per acre in sufficient water to achieve thorough coverage when infestations first appear. Repeat as necessary. Remove tops before shipping or using. Tops should not be used for feed or food.

COLLARDS (10), KALE (10), SWISS CHARD (10), TURNIP TOPS (10): Aphids, Imported Cabbageworms, Diamondback Moth Larvae, Imported Cabbage Bugs, Corn Earworms - Use 2 3/4 to 1 1/2 pounds per acre.

CORNER (10): Aphids, serpentine leafminers, trips - Use 1 1/2 pounds per acre.

ENDIVE (10): Aphids and Dipterous Leafminers - Use 1 1/2 pounds per acre.

GREEN (10): Imported Cabbage Bugs - Apply 1 1/2 pounds per acre.

HEAD (10): Leafminers - Use 1 1/2 pounds per acre. Imported Cabbage Bugs - Use 1 1/2 pounds per acre. Imported Cabbageworms - Use 1 1/2 pounds per acre. Imported Cabbage Bugs - Use 1 1/2 pounds per acre. Imported Cabbageworms - Use 1 1/2 pounds per acre. Imported Cabbage Bugs - Use 1 1/2 pounds per acre. Imported Cabbageworms - Use 1 1/2 pounds per acre.

IMPORTANT

RECEIVED
APR 10 1973