


Code 3226

Net Weight
Pounds

Diazinon[®] 40 WP



Insecticide

For Agricultural or Commercial Use Only.

EPA Reg. No. 279-2932AA EPA Est. 279

Active ingredient	34.4%
0,0-diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	40.0%
Inert ingredients	60.0%
	100.0%

Diazinon[®] is a registered trademark of FMC Corporation.

**KEEP OUT OF REACH OF CHILDREN
WARNING**

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Warning


May be fatal if swallowed. May be absorbed through skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or household pets. Do not permit children to get into sprayed grass until spray has completely dried.

For Emergency Assistance Call 716-735-3765

See Other Panels for Additional Precautionary Information

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia, Pennsylvania 19103



Environmental Hazards

Keep out of reach of fish, birds and other wildlife. Keep out of reach of mammals and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures.

Mix with sufficient water (unless otherwise noted) to obtain thorough coverage. Apply when pests first appear and repeat as necessary to maintain control. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in following crop.

Alfalfa, Clover: Spotted Alfalfa Aphids, Yellow Clover Aphids, Pea Aphids, Leafhoppers, Plant Bug Nymphs, Mites— Use 1 to 1 1/4 pounds per acre. Grasshoppers— Apply 1 1/4 pounds per acre. Cutworms (Surface and Subterranean)— Broadcast 5 to 10 pounds per acre just prior to cutting. Immediately work 2 to 3 inches into the soil for Surface Cutworms and 3 to 6 inches for Subterranean Cutworms. Dairy and beef cattle and sheep may be grazed immediately following application or may be fed green alfalfa or clover immediately following cutting. If treated alfalfa or clover is to be cut for hay, do not cut for 7 days following treatment. Alfalfa Weevil Larvae— For spring control apply 1 to 2 1/2 pounds per acre when feeding damage to growing tips is observed. Repeat application if necessary if damage is already extensive at the hay and make application to the stubble. Do not graze or feed beef animals or sheep for 2 days. Do not cut for hay for 7 days following treatment. Do not spray livestock directly except for control of biting insects. Do not treat during bloom to avoid pollinating insects.

Almonds (0): Mites, Twig Borers— See label for application rates per 100 gallons of water. Olive and San Jose Scale— Mix 1 1/2 to 2 1/4 pounds with 100 to 200 gallons of a dormant oil per 100 gallons of water and apply as a dormant spray. May not be fed to livestock.

Mustard (10)

Onions (10)

Peas (10)

Parsley

Peppers (5)

Pineapple (0)

Plums (10), Prunes (10)

Potatoes (35)

Radishes (10), Turnips (10)

Red Beets (14)

Shrimp

Strawberries

Turnip tops (10)

Watermelons


Apples

Oranges

Pears

Plums

Prunes



Peas (10)

Peppers (5)

Pineapple (0)

Plums (10), Prunes (10)

Potatoes (35)

Radishes (10), Turnips (10)

Red Beets (14)

Shrimp

Strawberries

Turnip tops (10)

Watermelons

Apples

Oranges

Pears

Plums

Prunes

When used on grapes, this product should be applied to the foliage with a spray nozzle. For use on other crops, apply to the soil surface with a broadcast or banding application. Broadcast applications should be made at the time of planting. Work to the soil surface with a hoe or similar implement immediately after application.

Spinach (10) Aphids, Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre immediately after planting. For control of Spider Mites and Leafminers, apply to the soil surface with a broadcast or banding application immediately after planting.

Sugar Beets (0) Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement.

Sugarcane (1) Aphids, Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement.

Strawberries (5) Aphids, Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement. Use 2 to 3 quarts in 100 gallons of water per acre for direct spraying of plants. Weeds should be sufficiently agitated to cause the spray to adhere to the foliage. Make additional post-harvest applications if necessary. Strawberry Leafrollers. Use 1 to 2 pounds per 100 gallons of water. Apply 200 gallons of spray per acre when blossoms show.

Sweet Corn (0) Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement.

Sweet Potatoes (0) Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement.

Swiss Chard (12) Aphids, Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement.

Tomatoes (1) Aphids, Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement. Use 1 to 2 quarts per acre for direct spraying of plants. Weeds should be sufficiently agitated to cause the spray to adhere to the foliage. Garden Symphyla, Centipedes, and Leafminers should be controlled just prior to planting. Immediately work to the soil surface with a hoe or similar implement. Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement.

Walnuts (0) Aphids, Flea Beetles, Leafminers, Spider Mites, and Wireworms. Broadcast 7 to 10 pounds per acre at the time of planting and work immediately to the soil surface with a hoe or similar implement.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use This product is sold as a general use herbicide. It is not intended for use on food crops.

Warranty Manufacturer warrants that the product is free from defects in material and workmanship under normal use and conditions. The user assumes all risk of injury or damage to property or persons from the use of this product.

Directions and Recommendations Follow directions carefully. Timing of application should be based on weather and crop conditions. Mixture with other herbicides should be made as recommended and other influencing factors should be considered. For information on the use of this product, contact the manufacturer.

Use of Product Manufacturer's recommendations for the use of this product are based on tests believed to be reliable. The use of this product is subject to the terms of the manufacturer's guarantee expressed on the label. The user assumes all risk of injury or damage to property or persons from the use of this product.

Damages Manufacturer's liability for damages for breach of warranty is limited to the purchase price of the product and shall not include incidental or consequential damages.

