

[279-2940]

DIRECTIONS

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Use windows and doors and shut off ventilating systems. Use a non-thermal fogger or sprayer or a blower to deliver an aerosol spray (no droplets over 50 microns in diameter and not less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary.

The above described application will kill accessible, exposed stages of flies, including fruit flies, mosquitoes, silver fish, flying moths, gnats, crickets, roaches, spiders, scorpions, wasps, hornets, clover mites, cheese mites, cheese skippers, and these common stored product pests: granary weevils, rice weevils, confused flour beetles, saw toothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, Tobacco moths, meal worms, and grain mites.

IN MUSHROOM PRODUCTION AND PROCESSING: To control adult Phorids, Sciards, and Cecidomyiids in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off. In packing and canning plants, use at a rate of 1 ounce to 2000 cubic feet of space.

CAUTION: Harmful if swallowed. Avoid contamination of feed and food-stuffs.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on the label. Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

384

Printed in U.S.A.

AQUEOUS FOOD PLANT PYRENONE[®] FOGGING INSECTICIDE

EPA Reg. No. 79-2940

A Concentrated PYRENONE[®] Micro Emulsion

- Clear
- Stable
- Leaves No Oily or Objectionable Residue

Designed for Use in Edible Product Areas of Official Establishments Operating Under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs, and in Other Food Processing Plants.

Apply with Conventional Mechanical or Compressed Air Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

ACTIVE INGREDIENTS:

Pyrethrins	0.5%
*Piperonyl Butoxide, technical	5.0%
Petroleum distillate	2.0%
INERT INGREDIENTS:	92.5%
	100.0%

*Equivalent to 4.0% (butylcarbityl(6-propylpiperonyl) ether and to 1.0% related compounds
PYRENONE - FMC trademark


CAUTION

Keep Out Of Reach Of Children

See rear panel for additional precautions

Net Contents 1 Gallon

MANUFACTURED IN U.S.A. BY
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
MIDDLEPORT, N. Y. 14105



279-2940

