

Code: 30318

[279-2956]

MANUFACTURED:

SULFUR 25 DIBROM[®] 4 SEVIN[®] 10 DUST

DIRECTIONS FOR USE

Apply when insects or disease first appear and repeat as necessary. For insect or disease control beyond the time limitations of this product use a suitable material. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in () following the crop.

BEANS (4): Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Leafhoppers — Use 10 pounds per acre. Corn Earworm, Stink Bugs, Tarnished Plant Bug — Use 15 pounds per acre. **California:** Lygus Bug, Lima Bean Pod Borer — Use 20 pounds per acre. Sulfur may injure certain varieties; check tolerance to sulfur prior to use.

BROCCOLI (3), BRUSSELS SPROUTS (3), CABBAGE (3), CAULIFLOWER (3): Flea Beetles, Harlequin Bug — Use 10 pounds per acre. Armyworm, Corn Earworm, Cabbage Looper (small) — Use 20 pounds per acre. For control of Small Cabbage Loopers, after edible parts start to form, apply at 5 to 7 day intervals.

COLLARDS (14), KALE (14), TURNIPS (Roots - 3, Tops - 14): Flea Beetle, Harlequin Bug — Use 10 pounds per acre. Leafhoppers, Armyworm, Corn Earworm, Tarnished Plant Bug, Stink Bug, Cabbage Looper (small) — Use 20 pounds per acre. For control of Small Cabbage Loopers, after edible parts start to form, apply at 5 to 7 day intervals.

COTTON (4): Early Season: Thrips, Flea Beetle, Cotton Fleahopper, Cotton Leafworm, Striped Blister Beetle — Use 10 pounds per acre. Mid to Late Season: Boll Weevil, Bollworm, Fall Armyworm, Cotton Leafperforator, Leafrollers, Leafhoppers, Tarnished Plant Bug, Lygus Bug, Aphids (suppression) — Use 20 pounds per acre. Pink Bollworm, Stink Bug, Saltmarsh Caterpillar — Use 25 pounds per acre. Grasshoppers — Use 15 pounds per acre.

GRAPES (4): Powdery Mildew, Grape Leafhopper, Grape Leafroller, Grape Leaf-skeletonizer, Thrips, Drosophila — Use 20 pounds per acre. Ground application preferred. Do not apply to Concord, Italia, or other labrusca varieties as fruit injury may occur.

LETTUCE (Head - 3, Leaf - 14), SPINACH (14), SWISS CHARD (14): Flea Beetle, Harlequin Bug, Leafhoppers — Use 10 pounds per acre. Armyworms, Corn Earworm, Tarnished Plant Bug, Stink Bug, Cabbage Looper (small) — Use 20 pounds per acre. For control of Small Cabbage Loopers, after edible parts start to form, apply at 5 to 7 day intervals. Apply at 5 to 7 day intervals for control of Corn Earworm on Lettuce after heads start to form.

PEACHES (4), (Western States): Aphids, Fruit Fly, Red-Spider Mites, Lygus Bug, Thrips, Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth — Use 50 to 60 pounds per acre. For Codling Moth, apply at petal fall and every 10 to 14 days thereafter until control is achieved.

PEPPERS (1): Colorado Potato Beetle, Flea Beetle, Leafhoppers — Use 10 pounds per acre. Fall Armyworm, Corn Earworm, Tomato Hornworm, Tarnished Plant Bugs, Stink Bugs, Aphids, Leafminer — Use 20 pounds per acre.

FUNGICIDE-INSECTICIDE

EPA REG. NO. 279-2956 AA

ACTIVE INGREDIENTS:	BY WT.
*Carbaryl- 1-naphthyl N-methylcarbamate	10.00%
**Naied- 1, 2-dibromo- 2, 2-dichloroethyl dimethyl phosphate	4.00%
Sulfur	25.00%
INERT INGREDIENTS	61.00%
	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. May cause irritation of eyes, nose and throat. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

STRAWBERRIES (4): Powdery Mildew, Mealy-berry Weevil — Use 20 pounds per acre.

TOMATOES (1): Colorado Potato Beetle, Fall Armyworm, Tomato Fruitworm, Stink Bug, Tomato Russet Mite, Fruit Fly, etc. — Use 20 pounds per acre.

ENVIRONMENTAL AND

This product is toxic to fish, birds, and areas may be killed. Keep out of lakes, runoff is likely to occur. Do not apply to areas treated. Do not contaminate water wastes. Apply this product only as specified. This product is highly toxic to bees exposed. Protective information may be obtained from Extension Service.


DISPOSE OF WASTES by burying in non-empty bags should be buried with waste.

Some phytotoxicity may occur on tender foliage of several days duration following application.

Applications of materials containing sulfur at periods of excessively high temperatures or

May corrode metal dust equipment; clean equipment. STORE IN COOL, DRY PLACE.

For Emergency
716-73



DIBROM is a registered trademark of Chas. I. D. 297,862.

Trademark of Union Carbide for the active

[279-2956]

Net Weight
50 Pounds

SULFUR 25 DIBROM[®] 4 SEVIN[®] 10 DUST

GUARANTEE EXPIRES:

FOR USE

and repeat as necessary. For insect or
this product use a suitable material
interval between last application and
the crop.

rs, Flea Beetles, Leafhoppers — Use 10
Tarnished Plant Bug — Use 15 pounds per
— Use 20 pounds per acre. Sulfur may
prior to use.

AGE (3), CAULIFLOWER (3): Flea Beetles,
worm, Corn Earworm, Cabbage Looper
of Small Cabbage Loopers, after edible
parts

Tops (14): Flea Beetle, Harlequin Bug
worm, Corn Earworm, Tarnished Plant
— 20 pounds per acre. For control of
to form, apply at 5 to 7 day intervals

Cotton Fleahopper, Cotton Leafworm,
Mid to Late Season: Boll Weevil, Boll-
Leafrollers, Leafhoppers, Tarnished
— 20 pounds per acre. Pink Bollworm
— 15 pounds per acre. Grasshoppers — Use 15

opper, Grape Leafhopper, Grape Leaf
— 20 pounds per acre. Ground application
other labrusca varieties as fruit injury

ISS CHARD (14): Flea Beetle, Harlequin
armyworms, Corn Earworm, Tarnished
— Use 20 pounds per acre. For control
start to form, apply at 5 to 7 day
of Corn Earworm on Lettuce after

Red-Spider Mites, Lygus Bug, Thrips,
Cicada, Codling Moth, Tussock Moth
th apply at petal fall and every 10 to

Leafhoppers — Use 10 pounds per
worm, Tarnished Plant Bugs, Stink
acre.

FUNGICIDE-INSECTICIDE

EPA REG NO. 279-2956 AA

ACTIVE INGREDIENTS:	BY WT.
*Carbaryl: 1 naphthyl N-methylcarbamate	10.00%
**Naled: 1, 2-dibromo-2, 2-dichloroethyl dimethyl phosphate	4.00%
Sulfur	25.00%
INERT INGREDIENTS	61.00%
	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. May cause ir-
throat. Avoid breathing of dust. Do not take int-
be harmful. Avoid contact with skin. Wash hands and face before eating.
Take shower or bath after work. Wear regular long sleeved work clothing.
Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND
WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase
inhibitor. Atropine is antidotal.

STRAWBERRIES (4): Powdery Mildew, Meadow Spittlebug, Strawberry Leafroller, Straw-
berry Weevil — Use 20 pounds per acre

TOMATOES (1): Colorado Potato Beetle, Flea Beetle, Leafhoppers — Use 10 pounds
per acre. Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug,
Stink Bug, Tomato Russet Mite, Fruit Fly, Twospotted Spider Mite, Aphid, Leafminer —
Use 20 pounds per acre

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated
areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where
runoff is likely to occur. Do not apply when weather conditions favor drift from
areas treated. Do not contaminate water by cleaning of equipment or disposal of
wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on
crops. Protective information may be obtained from your Cooperative Agricultural
Extension Service.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies.
Empty bags should be buried with wastes or burned. Keep out of smoke.


Some phytotoxicity may occur on tender foliage in presence of rain or high humidity
of several days duration following application.

Applications of materials containing sulfur should not be made just prior to or during
periods of excessively high temperatures or within 10 days to 2 weeks of an oil spray.

May corrode metal dust equipment; clean equipment after use.

STORE IN COOL, DRY PLACE.

For Emergency Assistance Call
716-735-3765



DIBROM is a registered trademark of Chevron Chemical Company. U.S. Patent No.
2,971,882.

Trademark of Union Carbide for the active ingredient carbaryl.