



279-2972

DIRECTIONS FOR USE

Hold container upright, nozzle away from you and press actuator button spraying as directed.

To Treat Animals: To protect cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

In Animal Quarters & Milk Rooms: For rapid control of House-flies, Mosquitoes, Gnats, Wasps and Flying Moths close windows and doors and spray at the rate of one second per 1000 cubic feet of space. For best results keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

When using this product in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

Remove pets, birds and cover fish aquariums before spraying.

Flies, Mosquitoes, Gnats, Wasps, and Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

Roaches, Water Bugs, Silverfish, Crickets, Spiders, Centipedes, and Saw Bugs: Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moth, Angoumois Grain Moth: Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep areas closed for 15 minutes following treatment. Repeat application weekly or as needed.

Exposed stages of **Confused Flour Beetle, Saw Toothed Grain Beetle, Red Flour Beetle, Rice Weevil, Granary Weevil, Grain Mites:** Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Repeat treatment as necessary.

Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, and floors. Repeat treatment as necessary.

Fleas: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Pyrenone[®] Dairy and Industrial Aerosol

EPA Reg. No. 279-2972

Contains PYRENONE[®]

Industrial Strength - Low Dosage Required

For Use on Cattle, in Animal Quarters, and Milk Rooms.

KILLS AND REPELS Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats.

For use in Schools, Hospitals, Institutions, Dairies, Supermarkets, Hotels, Restaurants, Drive-Ins, Motels, Drug Stores, Food Plants, and Animal Quarters.

Active Ingredients:

Pyrethrins	0.5%
Piperonyl Butoxide, Technical*	5.0%
Petroleum Distillate	14.5%

Inert Ingredients: 80.0%

* Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and to 1.0% of related compounds.

Pyrenone - FMC Trademark

CAUTION Keep Out Of Reach Of Children

CAUTION: Do not apply over uncovered milk or food products. Any exposed utensils or food handling surfaces should be thoroughly cleaned before use. Do not use in areas where fish or reptiles may be exposed to the spray.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F. may cause bursting. Never throw container into fire or incinerator.

NET WEIGHT _____

MANUFACTURED IN U.S.A. BY
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
MIDDLEPORT, N. Y. 14105
084

