

279-2976J

## DIRECTIONS

## SHAKE BEFORE USING AND AT INTERVALS DURING PROLONGED USE

Hold container upright with nozzle away from you. Press valve down and spray as directed. To avoid possible staining of fabrics or surfaces, do not spray at close range.

**Flies, Mosquitoes, Gnats, Wasps, and Fleas:** Mix 1 oz. of this product with 1 gal. of water and spray at the rate of 1 second per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

**Roaches, Waterbugs, Silverfish, Crickets, Spiders, Centipedes, and Sow Bugs:** Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Repeat treatment as necessary.

**Ants:** Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

**Exposed Stages of Indian Meal Moth, Angoumois Grain Moth:** Close doors and windows, direct spray into all parts of room, especially around stored food product containers, pallets, and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

**Exposed Stages of Flour Beetles, Weevils, Grain Mites:** Spray into cracks and crevices, processing and handling equipment, around curtains or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat applications as necessary. Infested stored food products in the home should be disposed of.

**Carpet Beetles:** Spray directly on beetles, floors, baseboards, shelves, and carpet. Repeat treatment as necessary.

**Bedbugs:** Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, molding, and floors. Allow treated fabric to dry thoroughly before using. Repeat treatment as necessary.

**Fleas:** Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. Use only a recommended pet spray on pet.

**To Treat Animals:** To protect beef and dairy cattle from attacks of stable flies, horse flies, deer flies, house flies, horn flies, mosquitoes and gnats, spray about 1 second on each side being careful to spray back, withers and forelegs thoroughly. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

**In Animal Quarters and Milk Rooms:** For rapid control of Houseflies, Mosquitoes, Gnats, Wasps and Small Flying Moths close windows and doors and spray at the rate of one second per 1000 cubic feet of space. For best results keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

For the Control of the Foliage Garden Insects:

Spray with slow sweeping motion, not closer than 15 inches from plants. Avoid wetting plants. Spray plants only when air is still. For single plants, spray in short bursts. Repeat as necessary for control. Be cautious about wetting tender foliage, young plants, and new growth. Do not spray plants when temperatures exceed 90°F. Contact insect directly whenever possible. Repeat application at 7 to 10 day intervals or after heavy rains.

|                             |                                                                                      |
|-----------------------------|--------------------------------------------------------------------------------------|
| Roses                       | Aphids, Rose Chafer, Japanese Beetle, Rose Slug, Tent Caterpillar, Scales            |
| Asters                      | Leafhoppers, Tarnish Plant Bug, Lace Bugs, Thrips (exposed), Mealy Bugs              |
| Dahlias                     | Aphids, Red Spider Mites, Plant Bug                                                  |
| African Violets             | Broad Mites, Citrus Mealybug                                                         |
| Azaleas                     | Azalea Lace Bug, White Fly, Leaf Miner                                               |
| Eucalyptus                  | Eucalyptus Scale, Aphids, Thrips (exposed)                                           |
| Hydrangea                   | Aphids, Red Spider Mites, Hydrangea Leafhopper                                       |
| Juniper                     | Juniper Scale, Red Spider Mite, Bagworm                                              |
| Lilac                       | Leaf Miner, Leaf Roller, Scales                                                      |
| Tomatoes                    | Flea Beetle, Hornworm, Colorado Potato Beetle, Stink Bug, Green Peach Aphid          |
| Peppers, Eggplant, Potatoes | Colorado Potato Beetle, Leafhoppers, Blister Beetle, Green Peach Aphid, Flea Beetles |
| Cucumbers, Squash, Melons   | Striped & Spotted Cucumber Beetle                                                    |
| Beans                       | Mexican Bean Beetle, White Fly, 12-spotted Beetle                                    |
| Sweet Corn                  | Japanese Beetle                                                                      |



FMC Corporation  
Agricultural Chemical Division  
Middleport, New York 14105  
045

PYRENONE (P)

## Multi-Purpose

## Insecticide

Contains PYRENONE

Professional Strength - Low Dosage Required

For use in Schools, Hospitals, Institutions, Barns, Supermarkets, Hotels, Restaurants, Driveshops, Motels, Drive-Ins, Food Plants, Animal Quarters, and Milk Rooms.

Provides rapid control of Flies, Mosquitoes, Gnats, Wasps, Flying Moths (Millers) and accessible stages of Roaches, Apts, Water Bugs, Silverfish, Flour Beetles, Weevils, Grain Mites, and Spiders.

**KILLS AND REPELS Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats**

**Controls Garden Insects On Ornamental and Vegetable Plants.**

Acceptable for Use To Day of Harvest on Edible Plants.

EPA Reg. No. 279-2976

## ACTIVE INGREDIENTS:

Pyrethrins, 100% extract, 0.5%

\*Piperonyl Butoxide, Technical, 5.0%

Petroleum Hydrocarbons, 1.3%

INERT INGREDIENTS, 90.2%

\*Equivalent to 4.0% of dimethylallyl isopropyl piperonyl ether and 1.0% of related compounds.

PYRENONE - FMC trademark

## CAUTION

**Keep Out Of Reach Of Children**


**CAUTION:** Avoid breathing of vapor or contact with eyes or skin. Vacate premises after treatment and ventilate before reoccupying.

Avoid contamination of food and foodstuffs. When using this product on food processing plants, food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

Do not apply over uncovered milk or food products. Any exposed animals or food handling surfaces should be thoroughly cleaned before use. Do not use in areas where fish or reptiles may be exposed to the spray. Remove pets, birds and cover fish aquariums before spraying.

**WARNING:** Contents under pressure. Do not puncture. Do not use or store near heat or open flames. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

NET WEIGHT — 100 LBS.



ACCEPTED  
JUL 15 1977  
279-2976J