



Code 3510

ACCEPTED
Pesticide Labels
EPA-700-10-010-01

APR 12 1983

279-3014

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Pounce 3.2 EC

Insecticide

EPA Reg. No. 279-3014

EPA Est., 279-FL-1

Active Ingredients:

*Permethrin**	38.4%
Xylene range aromatic solvent	50.2%
Inert Ingredients:	11.4%
	100.0%

**3-Phenoxyphenyl methyl (±) *cis-trans* 3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate
***cis trans* ratio: Max. 55% (±) *cis* and min. 45% (±) *trans*

Contains 3.2 lbs permethrin per gallon.

U.S. Patent No. 4,024,163

KEEP OUT OF REACH OF CHILDREN

CAUTION

First Aid

If in eye, flush eyes with water for 15 minutes. Call a physician.
If swallowed, do not induce vomiting. Call a physician.
If on skin, remove contaminated clothing. Wash with plenty of soap and water. Get medical attention if irritation persists.
Note to Physician: Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages by aspiration of the solvent.

For Emergency Assistance Call 716-735-3765.

See Other Panels for Additional Precautionary Information.

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Remove and wash clothing before reuse.

Environmental Hazards

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service.

This product is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Physical or Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not store below 10° F (-12° C). Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or local procedures. To dispose of container, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures. Do not cut or weld this container.

This label must be in the possession of the user at the time of application.

274

Potatoes

To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Cutworms, European Corn Borer, Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm, Aster Leafhopper and Tarnished Plant Bug. Use Pounce 3.2 EC at a rate of 4-8 ounces (0.1 to 0.2 pounds active) per acre. Apply when insects appear or feeding is noticed and repeat as necessary to maintain control. Use higher recommended dosage rates as pest populations increase. Apply in a minimum of 3 gallons of finished spray per acre by air or 20 gallons of finished spray per acre using ground equipment. Do not make more than 12 applications per season. Do not apply within 7 days of harvest. Do not graze or feed potato forage. Do not plant rotational crops within 60 days of the last application.

Spinach - To control Beet Armyworm, Corn Earworm, Cabbage
Looper, Cutworms - Use Pounce 3.2 EC at a rate of 2-4 oz.
(0.05-0.1 lb. active) per acre. Apply when insects appear.
Make applications with a minimum of 20 gallons of finished
spray per acre with ground equipment. Do not make more than
7 applications per season. Do not apply within 7 days of
harvest.

d316A3
dd30

Broccoli, Brussels Sprouts, Cauliflower

To control Beet Armyworm, Cabbage Looper, imported Cabbage-worm and Diamondback Moth—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05-0.1 pounds active) per acre. Apply when insects appear and repeat as required to maintain control. Apply in a minimum of 3 gallons of finished spray per acre by aircraft and 20 gallons per acre by ground equipment. Do not make more than 3 applications per season. Pounce 3.2 EC may be applied up to the day of harvest. Do not plant rotational crops within 60 days of last application. Do not feed trimmings from treated fields to livestock.

Cabbage

To control Beet Armyworm, Cabbage Looper, Diamondback Moth, and imported Cabbage-worm—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05-0.1 pounds active) per acre. Apply when insects appear and repeat as required to maintain control. Apply in a minimum of 1 gallon of finished spray per acre by aircraft and 20 gallons per acre with ground equipment. Do not make more than 10 applications per season. Do not apply within 1 day of harvest. Do not plant rotational crops within 60 days of last application. Do not feed cabbage trimmings from treated fields to livestock.

Cotton

To control beet armyworm (California and Arizona only), bollworm, cabbage looper, cotton fleahopper, cotton leaf perforator, Lygus bugs, pink bollworm, tarnished plant bug and tobacco budworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Use higher dosage rates as population pressure increases. To control boll weevil, alternate Pounce 3.2 EC at 4 ounces (0.1 lb. active) per acre with Pounce 3.2 EC at 2 ounces (0.05 lb. active) per acre every 3-4 days until control is achieved.

Apply the above rates in a minimum of one gallon of water per acre by aircraft and 5 gallons of water per acre with ground equipment. Pounce 3.2 EC may also be applied using refined non-volatile vegetable oil for control of the above named insects. Pounce 3.2 EC should be diluted with oil and applied in a minimum of one quart total volume/acre using equipment calibrated to give adequate coverage. To control bollworm and tobacco budworm use 2 to 8 ounces (0.05 to 0.2 lb active) per acre. The 2 ounce rate per acre should be used only when insect infestation levels are light and the cotton is closely scouted before and after application.

Do not make more than 15 applications per season. Do not apply within 14 days of harvest. Do not graze or feed cotton forage. Do not plant rotational crops within 60 days of last application.

Cotton Tank-Mix

Pounce 3.2 EC Plus Tank Mixtures (all cotton growing states). Mix Pounce 3.2 EC at 4.0-8.0 ounces/acre (0.1-0.2 lb. a/acre) with other insecticides recommended on this label for control of cotton insect pests. Pounce 3.2 EC has been found to be effective and compatible in tank mixtures with the following products.

Methyl Parathion
EPN Plus Methyl Parathion
Methomyl
Guthion
Azinphos-Methyl
Malathion
Chlordimeform

When using Pounce 3.2 EC in tank mixtures, observe all restrictions and precautions which appear on the labels of these products. Apply tank mixtures immediately after mixing.

Celery

To control Leafminer, Cutworms and Southern Armyworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1-0.2 pounds active) per acre on a 3 to 5 day schedule, or as needed to maintain control. Apply in a minimum of 5 gallons of finished spray per acre by aircraft and 10 gallons per acre by ground equipment. Do not make more than 10 applications per season in California. Do not apply within 3 days of harvest. Do not graze celery forage nor feed trimmings from treated fields to livestock.

Head Lettuce

To control Alfalfa Looper and Cabbage Looper—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05-0.2 pounds active) per acre. To control Beet Armyworm, Corn Earworm and Tobacco Budworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1-0.2 pounds active) per acre. Apply when insects appear and repeat as required to maintain control. Use higher dosage rates as population pressure increases. Apply a minimum of 5 gallons of finished spray per acre by aircraft and 15 gallons per acre with ground equipment. Do not make more than 15 applications per season. Do not plant rotational crops within 60 days of last application.

Pears

To control Pear Psylla—Apply Pounce 3.2 EC at a rate of 8 to 16 oz. (0.2-0.4 pounds active) per acre. Pounce 3.2 EC may be combined with 2 to 8 gallons of spray oil per acre. Apply during the dormant through delayed dormant growth periods only. Apply in 3 to 20 gallons of finished spray per acre by aircraft and 25 to 400 gallons per acre by ground equipment. Do not make more than two applications per season. Do not graze orchard floor.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Pounce and FMC are registered trademarks of FMC Corporation.

(3510-PA-10/82)

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia, Penna. 19103

Attention: Dr. Mark Galley

SUBJECT: Pounce 3.2 EC/Spinach
E.P.A. Registration Number 279-3014
Your letter dated March 16, 1983

The amendment referred to above, submitted in connection with registration under FIFRA sec. 3 (c)(5) is acceptable.

Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6 (e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for records.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

RES: [illegible]

310
17
87369