

300 + 231971
1714

SEP 22 1988

Ms. Susan E. Burkart
Senior Registration Specialist
FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia, PA 19103

Dear Ms. Burkart:

Subject: Correct Dilution Rate and Draft Label
Amendment to Add Ants to Almonds and Pistachios
Pounce 3.2 EC Insecticide
EPA Registration No. 279-3014
Correct Dilution Rate and Preemergent Use Directions
EPA Registration No. 279-3051
Pounce 3.2 Miscible Insecticide
EPA File Symbol 279-GNTG
Draft Amended Label for Pine Seed Orchards
Your Submissions Dated February 16 and 17, 1988

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling.

- 1. For EPA Registration Nos. 279-3014 and 279-3051, in the Environmental Hazards statement pertaining to water contamination, revise the statement to read:

Do not contaminate water when disposing of equipment washwaters.

- 2. Since Pounce 3.2 Miscible Insecticide - EPA File Symbol 279-GNTG is not a registered product, we will not act on this submission at this time.

11165:I/C:Heyward:LR-7:KENCO:3/10/88:4/20/88:CB:EK:VO:CB

CONCURRENCES

SYMBOL	ORIGINATOR							
SURNAME								
DATE								

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of each product's labeling is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosures

Code 3510

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Net Contents

Pounce 3.2 EC

Insecticide

EPA REG. NO. 279-3014

EPA Est., 279-NJ-2

ACTIVE INGREDIENTS:

*Permethrin** 38.4%

INERT INGREDIENTS:*** 61.6%

100.0%

*(3-Phenoxyphenyl)methyl (±) *cis-trans* 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate
***cis/trans* ratio. Max 55% (±) *cis* and min. 45% (±) *trans*
***Contains xylene range aromatic solvents

Contains 3.2 pounds permethrin per gallon.

U.S. Patent No. 4,024,163

KEEP OUT OF REACH OF CHILDREN

CAUTION 279-3014

STATEMENT OF PRACTICAL TREATMENT

If in eye: flush eyes with water for 15 minutes. Call a physician.

If swallowed: do not induce vomiting. Call a physician.

If on skin: remove contaminated clothing. Wash with plenty of soap and water. Get medical attention if irritation persists.

Note to Physician: Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages by aspiration of the solvent

For Emergency Assistance Call 716-735-3765.

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

CAUTION
Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Remove and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service.

This product is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment, or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.



FMC Corporation
Agricultural Chemical Group
Philadelphia PA 19103

1087

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION

Area treated with Pounce 3.2 EC on (insert date). Do not enter without appropriate protective clothing until sprays have dried (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

CHEMIGATION USE DIRECTIONS

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hard move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Pounce 3.2 EC insecticide should be applied continuously for the duration of the water application. Pounce 3.2 EC should be diluted in sufficient volume to insure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation is not required when a suitable diluent is used.

Missouri

Barry, Benton, Camden, Christian, Dallas, Greene, Hickory, Jasper, Lawrence, Miller, Newton, Osage Park, St. Clair, Stone and Webster

Nevada

Clark

New Mexico

Chaves, DeBaca and Eddy

North Carolina

Edgecombe, Nash and Pitt

Ohio

Pickaway

Oklahoma

Delaware, McCurtain and Pushmataha

Oregon

Lake

Tennessee

Bedford, Blount, Claiborne, Decatur, Franklin, Hancock, Hardin, Hickman, Knox, Lawrence, Lincoln, Loudon, Marshall, Maury, Meigs, Monroe, Rhea, Roane, Scott, Sequatchie, Smith, Sullivan and Wayne

Texas

Bastrop, Burleson, Comal, Hams, Hays, Jeff Davis, Pecos and Reeves

Utah

Utah and Washington

Virginia

Lee, Russell, Scott, Smyth, Tazewell, Washington and Wise

STORAGE AND DISPOSAL

Pesticide Storage

Do not store below 10°F. (-12°C)

Do not use or store near heat, open flame or hot surfaces.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect (716) 735-3765.

To confine spill, if liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved State and local procedures. Do not cut or weld metal containers.

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ENDANGERED SPECIES RESTRICTIONS

The following restrictions apply to use of this product after February 1, 1988.

Before using this pesticide on corn, cotton or soybeans in the counties listed below, you must obtain the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES for the county in which the product is to be used. The bulletin is available from your County Extension Agent, State Fish and Game Office, or your pesticide dealer. Use of this product in a manner inconsistent with the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES is a violation of Federal laws.

Alabama

Colbert, Greene, Jackson, Lamar, Lauderdale, Limestone, Madison, Marshall, Morgan, Pickens and Sumter

Arizona

Graham, Maricopa, Mohave, Pima, Pinal and Santa Cruz

Arkansas

Benton, Clay, Clark, Cross, Lawrence, Lee, Poinsette, Polk, Randolph, Sharp and St. Francis

California

Butte, Colusa, Glenn, Imperial, Kern, Merced, Modoc, Riverside, Sacramento, Solano, Sutter, Tehama and Yolo

Florida

Florida

Kentucky

Ballard, Butler, Edmundson, Green, Hart, Jackson, Laurel, Livingston, Marshall, McCracken, McCreary, Pulaski, Rockcastle, Taylor, Warren and Wayne

Mississippi

Claiborne, Copiah, Hinds, Itawamba, Lowndes, Monroe and Noxubee

GENERAL INSTRUCTIONS

Pounce 3.2 EC is a 3.2 pounds per gallon formulation of the insecticide permethrin. Apply Pounce 3.2 EC when insects appear or feeding is noticed. The higher rate should be used as pest populations increase. Repeat the application as necessary to maintain control. Pounce 3.2 EC may be applied by both ground and aerial equipment. Use sufficient water to obtain full coverage. With the exception of crops listed below, rotational crops should not be planted within 60 days of last application.

This label must be in the possession of the user at the time of application.

Almonds: To control Navel Orangeworm, Peach Twig Borer—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pounds active) per acre. Apply when insects appear. Do not make more than two applications during hull split. Do not make more than 5 applications per season. Apply in a minimum of 15 gallons of finished spray per acre by aircraft or 20 gallons of finished spray per acre with ground equipment. Do not apply within seven days of harvest. Do not graze livestock in treated orchards.

Apples: To control Plum Curculio, Redbanded Leafroller, Rosy Apple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Oblique Banded Leafroller and Green Fruitworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pounds active) per acre with ground equipment only. Apply in 25 to 400 gallons of finished spray per acre when insects appear and repeat as required to maintain control. Do not make more than three applications per season. Do not apply after petal fall.

Asparagus: To control Cutworms and Asparagus Beetle—Use Pounce 3.2 EC at the rate of 2 to 4 ounces (0.05 to 0.1 lbs. active) per acre. Apply by ground equipment using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 4 applications. Do not apply within 3 days of harvest.

For post harvest application, use Pounce 3.2 EC at a rate of 4 ounces (0.1 lbs. active) per acre. Apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the Asparagus Beetle, Tarnished Plant Bug and Lygus Bugs and the adult stage of the Japanese beetle are present.

Broccoli, Chinese Broccoli (glaucan, white flowering broccoli), Brussels Sprouts, Cauliflower: To control Beet Armyworm, Cabbage Looper, Imported Cabbageworm and Diamondback Moth—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre. Apply in a minimum of 3 gallons of finished spray per acre by aircraft or 20 gallons per acre by ground equipment. Do not make more than 8 applications per season. Do not apply within one day of harvest.

Cabbage, Chinese Cabbage (tight-heading varieties only): To control Cabbage Looper, Diamondback Moth, Imported Cabbageworm, and Southern White Butterfly—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05 to 0.2 pound active) per acre. To control Beet Armyworm, Cutworms, and Flea Beetle—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply in a minimum of 1 gallon of finished spray per acre by aircraft or 20 gallons per acre with ground equipment. Do not apply more than 1 pound active ingredient per acre per season. Do not apply within 1 day of harvest.

Cantaloupes: To control Aphids, Leafminers, Squash Bug, and Whiteflies—Use Pounce 3.2 EC at a rate of 8 ounces (0.2 pound active) per acre. To control Cucumber Beetle, Cutworms, Melon Worm, Pickleworm, and Squash Vine Borer—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active ingredient per acre per season. Do not apply within one day of harvest.

Celery, Florence fennel (sweet anise, sweet fennel, finocchio): For general use directions refer to the Leafy Vegetable crop grouping. In California do not apply more than 2.0 pounds active ingredient per acre per season.

Cherries: To control Lesser Peach Tree Borer, Rose Chafer, Green Fruitworm, Red-banded Leafhopper, Plum Curculio and Tarnished Plant Bug—Use Pounce 3.2 EC insecticide at a rate of 1 to 2 ounces (0.025 to 0.05 pounds active) per 100 gallons of water. Do not exceed 400 gallons per acre. Apply when insects appear. East of the Rockies, do not exceed 6 applications per season, with no more than 4 applications after petal fall. West of the Rockies, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not graze livestock or feed crop forage from treated orchards.

Chrysanthemums (Greenhouse): For control of *Liriomyza* leafminer flies, use Pounce 3.2 EC at a rate of 20 fluid ounces (0.5 pound active) per 100 gallons (1 teaspoon per gallon). Avoid spraying the blooms. Pounce 3.2 EC may be applied on a weekly schedule.

Caution: Pounce 3.2 EC has demonstrated excellent plant safety, however, not all cultivars have been tested. Before treating large numbers of plants of a particular cultivar, treat a few plants and observe prior to full scale applications.

Conifers (Container and Field Grown): For control of Nantucket Pine Tip Moth use Pounce 3.2 EC at a rate of 4 to 8 fluid ounces (0.1 to 0.2 pound active) per acre. Pounce 3.2 EC may be diluted in a non-volatile vegetable oil or water in a minimum of 1 gallon of finished spray per acre using equipment calibrated to give adequate coverage. Begin application when the adults appear and repeat at 5 to 7 day intervals or as needed throughout the season.

Cotton: Preemergent Use: To control Cutworms—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply prior to planting or preemergent to crop. Use a minimum of 1 gallon of water per acre by aircraft or 5 gallons of water per acre by ground equipment. Pounce 3.2 EC has been found to be effective and compatible in tank mixtures with herbicides and liquid fertilizers.

Foliar Use: To control: Beet Armyworm (California and Arizona only), Bollworm, Cabbage Looper, Cotton Fleahopper, Cotton Leaf Perforator, Cutworms, Lygus Bugs, Pink Bollworm, Tarnished Plant Bug and Tobacco Budworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. To control Boll Weevil, alternate Pounce 3.2 EC at 4 ounces (0.1 pound active) per acre with Pounce 3.2 EC at 2 ounces (0.05 pound active) per acre every 3 to 4 days until control is achieved. Apply the above rates in a minimum of 1 gallon of water per acre by aircraft or 5 gallons of water per acre with ground equipment. Pounce 3.2 EC may also be applied using refined non-volatile vegetable oil for control of the above named insects. When applying in water by aircraft, 1 quart of oil may be substituted for 1 quart of water per gallon of finished spray. Pounce 3.2 EC should be diluted with oil and applied in a minimum of 1 quart total volume acre using equipment calibrated to give adequate coverage. To control Bollworm and Tobacco Budworm use 2 to 8 ounces (0.05 to 0.2 pound active) per acre. The 2 ounce rate per acre should be used only when insect infestation levels are light and cotton is closely scouted before and after application.

Do not apply more than 3.0 pounds active ingredient per acre per season. Do not apply within 14 days of harvest. Do not graze or feed cotton forage.

Cotton Tank-Mix: Pounce 3.2 EC insecticide Plus Tank Mixtures (all cotton growing states). Mix Pounce 3.2 EC at 4.0 to 8.0 ounces (0.1 to 0.2 pound active) per acre with other insecticides recommended on this label for control of cotton insect pests. Pounce 3.2 EC has been found to be effective and compatible in tank mixtures with the following products:

Methyl Parathion
EPN plus Methyl Parathion
Methomyl
Guthion

Azinphos-Methyl
Malathion
Chlordimeform

When using Pounce 3.2 EC in tank mixtures, observe all restrictions and precautions which appear on the labels of these products. Apply tank mixtures immediately after mixing.

Eggplants: To control Colorado Potato Beetle—Use Pounce 3.2 EC at the rate of 8 ounces (0.2 lbs. active) per acre. To control Vegetable Leafminer—Use Pounce 3.2 EC at the rate of 4 to 8 ounces (0.1 to 0.2 lbs. active) per acre. Apply using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 16 applications. Do not apply within 3 days of harvest.

Field Corn, Popcorn, Field Corn Grown for Seed: Preemergent Use: To control Armyworms, Cutworms, and Stalk Borers—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pounds active) per acre in the time period from five days prior to planting up to emergence of the crop. Apply either as a banded or broadcast spray in a minimum of 10 gallons finished spray per acre with ground equipment or 1 gallon per acre by air. Use a higher rate of Pounce 3.2 EC when incorporating into the

soil. Pounce 3.2 EC has been found compatible and may be tank mixed with the following herbicides:

AATC*	Lasso*	Prowl*
Banve*	Lorox*	Ramrod*
Blade*	Paraquat CL	Roundup*
Dual*	Princep*	Sutan*
Eradicane*		

When using Pounce 3.2 EC in tank mixtures, observe all restrictions and precautions which appear on the labels of these products. Provide constant agitation during mixing and application to keep the mixture in suspension. Label dosage rates should not be exceeded.

Foliar Use: For control of Armyworm, Corn Earworm, Corn Rootworm Beetles, Cutworms, European Corn Borer, Flea Beetle, Southwestern Corn Borer and Stalk Borers—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pounds active) per acre prior to the brown silk stage. For control of Western Bean Cutworm—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre prior to the brown silk stage.

When treating for stalk borer, Pounce 3.2 EC must be applied when or shortly before the stalk borer larvae are moving into the corn from surrounding weeds and grasses. Mowing or burndown herbicide are suggested to initiate movement. For control of Corn Earworm apply just before silking and continue as necessary to maintain control.

Apply a minimum of 1 gallon of finished spray per acre by air or 10 gallons per acre with ground equipment.

Filberts: To control Filbertworm and Oblique Banded Leafroller—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pounds active) per acre. For full coverage application apply 0.05 to 0.1 pounds active per 100 gallons (based on 400 gallons finished spray per acre), spray to run-off, OR, for low volume application apply 0.2 to 0.4 pounds active per acre (50 to 200 gallons finished spray per acre). For aerial application apply 0.2 to 0.4 pounds active in a minimum of 15 gallons of finished spray per acre. Apply when insects appear. Do not apply more than 1.6 pounds active per acre per season. Do not apply within 14 days of harvest. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock.

Head Lettuce: For general use directions refer to the Leafy Vegetable crop grouping.

Horseradish: To control Imported Crucifer Weevil (*Baris lepidii*).

As a spring preplant dip—Use a 0.1% active solution (2 pints, 1 ounce of Pounce 3.2 EC per 100 gallons). Soak sets for 30 minutes and air-dry before planting.

For foliar application, use Pounce 3.2 EC at a rate of 8 ounces (0.2 pound active) per acre. Apply by ground equipment in a minimum spray volume of 20 gallons per acre. Make up to 3 foliar applications as needed to control weevil adults during oviposition.

Do not apply within 22 days of harvest.

Leafy Vegetables (except Brassica): Amaranth; arrugula; celery; celtuce; chervil; corn salad; chrysanthemum (edible-leaved and garland); cress (garden and upland); dandelion; dock; endive; fennel; lettuce (leaf and head); orchard; parsley; purslane (garden and winter); rhubarb; spinach; Swiss chard. To control Beet Armyworm, Corn Earworm, Cutworms, European Corn Borer, Fall Armyworm, Green Cloverworm, Southern Armyworm, Tobacco Budworm and Vegetable Leafminer—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. To control Alfalfa Looper, Cabbage Looper and Leafhoppers—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05 to 0.2 pound active) per acre. Apply every 3 to 5 days or as needed. Use sufficient water to obtain full coverage of foliage. Do not apply within 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock.

Ornamental Nursery Stock (Field Grown): To control Bagworm, Beet Armyworm, Cabbage Looper, Citrus Thrips, *Malothus* spp., Lace Bug, Leafhoppers, Leafminers and Whiteflies—Use 4 to 8 ounces per 100 gallons of water. Pounce 3.2 EC may be used to control specified pests on non-edible ornamentals and non-bearing plants of fruiting species.

Caution: Marginal leaf burn may occur on *Salvia*, *Dietenbachia* and *Pteris* Fern. Application to blooming plants may cause browning of petals. Pounce 3.2 EC has demonstrated excellent plant safety; however, not all species and varieties have been tested. Before treating large numbers of plants of a particular variety, treat a few plants and observe prior to full scale application.

Peaches: To control Lesser Peach Tree Borer, Peach Twig Borer, Rose Chafer, Plum Curculio, Tarnished Plant Bug, Green Fruitworm, Oriental Fruit Moth—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per 100 gallons of water. Do not exceed 400 gallons per acre. Apply when insects appear. Make applications with ground equipment, using sufficient water to obtain uniform coverage. Do not apply within 7 days of harvest. Do not make more than 8 applications per season. Do not graze livestock in treated orchards.

Pears (Dormant through Delayed Dormant): To control Pear Psylla—Apply Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Pounce 3.2 EC may be combined with 2 to 8 gallons

of spray oil per acre. Apply during the dormant through delayed dormant growth periods only. Apply in 3 to 20 gallons of finished spray per acre by aircraft and 25 to 400 gallons per acre by ground equipment. Do not make more than two applications per season.

rs (Summer): To control Pear Psylla, Green Fruitworm, Codling Moths—Use Pounce 3.2 EC insecticide at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Use in sufficient water to achieve runoff and apply by ground equipment. Do not make more than 3 applications per season. Do not apply within 14 days of harvest.

Peppers (Bell): To control Vegetable Leafminer, Cabbage Looper, Flea Beetle, Corn Earworm, Pepper Weevil—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 lbs. active) per acre. To control European Corn Borer—Use Pounce 3.2 EC at the rate of 8 ounces (0.2 lbs. active) per acre. Apply using sufficient water to obtain uniform coverage. Apply as needed. Do not exceed 8 sprays per season. Do not apply within 30 days of harvest.

Pine Seed Orchards: To control Coneworms, Seed Bugs—Use Pounce 3.2 EC at the following rates:

13 ounces in 100 gallons of water (0.025% dilution by weight) for high volume sprayers.

68 ounces in 100 gallons of water (0.125% dilution by weight) for low volume sprayers;

48 ounces in not less than 10 gallons of water per acre for aerial application.

Apply first application within 1 week of female flower closure or peak pollen flight for webbing coneworm control. For other coneworms and seed bugs, apply first application within 30 days following female flower closure. Repeat applications at 4 week intervals, but do not apply more than 6 applications. Apply approximately 5 to 10 gallons of the 0.025% dilution per tree with high volume sprayers. With low volume sprayers, apply 100 gallons of the 0.125% dilution per acre. Do not graze or harvest cover crop. Avoid contact with open water.

Pistachios: To control Peach Twig Borer and Navel Orangeworm—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Use sufficient water to obtain full coverage of foliage. Do not make more than two applications per season. Do not apply after 10% split. Nuts may be harvested on the day of application. Do not graze livestock in treated orchards.

Potatoes: To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Cutworms, European Corn Borer, Potato Flea Beetle, Potato Leafhopper, Potato Psyllid, Potato Tuberworm, Aster Leafhopper and Tarnished Plant Bug. Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply the above rates in a minimum of 3 gallons of water per acre by aircraft or 20 gallons of water per acre with ground equipment. Pounce 3.2 EC may also be applied using refined non-volatile vegetable oil for control of the above-named insects. Pounce 3.2 EC should be diluted with oil and applied in a minimum of one quart total volume per acre using equipment calibrated to give adequate coverage. Do not make more than 12 applications per season. Do not apply within 7 days of harvest. Do not graze or feed potato forage.

Pumpkins: To control Aphids, Leafminers, Squash Bug, and Whiteflies—Use Pounce 3.2 EC at a rate of 8 ounces (0.2 pound active) per acre. To control Cucumber Beetle, Cutworms, Melon Worm, Pickleworm and Squash Vine Borer—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply within 1 day of harvest.

Range Grass: To control Range Caterpillar—Use Pounce 3.2 EC at a rate of 0.4 ounces (0.01 lbs. active) per acre. Apply using sufficient water to obtain uniform coverage. Do not apply more than once per year. Cattle may be present during application. Do not harvest or feed hay to livestock.

Roses (Field Grown): For control of *Heliothis* sp., use 4 to 8 fluid ounces (0.1 to 0.2 pound active) per acre. Pounce 3.2 EC may be applied in 5 gallons of finished spray per acre with ground equipment and 1 gallon per acre by air.

Roses (Greenhouse): For control of Beet Armyworm, Cabbage Looper, Omnivorous Leafroller, use Pounce 3.2 EC at a rate of 8 fluid ounces (0.2 pound active) per 100 gallons of water.

Caution: Pounce is not phytotoxic to the following varieties of greenhouse roses: Ballena, Bettina, Carolina, Coquette, Excitement, Forever Yours, G Wave, Jack Frost, The Bride's Maid, Malador, Paul's Pink, Samantha, Seventeen, Sonia, Town Crier, Tropicana and Visa. Other varieties may vary in their sensitivity to Pounce 3.2 EC, and a small number of plants should be treated under local conditions to determine plant safety prior to commercial use.

Soybeans: To control Cabbage Looper, Green Cloverworm, Mexican Bean Beetle, Soybean Looper, Cutworms, Velvetbean Caterpillar, Bean Leaf Beetle and Potato Leafhopper—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre. To control Beet Armyworm and Corn Earworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply a minimum of 1 gallon of finished spray per acre by air or 5 gallons with ground equipment.

Pounce 3.2 EC may also be applied using refined non-volatile vegetable oil for control of the above named insects. Pounce 3.2 EC should be diluted with oil and applied in a minimum of 1 quart total volume per acre using equipment calibrated to give adequate coverage. When applying in water by aircraft, 1 quart of oil may be substituted for 1 quart of water per gallon of finished spray.

Do not apply more than 0.4 pound active ingredient per acre per season. Do not apply within 60 days of harvest. Do not graze or feed soybean forage or hay.

Spinach: Refer to Leafy Vegetable crop grouping for general use directions.

Sweet Corn: To control Corn Earworm, Corn Rootworm Beetle*, Cutworms, European Corn Borer, Fall Armyworm, Southern Armyworm, Flea Beetle, Leafhoppers, and Stalk Borers—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply every 3 to 5 days or as needed. Do not apply more than 1.2 pounds active ingredient per acre per season. Do not apply within 1 day of harvest of ears or forage or livestock grazing.

*Pest does not occur on this crop in California.

Sweet Corn (Florida only) fresh market use only: To control Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm—Use 4 to 10 ounces (0.1 to 0.25 pound active) per acre. Apply every 5 days or as needed.

Do not apply more than 3.25 pounds active ingredient per acre per crop season. Do not graze livestock in treated areas or harvest for forage.

Tomatoes (Only For Application To Tomatoes Grown In Florida For Final Marketing As Fresh Tomatoes)

To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Granulate Cutworm, Hornworms, Vegetable Leafminers, Southern Armyworm, Tomato Fruitworm and Tomato Pinworm—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05 to 0.2 pound active) per acre. Apply by aerial or ground equipment in sufficient water to obtain uniform coverage. Do not apply more than 1.2 pound active ingredient per acre per season. Can be applied up to the day of harvest. Do not apply to cherry tomatoes or other varieties which produce fruit less than one inch in diameter.

Walnuts: To control Navel Orangeworm, Codling Moth and Walnut Husk Fly—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pounds active) per acre. For full coverage application apply 0.05 to 0.1 pounds active per 100 gallons (based on 400 gallons finished spray per acre), spray to run-off; OR, for low volume application apply 0.2 to 0.4 pounds active per acre (50 to 200 gallons finished spray per acre). For aerial application apply 0.2 to 0.4 pounds active in a minimum of 20 gallons of finished spray per acre. Do not apply more than 1.6 pounds active ingredient per acre per season. Apply when insects appear. Do not apply within 1 day of harvest. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock.

Premises Spray

For agricultural use only. Spray directly to walls and ceiling as residual surface treatment only. Do not treat manure or litter. Avoid contamination of feed and water. Do not apply directly to livestock or poultry.

For Application In	Target Insects	Method of Applic.	Dilute:	Applic. Rate
Dairies, Barns, Feedlots, Stables, Poultry Houses, Swine and Livestock Houses	House Flies, Stable Flies and other Manure Breeding Flies. Also aids in the reduction of Cockroaches, Mosquitoes and Spiders	Sprayer	4 oz. per 12 1/2 gal. of water	1 gal. per 750 sq. ft. of surface

Re-treat as necessary, but not more often than once every 2 weeks. The use of any residual fly spray should be supplemental with proper manure management and general sanitation to reduce or eliminate fly breeding site.


Pounce 3.2 EC

8/27/87

Almonds: To control Naval Orangeworm and Peach Twig Borer -- Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Apply when insects appear. Apply in a minimum of 15 gallons of finished spray per acre by aircraft or 20 gallons of finished spray per acre with ground equipment.

To control ants -- Apply Pounce 3.2 EC at a rate of 16 ounces (0.4 pound active) per acre by ground equipment in a minimum of 15 gallons of finished spray per acre. Application should follow mowing of weed growth to insure maximum coverage of the soil surface. Overhead moisture following application will enhance activity.

Do not apply more than 0.8 pound active per acre during hull split. Do not apply more than 2 pounds active per acre per season. Do not apply within seven days of harvest. Do not graze livestock in treated orchards.



Pounce 3.2 EC

10/21/87

Pistachios: To control Peach Twig Borer, Navel Orangeworm, Plant Bugs, Stink Bugs, and Leaf-footed Bugs -- Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Use sufficient water to obtain full coverage of foliage.

To control ants -- Apply Pounce 3.2 EC at a rate of 16 ounces (0.4 pound active) per acre by ground equipment in a minimum of 15 gallons of finished spray per acre. Application should follow mowing of weed growth to insure maximum coverage of the soil surface. Overhead moisture following application will enhance activity.

Do not apply more than 0.8 pound active per acre per crop season. Do not apply after 10% hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated orchards.

