

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

316-2766021  
7/24/94

Ms. Linda Dansbury  
FMC Corporation  
Agricultural Chemical Group  
1735 Market Street  
Philadelphia, PA 19103

AUG 11 1994

Dear Ms. Dansbury:

Subject: Updated Labeling  
Pounce 3.2 EC Insecticide  
EPA Registration No. 279-3014  
Pounce 25 WP Insecticide  
EPA Registration No. 279-3051  
Pounce WSB Insecticide  
EPA Registration No. 279-3083  
Your Letter Dated October 25, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable, provided that you make the following labeling changes listed below before you release the product for shipment:

1. For 279-3014, delete the section Commercial Impregnation and Application of Pounce 3.2 EC on Dry Bulk Fertilizers. Furthermore, once fertilizer is impregnated it would considered a new pesticide subject to FIFRA if it was resold. Your directions for use do not contain any safeguards to prevent th s from occurring.
2. When submitting finished labeling incorporate all Worker Protection Standard Labeling changes previously approved for the subject products.

Submit three copies of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely,

George T. LaRocca  
Product Manager 13

Insecticide-Rodenticide Branch  
Registration Division (7505C)

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

Code 3510

## RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

Net Contents

# Pounce® 3.2 EC Insecticide

EPA REG.NO. 279-3014

EPA Est. 279-

### ACTIVE INGREDIENTS:

\*Permethrin\*\* .....38.4%

INERT INGREDIENTS:\*\*\* .....61.6%

100.0%

\*(3-Phenoxyphenyl)methyl (±) cis-trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate

\*\*cis/trans ratio: Max. 55% (±) cis and min. 45% (±) trans

\*\*\*Contains xylene range aromatic solvents.

Contains 3.2 pounds permethrin per gallon.

U.S. Patent No. 4,024,163

## KEEP OUT OF REACH OF CHILDREN CAUTION

### STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush eyes with plenty of water. Call a physician.

If swallowed: Do not induce vomiting. Call a physician.

If on skin: Remove contaminated clothing. Wash with plenty of soap and water. Get medical attention if irritation persists.

Note to Physician: Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages by aspiration of the solvent.

For Emergency Assistance Call (800) 331-3148.

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans (and Domestic Animals)

##### CAUTION


Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Remove and wash clothing before reuse.

See other panels for additional precautionary information.

BEST AVAILABLE COPY

# FMC.

FMC Corporation  
Agricultural Chemical Group  
Philadelphia PA 19103



Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

279-3014

### Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water or wetlands (swamps, bogs, marshes, and pot-holes). Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

### Physical/Chemical Hazards

Do not use or store near heat or open flame.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

##### CAUTION

Area treated with Pounce 3.2 EC on (insert date). Do not enter without appropriate protective clothing until sprays have dried (insert State Department of Agriculture's reentry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

## STORAGE AND DISPOSAL

### Pesticide Storage

Do not store below 10° F. (- 12°C).

Do not use or store near heat, open flame or hot surfaces.

Keep out of reach of children and animals. Store in original containers only. Store in a cool dry place and avoid excess heat. Carefully open containers. After partial use replace lids and close tightly. Do not concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### Container Disposal

**Metal Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved State and local procedures. Do not cut or weld metal containers.

**Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning or dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Returnable/Refillable Sealed Container:** Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

## Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move and tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Pounce® 3.2 EC insecticide should be applied continuously for the duration of the water application. Pounce 3.2 EC should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.1 inch per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

## COMMERCIAL IMPREGNATION AND APPLICATION OF POUNCE 3.2 EC ON DRY BULK FERTILIZERS

Pounce 3.2EC insecticide may be impregnated on dry bulk fertilizers. When applied as directed, Pounce/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of Pounce applied in water.

The Pounce/fertilizer mixtures may be surface applied or shallow incorporated incorporated. The higher rate should be used if incorporation is used.

**Impregnation:** Apply using a minimum of 200 pounds of dry bulk fertilizer per acre and up to a maximum of 450 pounds per acre with the recommended amount of Pounce 3.2EC insecticide per acre. Use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform, fine spray pattern over the tumbling fertilizer for thorough coverage. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Pounce provides a satisfactory, dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry, flowable mixture. Microcel E (Johns-Manville Products Corporation) is a recommended absorbent powder. Generally less than 2% by weight of Microcel E is required. **DO NOT** impregnate Pounce 3.2EC onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the insecticide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with Pounce.

The amount of Pounce actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with Pounce 3.2EC insecticide should be applied immediately, not stored.

All state regulations, labeling, etc., of the mixtures are the responsibility of the seller.

## RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZERS WITH POUNCE 3.2 EC INSECTICIDE

PINTS OF POUNCE 3.2 EC PER TON OF FERTILIZER

Fertilizer Rate Lbs./Acre	POUNCE 3.2 EC Rate Per Acre		
	2 oz. (.05 lbs A.I.)	4 oz. (0.1 lb A.I.)	8 oz. (0.2 lbs A.I.)
200	20	40	80
250	16	32	64
300	13.3	26.7	53.3
350	11.4	22.9	45.7
400	10	20	40
450	8.9	17.8	35.6

If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will result in poor distribution during application.

## GENERAL INSTRUCTIONS

Pounce 3.2 EC is a 3.2 pounds per gallon formulation of the insecticide permethrin. Apply Pounce 3.2 EC when insects appear or feeding is noticed. The higher rate should be used as pest populations increase. Repeat the application as necessary to maintain control. Pounce 3.2 EC may be applied by both ground and aerial equipment. Use sufficient water to obtain full coverage. With the exception of crops listed below, rotational crops should not be planted within 60 days of last application.

This label must be in the possession of the user at the time of application.

**Alfalfa:** To control Alfalfa Cat, pillar, Armyworms, Cutworms, Loopers, Webworms, Blue Alfalfa Aphid, Green Peach Aphid, Pea Aphid, Green Cloverworm, Potato Leafhopper (knockdown only), Spotted Alfalfa Aphid, and Velvetbean Caterpillar—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05 to 0.2 pound active) per acre. To control Alfalfa Weevil, Egyptian Alfalfa Weevil, Cucumber Beetle, Meadow Spittlebug, Plant Bugs (including Lygus spp.), Potato Leafhopper, and Stink Bugs—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre.

Use higher recommended dosage for increased pest pressure or for increased residual pest control. Apply in a minimum of 1 gallon of finished spray per acre by aircraft or 10 gallons of finished spray per acre with ground equipment. Do not apply more than 0.2 pound active per cutting. When rates of 0.1 pound active per acre or less are used, application may be made on day of harvest. When rates greater than 0.1 pound active per acre are used, do not apply within 14 days of harvest.

**BEST AVAILABLE COPY**

**Almonds:** To control Navel Orangeworm, Peach Twig Borer—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Apply when insects appear. Apply in a minimum of 15 gallons of finished spray per acre by aircraft or 20 gallons of finished spray per acre with ground equipment.

To control ants—Apply Pounce 3.2 EC at a rate of 16 ounces (0.4 pound active) per acre by ground equipment in a minimum of 15 gallons of finished spray per acre. Application should follow mowing of weed growth to insure maximum coverage of the soil surface. Overhead moisture following application will enhance activity.

Do not apply more than 0.8 pound active per acre during hull split. Do not apply more than 2 pounds active per acre per season. Do not apply within 7 days of harvest. Do not graze livestock in treated orchards.

**Apples:** To control Plum Curculio, Redbanded Leafroller, Rosy Apple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Oblique Banded Leafroller and Green Fruitworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre with ground equipment only. Apply in 25 to 400 gallons of finished spray per acre when insects appear and repeat as required to maintain control. Do not apply more than 0.6 pound active per acre per season. Do not apply after petal fall.

**Artichoke:** To control Artichoke Plume Moth and Leafminers—Use Pounce 3.2 EC at a rate of 4 to 12 ounces (0.1 to 0.3 pound active) per acre. Apply with ground equipment using sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.5 pounds active per acre per season. Buds may be harvested on the day of application.

**Asparagus:** To control Cutworms and Asparagus Beetle—Use Pounce 3.2 EC at the rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre. Apply by ground equipment using sufficient water to obtain uniform coverage. Do not apply more than 0.4 pound active ingredient per acre per season. Do not apply within 1 day of harvest.

For post harvest application, use Pounce 3.2 EC at a rate of 4 ounces (0.1 pound active) per acre. Apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the Asparagus Beetle, Tarnished Plant Bug and Lygus Bugs and the adult stage of the Japanese beetle\* are present.

\*Not for control of this insect in California.

**Avocado:** To control Avocado Caterpillar, Avocado Lace Bug, Avocado Leafhopper, Avocado Leafroller, Orange Tortrix, Avocado Looper, Omnivorous Looper, Avocado Tree Girdler, Avocado Whitefly, Brown Soft Scale, Spanworm, Twig Borers, Mirids, Scale Crawlers, and Thrips—Use Pounce 3.2 EC at a rate of 8 ounces (0.2 pound active) per acre. Apply with ground equipment in a minimum of 20 gallons of water per acre. Apply when insects first appear and repeat at 7 to 10 day intervals as needed to provide control. Do not apply more than 1.2 pounds active per acre per season. Do not apply within 7 days of harvest.

**Broccoli, Chinese Broccoli (gai lan, white flowering broccoli), Brussels Sprouts, Cauliflower:** To control Armyworm spp., Cabbage Looper, Imported Cabbageworm, Diamondback Moth, Plant Bugs, and Thrips—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre. Apply in a minimum of 3 gallons of finished spray per acre by aircraft or 20 gallons per acre by ground equipment. Do not apply more than 0.8 pound active per acre per season. Do not apply within 1 day of harvest.

**Bulb Onions and Garlic:** To control Onion Thrips and Armyworms—Use Pounce 3.2 EC at a rate of 6 to 12 ounces (0.15 to 0.3 pound active) per acre. To control Cutworms, Leafminers, Onion Maggots (adults), and Stink Bugs—Use Pounce 3.2 EC at a rate of 4 to 12 ounces (0.1 to 0.3 pound active) per acre.

Apply in a minimum of 20 gallons per acre with ground equipment or in a minimum of 5 gallons per acre by aircraft. Begin applications when pests appear and repeat as necessary to maintain control. Use the higher label rates as Onion Thrips population increases and avoid rescue situations.

Do not apply more than 2.4 pounds active per acre per season. Do not apply within 1 day of harvest. Do not graze livestock in treated areas or cut treated crops for feed.

**Cabbage, Chinese Cabbage (tight-heading varieties only):** To control Cabbage Looper, Diamondback Moth, Imported Cabbageworm, and Southern White Butterfly—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05 to 0.2 pound active) per acre. To control Armyworm spp., Cutworms, and Flea Beetles—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply in a minimum of 1 gallon of finished spray per acre by aircraft or 20 gallons per acre with ground equipment. Do not apply more than 1 pound active ingredient per acre per season. Do not apply within 1 day of harvest.

**Cantaloupes:** For general use directions refer to the Cucurbit Vegetables crop grouping.

**Celery Florence fennel (sweet anise, sweet fennel, finocchio):** For general use directions refer to the Leafy Vegetable crop grouping. In California do not apply more than 2.0 pounds active ingredient per acre per season.

**Cherries:** To control Lesser Peach Tree Borer, Rose Chafer, Green Fruitworm, Red-banded Leafroller, Plum Curculio and Tarnished Plant Bug—Use Pounce 3.2 EC insecticide at a rate of 4 to 8 ounces (0.10 to 0.20 pound active) per acre as a dilute spray. Do not exceed 400 gallons per acre. Apply when insects appear. East of the Rockies, do not exceed 6 applications per season, with no more than 4 applications after petal fall. West of the Rockies, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not graze livestock or feed crop forage from treated orchards.

**Chrysanthemums:** For control of Liomyza Leafminer flies, use Pounce 3.2 EC at a rate of 20 fluid ounces (0.5 pound active) per 100 gallons (1 teaspoon per gallon). Avoid spraying the blooms. Pounce 3.2 EC may be applied on a weekly schedule.

**Caution:** Pounce® 3.2 EC has demonstrated excellent plant safety, however not all cultivars have been tested. Before treating large numbers of plants of a particular cultivar, treat a few plants and observe prior to full scale applications.

**Collards and Turnips:** To control Beet Armyworm, Corn Earworm, Southern White Butterfly, Leafminer, Leafhoppers, European Corn Borer, Fall Armyworm Cutworms, Green Cloverworm, Southern Armyworm, Tobacco Budworm, Vegetable Leafminer, Cabbage Looper Imported Cabbageworm, and Diamondback Moth—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre. Do not apply within 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock. Apply with ground equipment only. For use on Collards in AR, AZ, GA, IL, NC, OK, SC, and TX and on Turnips in FL, GA, IL, IN, OK, SC, TX, and WA. Do not apply more than 0.8 pound active ingredient per acre per season on Collards and Turnips (AR, TX & IN). Do not apply more than 0.4 pound active ingredient per acre per season on Turnips (SC, GA, FL & WA).

**Conifers (Container and Field Grown):** For control of Nantucket Pine Tip Moth use Pounce 3.2 EC at a rate of 4 to 8 fluid ounces (0.1 to 0.2 pound active) per acre. Pounce 3.2 EC may be diluted in a non-volatile vegetable oil or water in a minimum of 1 gallon of finished spray per acre using equipment calibrated to give adequate coverage. Begin application when the adults appear and repeat at 5 to 7 day intervals or as needed throughout the season.

**Cotton:** Preemergent Use: To control Cutworms—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply prior to planting or preemergent to crop. Use a minimum of 1 gallon of water per acre by aircraft or 5 gallons of water per acre by ground equipment. Pounce 3.2 EC has been found to be effective and compatible in tank mixtures with herbicides and liquid fertilizers.

**Foliar Use:** To control Beet Armyworm (California and Arizona only), Bollworm, Cabbage Looper, Cotton Fleahopper, Cotton Leaf Perforator, Cutworms, Lygus Bugs, Pink Bollworm, Tarnished Plant Bug and Tobacco Budworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. To control Boll Weevil, alternate Pounce 3.2 EC at 4 ounces (0.1 pound active) per acre with Pounce 3.2 EC at 2 ounces (0.05 pound active) per acre every 3 to 4 days until control is achieved. Apply the above rates in a minimum of 1 gallon of water per acre by aircraft or 5 gallons of water per acre with ground equipment. Pounce 3.2 EC may also be applied using refined non-volatile vegetable oil for control of the above named insects. When applying in water by aircraft, 1 quart of oil may be substituted for 1 quart of water per gallon of finished spray. Pounce 3.2 EC should be diluted with oil and applied in a minimum of 1 quart total volume/acre using equipment calibrated to give adequate coverage. To control Bollworm and Tobacco Budworm use 2 to 8 ounces (0.05 to 0.2 pound active) per acre. The 2 ounce rate per acre should be used only when insect infestation levels are light and cotton is closely scouted before and after application.

Do not apply more than 3.0 pounds active ingredient per acre per season. Do not apply within 14 days of harvest. Do not graze or feed cotton forage.

**Cotton Tank-Mix:** Pounce 3.2 EC insecticide and Tank Mixtures (all cotton growing states). Mix Pounce 3.2 EC at 0 to 8.0 ounces (0.1 to 0.2 pound active) per acre with other insecticides recommended on this label for control of cotton insect pests. Pounce 3.2 EC has been found to be effective and compatible in tank mixtures with the following products:

Methyl Parathion	Azinphos-Methyl
Metomyl	Malathion
Guthion®	

When using Pounce 3.2 EC in tank mixtures, observe all restrictions and precautions which appear on the labels of these products. Apply tank mixtures immediately after mixing.

**BEST AVAILABLE COPY**

#### Spray Drift Precautions—Cotton

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

Do not make aerial or ground applications during temperature inversions.

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. Boom length must not exceed 75% of wing span or rotor diameter.

Do not apply by ground equipment within 25 feet, or by air within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush<sup>®</sup>, Ammo<sup>2</sup>, Asana<sup>®</sup> XL, Baythroid<sup>®</sup>, Capture<sup>®</sup>, Cymbush<sup>®</sup>, Karate<sup>®</sup>, Pounce<sup>®</sup> and Scout<sup>®</sup> X-TRA.

**Cucurbits:** Balsam pear (bitter melon); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourds, edible; melons, including hybrids such as cantaloupe, casaba, crenshaw, honeydew melons, honey balls, mango melon, muskmelon, Persian melon; pumpkin; squash, summer and winter; watermelon, including hybrids. To control Aphids, Leafminers, and Squash Bug—Use Pounce 3.2 EC at a rate of 8 ounces (0.2 pound active) per acre. To control Cabbage Looper, Cucumber Beetle (adults), Cutworms, Leafhoppers, Melonworm, Pickleworm, Plant Bugs (including Lygus and Stink Bugs), Rindworms, and Squash Vine Borer—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply a minimum of 4 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active ingredient per acre per season. Pounce 3.2 EC may be applied up to harvest.

**Eggplants:** To control Colorado Potato Beetle—Use Pounce 3.2 EC at the rate of 8 ounces (0.2 pound active) per acre. To control Cabbage Looper, Flea Beetles, and Vegetable Leafminer—Use Pounce 3.2 EC at the rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply using sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 3.2 pounds active ingredient per acre per season. Do not apply within 3 days of harvest.

**Field Corn, Popcorn, Field Corn Grown for Seed: Preemergent Use:** To control Armyworms, Cutworms and Stalk Borers—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre in the time period from five days prior to planting up to emergence of the crop. Apply either as a banded or broadcast spray in a minimum of 10 gallons finished spray per acre with ground equipment or 1 gallon per acre by air. Use a higher rate of Pounce 3.2 EC when incorporating into the soil. Pounce 3.2 EC has been found compatible and may be tank mixed with the following herbicides:

AAtrex <sup>®</sup>	Lasso <sup>®</sup>	Prow <sup>®</sup>
Banvel <sup>®</sup>	Lorox <sup>®</sup>	Ramrod <sup>®</sup>
Bladex <sup>®</sup>	Paraquat CL	Roundup <sup>®</sup>
Dual <sup>®</sup>	Princep <sup>®</sup>	Sutan <sup>®</sup>
Eradicane <sup>®</sup>		

When using Pounce 3.2 EC in tank mixtures, observe all restrictions and precautions which appear on the labels of these products. Provide constant agitation during mixing and application to keep the mixture in suspension. Label dosage rates should not be exceeded.

**Foliar Use:** For control of Armyworm (including Fall Armyworm), Corn Earworm, Corn Rootworm Beetles, Cutworms, European Corn Borer, Flea Beetle, Hop Vine Borer, Southwestern Corn Borer, Stalk Borers, and Tobacco Budworm—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre prior to the brown silk stage. For control of Western Bean Cutworm—Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre prior to the brown silk stage.

When treating for stalk borer, Pounce 3.2 EC must be applied when or shortly before the stalk borer larvae are moving into the corn from surrounding weeds and grasses. Mowing or burndown herbicide are suggested to initiate movement. For control of Corn Earworm apply just before silking and continue as necessary to maintain control.

Apply a minimum of 1 gallon of finished spray per acre by air or 10 gallons per acre with ground equipment.

**Filberts:** To control Filbertworm and Oblique Banded Leafroller—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. For full coverage application apply 0.05 to 0.1 pound active per 100 gallons (based on 400 gallons finished spray per acre) and spray to runoff, OR for low volume application apply 0.2 to 0.4 pound active per acre (50 to 200 gallons finished spray per acre). For aerial application apply 0.2 to 0.4 pound active in a minimum of 15 gallons of finished spray per acre. Apply when insects appear. Do not apply more than 1.6 pounds active per acre per season. Do not apply within 14 days of harvest. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock.

**Head Lettuce:** For general use directions refer to the Leafy Vegetable crop grouping.

**Horseradish:** To control imported Crucifer Weevil (*Bans lepidi*).

As a spring preplant dip—Use a 0.1% active solution (2 pints, 1 ounce of Pounce 3.2 EC per 100 gallons). Soak sets for 30 minutes and air-dry before planting.

For foliar application, use Pounce 3.2 EC at a rate of 8 ounces (0.2 pound active) per acre. Apply by ground equipment in a minimum spray volume of 20 gallons per acre. Make up to 3 foliar applications as needed to control weevil adults during oviposition.

Do not apply within 22 days of harvest.

**Leafy Vegetables (except Brassica):** Amaranth; arrugula; celery; celtuce; chervil; corn salad; chrysanthemum (edible-leaved and garland); cress (garden and upland); dandelion; dock; endive; fennel; lettuce (leaf and head); orach; parsley; purslane (garden and winter); rhubarb; spinach; Swiss chard. To control Aphids, Beet Armyworm, Corn Earworm, Cutworms, European Corn Borer, Fall Armyworm, Green Cloverworm, Southern Armyworm, Tobacco Budworm and Vegetable Leafminer—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. To control Alfalfa Looper, Cabbage Looper and Leafhoppers—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05 to 0.2 pound active) per acre. Apply every 3 to 5 days or as needed by air or ground. Use sufficient water to obtain full coverage of foliage. Do not apply more than 2.0 pounds active ingredient per acre per season. Can be applied up to 1 day of harvest.

**Ornamental Nursery Stock (Field Grown):** To control Bagworm, Beet Armyworm, Cabbage Looper, Citrus Thrips, *Heliothis* spp., Lace Bug, Leafhoppers, Leafminers, and Whiteflies—Use 4 to 8 ounces per 100 gallons of water. Pounce 3.2 EC may be used to control specified pests on non-edible ornamentals and non-bearing plants of fruiting species.

**Caution:** Marginal leaf burn may occur on *Salvia*, *Dieffenbachia* and *Pteris* Fern. Application to blooming plants may cause browning of petals. Pounce 3.2 EC has demonstrated excellent plant safety; however, not all species and varieties have been tested. Before treating large numbers of plants of a particular variety, treat a few plants and observe prior to full scale application.

**Papaya (Florida Only):** To control Papaya Fruit Fly, Papaya Webworm, Papaya Whitefly, Brown Soft Scale, Scale Crawlers, Aphids, and Mealybug—Use Pounce 3.2 EC at a rate of 8 ounces (0.2 pound active) per acre. Apply with ground equipment in a minimum of 20 gallons of water per acre. Apply when insects first appear and repeat at 7 to 10 day intervals as needed to provide control. Do not apply more than 1.2 pounds active per acre per season. Do not apply within 7 days of harvest.

**Peaches:** To control Lesser Peach Tree Borer, Peach Twig Borer, Rose Chafer, Plum Curculio, Tarnished Plant Bug, Green Fruitworm, Oriental Fruit Moth—Use Pounce 3.2 EC at a rate of 4 to 16 ounces (0.10 to 0.40 pound active) per acre. Apply using 25-400 gallons of spray per acre with ground equipment or a minimum of 10 gallons per acre by air. Spray to wet all foliage. Can be applied up to 7 days before harvest. Do not apply more than 3.0 pounds active ingredient per acre per season. Do not graze livestock in treated orchards.

**Pears (Dormant through Delayed Dormant):** To control Pear Psylla—Apply Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Pounce 3.2 EC may be combined with 2 to 8 gallons of spray oil per acre. Apply during the dormant through delayed dormant growth periods only. Apply in 3 to 20 gallons of finished spray per acre by aircraft and 25 to 400 gallons per acre by ground equipment. Do not apply more than 0.8 pound active per acre per season.

**Pears (Summer):** To control Pear Psylla, Green Fruitworm, Codling Moths—Use Pounce 3.2 EC insecticide at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Use in sufficient water to achieve runoff and apply by ground equipment. Do not apply more than 1.2 pounds active per season. Do not apply within 14 days of harvest.

**Peppers (Bell):** To control Vegetable Leafminer, Cabbage Looper, Cutworms, Flea Beetle, Corn Earworm, Pepper Weevil—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. To control European Corn Borer—Use Pounce 3.2 EC at the rate of 8 ounces (0.2 pound active) per acre. Apply using sufficient water to obtain uniform coverage. Apply as needed. Do not apply more than 1.6 pounds active ingredient per acre per season. Do not apply within 3 days of harvest.

**BEST AVAILABLE COPY**

**Pine Seed Orchards:** To control Coneworms, Seed Bugs—Use Pounce 3.2 EC at the following rates.

For high volume sprayers: Use 8 ounces in 100 gallons of water (0.025% dilution by weight). Apply 5 to 10 gallons of finished spray per tree.

For low volume sprayers: Use 42 ounces in 100 gallons of water (0.125% dilution by weight). Apply 100 gallons per acre.

For aerial application: Use 30 ounces in not less than 10 gallons of water per acre.

To control Webbing Coneworm—make first application within 1 week of female flower closure or peak pollen flight.

To control other coneworms and seed bugs—make first application within 30 days following female flower closure.

Repeat applications at 4 week intervals, but do not apply more than 6 applications. Do not graze or harvest cover crop. Avoid contact with open water.

**Pistachios:** To control Peach Twig Borer, Navel Orangeworm, Plant Bugs, Stink Bugs, and Leaf-footed Bugs—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. Use sufficient water to obtain full coverage of foliage.

To control ants—Apply Pounce 3.2 EC at a rate of 16 ounces (0.4 pound active) per acre by ground equipment in a minimum of 15 gallons of finished spray per acre. Application should follow mowing of weed growth to insure maximum coverage of the soil surface. Overhead moisture following application will enhance activity.

Do not apply more than 0.8 pound active per acre per crop season. Do not apply after 10% hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated orchards.

**Potatoes:** To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Cutworms, European Corn Borer, Potato Aphid, Potato Flea Beetle, Potato Leafhopper, Potato Psyllid, Potato Tuberworm, Aster Leafhopper and Tarnished Plant Bug. Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply the above rates in a minimum of 3 gallons of water per acre by aircraft or 20 gallons of water per acre with ground equipment. Pounce 3.2 EC may also be applied using refined non-volatile vegetable oil for control of the above-named insects. Pounce 3.2 EC should be diluted with oil and applied in a minimum of one quart total volume per acre using equipment calibrated to give adequate coverage. Do not apply more than 2.4 pounds active ingredient per acre per season. Do not apply within 7 days of harvest. Do not graze or feed potato forage.

**Pumpkins:** For general use directions refer to the Cucurbit Vegetables crop grouping.

**Range Grass:** To control Range Caterpillar—Use Pounce 3.2 EC at a rate of 0.4 ounces (0.01 lb. active) per acre. Apply using sufficient water to obtain uniform coverage. Do not apply more than once per year. Cattle may be present during application. Do not harvest or feed hay to livestock.

**Roses (Field Grown):** For control of *Heliothis* spp., use 4 to 8 fluid ounces (0.1 to 0.2 pound active) per acre. Pounce 3.2 EC may be applied in 5 gallons of finished spray per acre with ground equipment and 1 gallon per acre by air.

**Roses (Greenhouse):** For control of Beet Armyworm, Cabbage Looper and Omnivorous Leafroller, use Pounce 3.2 EC at a rate of 8 fluid ounces (0.2 pound active) per 100 gallons of water.

**Caution:** Pounce is not phytotoxic to the following varieties of greenhouse roses: Ballena, Bettina, Cara Mia, Coquette, Excitement, Forever Yours, G. Wave, Jack Frost, Jr. Bridesmaid, Matador, Paul's Pink, Samantha, Seventeen, Sonia, Town Crier, Tropicana and Visa. Other varieties may vary in their sensitivity to Pounce 3.2 EC, and a small number of plants should be treated under local conditions to determine plant safety prior to commercial use.

**Soybeans:** To control Bean Leaf Beetle, Cabbage Looper, Cutworms, Flea Beetles, Green Cloverworm, Japanese Beetle, Mexican Bean Beetle, Potato Leafhopper, Saltmarsh Caterpillar (Woolly Bear Caterpillar), Thistle Caterpillar, and Velvetbean Caterpillar. —Use Pounce 3.2 EC at a rate of 2 to 4 ounces (0.05 to 0.1 pound active) per acre. To control Beet Armyworm, Corn Earworm, Soybean Looper and Webworms—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply a minimum of 1 gallon of finished spray per acre by air or 5 gallons with ground equipment.

Pounce 3.2 EC may also be applied using refined non-volatile vegetable oil for control of the above named insects. Pounce 3.2 EC should be diluted with oil and applied in a minimum of 1 quart total volume/acre using equipment calibrated to give adequate coverage. When applying in water by aircraft, 1 quart of oil may be substituted for 1 quart of water per gallon of finished spray.

Do not apply more than 0.4 pound active ingredient per acre per season. Do not apply within 60 days of harvest. Do not graze or feed soybean forage or hay.

**Spinach:** Refer to Leafy Vegetable crop grouping for general use directions.

**Sweet Corn:** To control Corn Earworm, Corn Rootworm Beetle\*, Cutworms, European Corn Borer, Fall Armyworm, Flea Beetle, Hop Vine Borer, Leafhoppers, Southern Armyworm, and Stalk Borers—Use Pounce 3.2 EC at a rate of 4 to 8 ounces (0.1 to 0.2 pound active) per acre. Apply every 3 to 5 days or as needed. Do not apply more than 1.2 pounds active ingredient per acre per season. Do not apply within 1 day of harvest of ears or forage or livestock grazing.

\*Pest does not occur on this crop in California.

**Sweet Corn (Florida only) fresh market use only:** To control Aster Leafhopper, Corn Earworm, Cutworms, European Corn Borer, Fall Armyworm, Southern Armyworm—Use 4 to 10 ounces (0.1 to 0.25 pound active) per acre. Apply every 3 to 5 days or as needed.

Do not apply more than 3.25 pounds active ingredient per acre per crop season. Do not graze livestock in treated areas or harvest for forage.

**Tomatoes (Only For Application To Tomatoes Grown In Florida For Final Marketing As Fresh Tomatoes)**

To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Granulate Cutworm, Hornworms, Vegetable Leafminers, Southern Armyworm, Tomato Fruitworm and Tomato Pinworm—Use Pounce 3.2 EC at a rate of 2 to 8 ounces (0.05 to 0.2 pound active) per acre. Apply by aerial or ground equipment in sufficient water to obtain uniform coverage. Do not apply more than 1.2 pound active ingredient per acre per season. Can be applied up to the day of harvest. Do not apply to cherry tomatoes or other varieties which produce fruit less than one inch in diameter.

**Walnuts:** To control Navel Orangeworm, Codling Moth and Walnut Husk Fly—Use Pounce 3.2 EC at a rate of 8 to 16 ounces (0.2 to 0.4 pound active) per acre. For full coverage application apply 0.05 to 0.1 pound active per 100 gallons (based on 400 gallons finished spray per acre), spray to run-off; OR, for low volume application apply 0.2 to 0.4 pound active per acre (50 to 200 gallons finished spray per acre). For aerial application apply 0.2 to 0.4 pound active in a minimum of 20 gallons of finished spray per acre. Do not apply more than 1.6 pounds active ingredient per acre per season. Apply when insects appear. Do not apply within 1 day of harvest. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock.

#### Premises Spray

For agricultural use only.

Spray directly to walls and ceiling as residual surface treatment only. Do not treat manure or litter. Avoid contamination of feed and water. Do not apply directly to livestock or poultry.

For Application in	Target Insects	Method of Applic.	Dilute	Applic. Rate
Dairies, Barns, Feedlots, Stables, Poultry Houses, Swine and Livestock Houses	House Flies, Stable Flies and other Manure Breeding Flies. Also aids in the reduction of Cockroaches, Mosquitoes and Spiders.	Sprayer	4 ounces to 12.5 gallons water	1 gallon per 750 square feet of surface

Re-treat as necessary, but not more often than once every 2 weeks. The use of any residual fly spray should be supplemental with proper manure management and general sanitation to reduce or eliminate fly breeding site.

**BEST AVAILABLE COPY**

## Treatment of Preconstruction Lumber and Logs

### General Information

Pounce 3.2 EC insecticide should be diluted with water. To prepare the spray, dilute Pounce 3.2 EC as shown in the following spray dilution chart.

Spray Dilution Chart

Gallons of Spray Mixture Desired	Gallons of Pounce® 3.2 EC Insecticide To Use		
	0.5% Solution	0.75% Solution	1.0% Solution
40	1/2	3/4	1
80	1	1 1/2	2
200	2 1/2	3 3/4	5
400	5	7 1/2	10
800	10	15	20

### Directions for Application

To protect unseasoned lumber and logs from wood destroying insects, such as Termites, Carpenter Ants and Beetles (Ambrosia, Powder-post, Old house borers and others), totally treat wood with a 0.5% to 1.0% solution of Pounce® 3.2 EC. This solution can be applied by various methods including spraying, brushing, dipping, and pressure treatment. Frequent monitoring of dip and pressure systems are necessary to insure that the desired level of Pounce 3.2 EC is maintained. Wood can be handled after treatment when dry.

1. For dip treatments, the wood should be totally submersed in the solution until thoroughly wet and then allowed to dry in a suitable location. Dipping solutions to which Pounce 3.2 EC has been added should be agitated before use if left unused for long periods of time. Sediment, debris and other deposits should be periodically cleaned from the tank.
2. For pressure treatments the wood should be placed in the treatment chamber, the Pounce 3.2 EC solution added and the system pressurized up to 250 psi for up to one hour depending on the density and type of wood treated. After the pressure is released and the system drained, the wood should be placed in a suitable location for drying.
3. For spray treatments, the wood should be sprayed thoroughly including back and ends.
4. For brush treatments, all parts of wood surfaces should be thoroughly treated.

### Dealers Should Sell in Original Packages Only.

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

AAtrex, Dual and Princep—Trademarks of Ciba Geigy Corporation

Ambush, Cymbush, Eradicane, Karate and Sutan—Trademarks of ICI Americas Inc.

Asana, Bladex and Lorox—Trademarks of E.I. duPont de Nemours and Co., Inc.


Banvel—Trademark of Sandoz AG

Baythroid and Guthion—Trademarks of Bayer AG

Lasso, Ramrod and Roundup—Trademarks of Monsanto Company

Prowl—Trademark of American Cyanamid Company

Scout—Trademark of Hoechst Roussel Agri-Vet Company

Am. 3, Capture, Pounce and —Trademarks of FMC Corporation  
(3510-3/26/91-A)

**BEST AVAILABLE COPY**