



ACCEPTED
with COMMENTS
in EPA Letter Dated:

JUN 13 1984

Under the Federal Insecticide,
Fungicide, and Rodenticide
Act as amended, for the purposes
registered under FIFRA

279-3019

Code 3811

Captan 50-W Fungicide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-3019

EPA Est., 279-

Active Ingredients	By Wt.
Captan	47.3%
Related Derivatives	2.7%
Inert Ingredients	<u>50.0%</u>
	100.0%

*N-[(Trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

**DANGER
PELIGRO**

PRECAUCION AL USUARIO: Si usted no ha leído el etiquetado de este producto hasta que la etiqueta le haya sido aplicada, no use el producto.

Statement of Practical Treatment

If in eyes - immediately flush with large amounts of water for at least 15 minutes. Call a physician.

If swallowed - drink 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

If inhaled - remove victim to fresh air. If not breathing give artificial respiration preferably mouth-to-mouth.

If on skin - wash skin with soap and water.

For Emergency Assistance Call 716-735-3785.

See Other Panels for Additional Precautionary Information.

FMC[®]

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

FA 279-3019

EPA Reg. No. 279-3019
279-3019-1
Page 1 of 1

ALL INFORMATION AVAILABLE

**Net Weight
Lbs.**

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Danger

Corrosive, causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield, and rubber gloves when handling. Harmful if swallowed. Avoid breathing dust. Use with adequate ventilation. Wash thoroughly after handling. Do not contaminate food or foodstuffs.

Environmental Hazards

This pesticide is toxic to fish. Do not discharge into lakes, streams, ponds or public water unless in accordance with a National Pollution Discharge Elimination System permit. For guidance, contact your regional office of EPA. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until dusts have settled.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

DANGER

Area treated with Daptac 50 WP (insecticide). Do not enter without appropriate protective clothing until sprays have dried. (insert State Department of Agriculture re entry intervals if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash immediately with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with a lot of water. If inhaled, get fresh air and seek medical attention if needed.

279-3019

BEST DOCUMENT AVAILABLE

Storage and Disposal

Pesticide Storage

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Captan 50-W is a micro-fine wettable powder for use in water as a spray for the control of certain fungus diseases of fruit, vegetables and ornamental crops, and as a soil treatment for the control of certain seed rots and damping-off diseases.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to adjoining food, fiber or pasture crops.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as the timing of applications needed will vary with local conditions.

COMPATIBILITY AND PLANT SAFETY

Captan 50-W can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and bordeaux mixture will reduce the fungicidal activity of Captan. Do not apply Captan in combination with or immediately before or closely following oil sprays. The time factor governing the safe interval between Captan and oil sprays varies due to general climatic conditions; therefore consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of Captan and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in frequent sprays, Captan may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums and cherries. This type of injury is not likely to occur in the early cover sprays required to prevent warm, cloudy, humid weather. To avoid the hazard of leaf spotting, in trees, in conditions, use Captan and other spray materials at lowest recommended rates and avoid drenching trees.

MIXING INSTRUCTIONS

For water sprays, mix with water in the following order: 1. Add Captan 50-W to a material into spray tank. Add water. 2. Add other materials. 3. Mix thoroughly. 4. Spray. For soil treatments, mix with water in the following order: 1. Add Captan 50-W to a material into spray tank. Add water. 2. Mix thoroughly. 3. Apply to soil. For more information on safety, use of Captan 50-W, refer to Precautions.

EPA Reg. No. 279-3019
1973-11-23
Page 3 of 4

279 3019

BEST DOCUMENT AVAILABLE

RECOMMENDATIONS

Unless otherwise specified, rates are given in terms of Captan 50-W per 100 gallons water for fruit and nut crops and ornamentals. For vegetable crops, rates are expressed as Captan 50-W per acre in sufficient water to provide thorough coverage. Except as specified, begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, applications can be made on the day of harvest.

Maximum permissible rate per acre, per application of Captan 50-W is given in parentheses, (), after crop claim.

FRUIT AND NUT CROPS

IMPORTANT Read label carefully. Although most of the directions on this label may be followed nationwide, a few are limited to either the eastern or western U.S. Follow these directions for your growing area where specified.

Almonds: Brown rot twig and blossom blight, shothole, scab, leaf blight—Use 2 to 3 pounds. Apply at popcorn, bloom, and petal fall stages and up to 5 weeks after petal fall. Hulls may be fed to livestock (32 pounds).

Subsequent applications may be made up to 12 days of harvest in accordance with pest infestations and State Agricultural Extension Service recommendations, provided the hulls are not fed to dairy animals or animals being finished for slaughter (32 pounds).

Apples (Eastern U.S.): Primary scab, black rot (frog-eye), botrytis blossom rot—Use 1 1/4 to 2 pounds at 5 to 7 day intervals as needed to maintain cover in prebloom, bloom, petal fall and first cover sprays. Use at 2 pound rate if conditions favor scab or if active scab can be found. Secondary scab, Brooks fruit spot, sooty blotch, fly speck, black rot, black pox, botryospaeria rot, bitter rot—Use 1 to 2 pounds at 10 to 14 day intervals in second and later cover sprays up to start of harvest (16 pounds). Powdery mildew—Add 1/4 to 3/4 pound sulfur or 1/4 to 1/2 pound Karathane to all post bloom sprays until the foliage matures. Use 1/2 pound Karathane if interval between sprays is longer than 7 days. Do not use Captan in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Staygreen, Baldwin, King, etc., as severe injury and defoliation may occur.

Apples (Western U.S.): Apple scab—In preblossom sprays, use 2 lbs per 100 gallons of water in high volume application. In low volume sprays, apply 16 pounds per acre. In mid-summer, the dosage may be reduced to 1 pound per 100 gallons in high volume application or 10 pounds per acre in low volume sprays. **(Pacific Northwest)** Bull's eye rot, Botrytis rot—2 pounds per 100 gallons spray. Make 1 to 2 applications with late cover sprays and 1 to 2 preharvest applications (16 pounds).

Apricots: Brown rot twig blight, jacket rot—Use 2 pounds. Apply in bud, bloom, 75% petal fall, and cover stages (10 pounds).

Blackberries, Dewberries, Raspberries (Eastern U.S.): Anthracnose—Use 4 pounds per acre in sufficient water for thorough coverage in the preblossom (when new canes are 6 to 8 inches long) and first cover blight after petal fall sprays. **Raspberries - Botrytis fruit rot -** Use 2 pounds per 100 gallons water. Begin application in the first bloom. Make 2 or more repeat applications (as field conditions warrant) at 10 to 15 day intervals (16 pounds). **Raspberries (Western U.S.):** Anthracnose, Botrytis spur blight—Apply 2 pounds at bloom or when new canes are 6 to 10 inches long. Repeat 2 weeks later. Apply late spray after canes are removed. Fruit rot—Apply 2 pounds 3 to 5 days before harvest starts. Repeat at mid-harvest and 8 to 10 days later (16 pounds).

Blueberries (Western U.S.): Botrytis rot, Botrytis blight, mummy rot—Use 2 pounds per 100 gallons water in sufficient water for thorough coverage. Begin application at bloom and repeat at 7 to 10 day intervals (16 pounds).

Cherries (Eastern U.S.): Brown rot, leaf spot, Botrytis rot—Use 2 pounds in prebloom, bloom, petal fall, shuck cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat applications at 7 to 10 day intervals until 10 days before harvest. In prebloom, petal fall, and early cover sprays, if sulfur is added, Captan may be reduced to 1 pound in these sprays. Post harvest sprays—Apply 2 pounds immediately after harvest and repeat applications at 7 to 14 days.

Cherries (Western U.S.): Anthracnose, Botrytis rot—Use 2 pounds per 100 gallons water in sufficient water for thorough coverage. Begin application at bloom and repeat at 7 to 10 day intervals (16 pounds).

Citrus (Grapefruit, Lemons, Oranges, Tangelos, Tangerines): Brown rot (phytophthora)—Use 2 to 4 pounds. Apply to skirts of trees before winter rains. Wet tree thoroughly to height of 3 to 4 feet. Repeat 10 to 12 weeks later (20 pounds) **Oranges and Tangelos:** Melanose, scab—Use 2 pounds plus spray sticker. Apply as necessary from 1/4 petal fall until fruit is 1/2 inch in diameter. Do not apply after fruit size exceeds 1/2 inch in diameter. Do not feed raw by-products to livestock (10 pounds)

Cranberries (Western U.S.): Blotch rot (Acanthorynchus vaccinii), Guignardia blight—Apply 6 pounds per acre in 300 gallons of water in early spring. Repeat applications as required. Storage rots, twig blight—Apply 6 pounds per acre in 300 gallons of water during the first week of June for twig blight and in early and late August for storage rots (7 pounds)

Grapes (Eastern U.S.): Black rot, downy mildew—Use 2 pounds per 100 gallons. Make first application just before bloom. Repeat just after bloom and at 10 to 14 day intervals for 1 to 3 more applications depending upon the susceptibility of the grape variety. (Current season infection)—Use 2 to 4 pounds per 100 gallons dilute spray. Apply 50 gallons per acre when shoots are 1/2 to 1 inch long. Repeat application when shoots are 4 to 6 inches long (10 pounds)

Phomopsis Cane and Leafspot.

Grapes (California): Bunch rot (Botrytis)—Apply 2 pounds per acre. Make 2 applications before bloom and 1 just after bloom. Repeat periodically making 3 cover applications before the bunches close. (Current season infection)—Apply 4 pounds per 100 gallons dilute spray. Apply 3 to 6 pounds per acre by ground equipment. Apply 4 pounds in 7 to 20 gallons of water per acre by aircraft. Apply first spray when green tissue begins to show but before shoots are 1 inch long. Apply a second spray when shoots are 6 to 8 inches long (10 pounds)

Phomopsis cane and Leafspot

Mangoes: Cercospora leaf spot or blotch—Use 2 pounds. Begin at first sign of disease and repeat at weekly intervals to maintain control. (20 pounds)

Peaches, Nectarines: Brown rot, scab—Use 2 pounds in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat application at 7 to 14 day intervals as needed to maintain cover. Continue applications through harvest if conditions favor brown rot. If powdery mildew is a problem, add 3 pounds sulfur to the petal fall, shuck and early cover sprays. If sulfur is added, Captan may be reduced to 1 pound in these sprays. Coryneum blight (peach blight, shot-hole)—Use 2 pounds, apply in pink bud, full bloom, petal fall and cover periods as necessary and as a post harvest spray (but before leaves drop) (10 pounds) **Peaches (Nursery Stock) (California):** Preventive preplant dip treatment for crown gall—Use 4 pounds Captan 50-WP plus 3 2 pints diluted sodium hypochlorite (5 25% household bleach) per 100 gallons water. Wash nursery trees to remove soil from roots. Cut off all dormant buds and suckers in crown area and prune root system if necessary. Submerge the entire dormant tree for 5 minutes. Recharge dip during operation as necessary to maintain a proper concentration of 20 ppm chlorine. Check the concentration frequently using a chlorine test kit.

Pears (Eastern U.S.): Scab—Use 2 pounds in preblossom, petal fall and cover sprays. Repeat cover application at 7 to 14 day intervals under severe scab conditions. Russeting may be reduced on Bosc Pears. Do not use on D'Anjou Pears (16 pounds)

Pears (Western U.S.—except Pacific Northwest): Pear scab—Use 2 pounds. Apply during early finger stage, petal fall and first cover sprays. Under severe scab infection conditions, additional treatments may be necessary. Russeting may be reduced on Bosc Pears. Do not use on D'Anjou Pears (16 pounds)

Pineapples: Heart rot, phytophthora parasitica—Use 10 pounds. Dip slips into solution just prior to planting. Root rot (phytophthora cinnamomi)—Use 4 pounds. Spray immediately after planting. Repeat at weekly intervals. Post Harvest Dip—Use 10 pounds per 100 gallons water to wash harvested fruit.

Plums, Prunes (Eastern U.S.): Brown rot—Use 2 pounds in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain cover. Continue applications through harvest if conditions warrant. The addition of a neutral spray sticker may be necessary to improve coverage.

Plums, Prunes (Western U.S.): Brown rot—Use 2 pounds. Apply at green bud, bloom, prebloom and petal fall stages. Repeat in cover sprays as conditions warrant. Prune russet scab (facy scab) (In California)—Apply 2 pounds per 100 gallons water (pounds per acre) to cover sprays.

BEST DOCUMENT AVAILABLE

1950

Quince: Brown rot, scab—Use 2 pounds. Begin when disease threatens. Repeat at 7 to 10 day intervals. Do not apply within 7 days of harvest. (10 pounds)

Strawberries: Botrytis (gray mold), leaf spot—Apply 4 to 6 pounds per acre in sufficient water for thorough coverage. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at weekly intervals. Under conditions favorable to fruit rot continue applications through harvest period treating immediately after each picking. (8 pounds)

VEGETABLE CROPS

Asparagus (Preplanting Root Dip): Botrytis, root rot, phoma rot, penicillium rot, fusarium—Use 3 pounds per 100 gallons water (4 tablespoonfuls per gallon). Dip root sections for 2 minutes, drain and plant.

Beans, Peas: Damping-off (Beans only), fungus root rot—Use 10 to 12 pounds per acre as a broadcast treatment, working into top 3 to 4 inches of soil, or use 5 to 6 pounds per acre in the row at time of planting.

Cantaloupes, Cucumbers and Pumpkins, Summer and Winter Squash, Honeydew Melons, Watermelons: Angular leafspot, Anthracnose, Downy Mildew - use 4 lbs. per acre in sufficient water for thorough coverage. Begin applications at first signs of infection and repeat at 5-7 day intervals or as needed to maintain control. (7.6 pounds)

Carrots: Cercospora blight, septoria leaf spot—Use 2 pounds per 100 gallons water and apply the equivalent of 4 pounds Captan 50-WP per acre. Apply at first sign of disease and repeat at 6 to 10 day intervals. (6 pounds)

Celery: Late Blight (Septoria), Pink Rot (Sclerotinia) - 2 lbs. per 100 gallons spray. Apply 200 diluted gallons per acre or equivalent of 4 to 4-1/2 lbs. per acre in water for coverage. Apply at first signs of diseases. Repeat at 7 to 10 day intervals as necessary to maintain control.

Eggplant: Phomopsis blight, fruit rot, anthracnose—Use 2 pounds. Begin when plants emerge in plant bed or when disease first appears in field. Repeat at 3 to 7 day intervals through growing season. Damping-off—Use 10 to 12 pounds per acre in sufficient water for thorough coverage. Begin when all seeds have germinated. Repeat at 10 day intervals. (6 pounds as foliar spray, 12 pounds in plant beds before transplanting)

Lettuce, spinach: Downy mildew - 2 lbs. per 100 gals. spray. Apply 100-200 gals. of spray per acre. Apply at first signs of disease. Repeat at 7-10 day intervals as necessary to maintain control.

Onions (Green) (California): Downy mildew - Use 3 pounds per acre of 4 to 20 gallons water per acre by a sprayer or 20 to 100 gallons water per acre by ground equipment. Apply at first signs of infection and repeat at 7 to 10 day intervals as necessary to maintain disease control. (3 pounds)

Onions (Green and Bulb): Downy mildew, purple blotch—Use 2 pounds. Apply at first sign of infection and repeat at 7 to 10 day intervals. (2 pounds)

Peppers, Pimentos: Anthracnose, blight, purple blotch, leaf and stem rot—Use 3 pounds. Apply at 3 to 7 day intervals throughout the growing season. (6 pounds). Damping-off—Use 2 pounds. Plant root treatment. Begin when all seeds have germinated. Repeat at 10 day intervals. (2 pounds)

BEST AVAILABLE

1950

319-3019

Tomatoes: Anthracnose, gray leaf spot, septoria leaf spot—Use 4 to 6 pounds per acre in sufficient water for thorough coverage. Under severe infection conditions use 6 to 8 pounds per acre in sufficient water for thorough coverage. Apply at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control (10 pounds).

SPECIAL USES

Preplanting Soil Treatment: For control of damping-off and root rots for the crops listed below, apply spray broadcast to soil surface as a preplanting application, and cultivate into upper 3 to 4 inches of soil. Or apply to furrow over the seed when specified (14,500 linear feet of row equals 1 acre.) Close and pack the furrow in usual manner. Apply dosage rate in sufficient water for thorough soil coverage unless otherwise specified.

Beans, Peas: Use 10 to 12 pounds broadcast or 5 to 6 pounds in furrow (12 pounds)

Beets: Use 2 pounds per 100 gallons water, 15 gallons spray per 1,000 square feet (13 pounds)

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Kohlrabi, Mustard, Rutabagas, Turnips: Damping-off only—Use 15 pounds broadcast (28 pounds)

Cantaloupes, Cucumbers, Honeydew Melons, Pumpkins, Squash (Summer and Winter), Watermelons: Use 2 pounds per 100 gallons water, 600 gallons spray per acre broadcast (12 pounds)

Celery Seedbed Treatment: Damp Off, Pink Rot; Peppers: Damp Off - 2 lbs. per 100 gallons spray. Spray or drench at 10 to 15 gallons per 1,000 square feet of seedbed. Apply when all seeds have germinated. Repeat as necessary.

Spinach (Direct seeded): Use 5 to 7 pounds in 30 gallons water in furrow (12 pounds)

Post Harvest Dip: For control of various mold and storage rots (Botrytis, Geosporium, Rhizopus)—Use as a post harvest dip or wash on the following fruits and vegetables: Apples, Apricots, Cantaloupes, Cucumbers, Grapes, Lemons, Limes, Mangoes, Nectarines, Onions (Green and Bulb), Oranges, Peaches, Pears, Potatoes, Tangerines, Tangerines—Use 2 1/2 pounds. Apply as a spray or in dip tank. When used as a dip, recharge wash solution period daily when tank volume is reduced by 25%. Bring water back to volume and add 1 pound for each 25 gallons added. At end of every 8 hour shift, empty tank, flush and charge with fresh solution. Do not allow dip tank to stand overnight. Maintain continuous agitation during operation. Do not use on leafy greens, Mangos or Citrus by-products to dairy animals or animals being finished for slaughter.

ORNAMENTALS

Azaleas: Damping off of cuttings—Use 4 pounds. Dip cuttings before bedding. Repeat at 7 to 14 day intervals through bloom.

Begonias (Tuberous): Damping off tuber rot—Use 4 pounds. Dip tubers before bedding.

Camellias: Botrytis flower blight—Use 2 pounds. Apply to first signs of infection. Repeat at 7 to 10 day intervals through bloom.

Carnations: Botrytis flower blight—Use 2 pounds. Begin at first signs of disease. Repeat at 7 to 10 day intervals. Shorten intervals to 5 to 7 days if necessary. Damping off of cuttings—Use 4 pounds. Dip cuttings before bedding.

Chrysanthemum: Botrytis flower blight, Septoria leaf spot—Use 2 pounds. Apply at first signs of disease. Repeat at 7 to 10 day intervals. Damping off of cuttings—Use 4 pounds. Dip cuttings before bedding.

Dianthus (California Only): Botrytis flower blight—Use 2 pounds. Apply at first signs of disease. Repeat at 7 to 10 day intervals.

BEST DOCUMENT AVAILABLE

7 3019

Gladolus (Corms): Corm rot and decay, damping-off—Use $\frac{1}{4}$ to $1\frac{1}{4}$ pounds per 10 gallons, dip corms 20 to 30 minutes. Drain and plant.

Grasses (Ornamental in Non-Pastured Areas Only): Leaf spot, damping-off, brown patch, melting out, seedling blights, and brown spot on St. Augustine grass—Use 2 pounds. Apply 10 gallons spray per 1,000 square feet. Begin when growth starts in spring. Repeat at 7 to 14 day intervals throughout season. Do not graze treated areas or feed clippings to livestock.

Grasses (Lawn Seedbeds): Damping-off and other soil borne diseases—Use $\frac{1}{4}$ to $\frac{1}{2}$ pound per 1,000 square feet or 2 pounds in 100 gallons water using 15 gallons spray per 1,000 square feet. Cultivate into upper 3 to 4 inches before planting.

Roses: Black spot, Botrytis blossom blight—Use 2 pounds. Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals, and more frequently during frequent rains and heavy dews.

Soil and Greenhouse Bench Treatment: Preplanting for damping-off and root rot diseases on seedlings or transplants of roses (and other shrubs, trees, flowers) Lawn seedbeds, vegetables (such as beans, celery, crucifers, eggplant, peas, peppers, tomatoes)—Use 2 pounds per 100 gallons water at rate of 15 gallons per 1,000 square feet. Cultivate into upper 3 to 4 inches of soil before planting.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

DOCUMENT AVAILABLE

FMC 2011 No. 279-3111
12/13/11

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

1987 Corporation
Agricultural Chemical Division
2000 Market Street
Philadelphia, PA 19103

Attention: J. M. Currie

1/3/84

Gentlemen:

Subject: Captain 50-W
EPA Registration No. 279-6019
Captain 7.5 Dust
EPA Registration No. 279-1649
Your letter dated January 25, 1984

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted at the next printing.

- 1. Insert the heading "Reentry Statement" above pertinent reentry information.
- 2. Add "Do not apply directly to water or wetlands" to the environmental hazard statement in 279-6019. Correct the spelling of "or" in the same statement in 279-1649.

Three copies of the label are enclosed for your records.

Sincerely,
HP

Walter M. Jandy
Regional Manager (A)
Pesticide Registration Branch
Registration Division, EPA-WPHQ

BEST DOCUMENT AVAILABLE

Enclosure
EPA-279-6019-01501-01502-01503-01504-01505-01506-01507-01508-01509-01510-01511-01512-01513-01514-01515-01516-01517-01518-01519-01520-01521-01522-01523-01524-01525-01526-01527-01528-01529-01530-01531-01532-01533-01534-01535-01536-01537-01538-01539-01540-01541-01542-01543-01544-01545-01546-01547-01548-01549-01550-01551-01552-01553-01554-01555-01556-01557-01558-01559-01560-01561-01562-01563-01564-01565-01566-01567-01568-01569-01570-01571-01572-01573-01574-01575-01576-01577-01578-01579-01580-01581-01582-01583-01584-01585-01586-01587-01588-01589-01590-01591-01592-01593-01594-01595-01596-01597-01598-01599-01600

SYMBOL ▶
SURNAME ▶
DATE ▶
EPA

CONCURRENCES

[Empty box for CONCURRENCES]

OFFICIAL FILE COPY