

Code 3669

RESTRICTED USE PESTICIDE

Net Weight

Due To Avian Toxicity

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification

Furadan® 15 G Insecticide-Nematicide For Agricultural or Commercial Use Only

EPA Reg. No. 279-3023

EPA Est. 279-

Active Ingredient:

*Carbofuran 15.0%

Inert Ingredients: 85.0%

100.0%

*2,3-Dihydro-2,2-dimethyl-7-benzofuroxylmethylcarbamate

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

If **swallowed**: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in **eyes**: Flush with plenty of water for at least 15 minutes. Get medical attention.

If on **skin**: Wash immediately with soap and water. Remove contaminated clothing and wash before reuse.

Antidote

Note to Physician: Carbofuran is a reversible cholinesterase inhibitor. Do not use oximes such as 2-PAM. Start by giving 2 mg. atropine intramuscularly. According to clinical response, continue until signs of atropinization occur (dry mouth or dilated pupils). If in eye, instill one drop of homatropine.

For Emergency Assistance Call (800)-331-3148.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

Poisonous if swallowed. May be fatal or harmful as a result of skin or eye contact or by breathing dust. Causes cholinesterase inhibition. Warning symptoms of poisoning include weakness, headache, sweating, nausea, vomiting, diarrhea, tightness in chest, blurred vision, pinpoint eye pupils, abnormal flow of saliva, abdominal cramps, and unconsciousness. Atropine sulfate is antidotal.

Wear long-sleeved clothing and protective gloves when handling. Wash hands and face before eating or smoking. Bathe at the end of the work day. Change clothing daily and wash before reuse.

See other panels for additional precautionary information.

Environmental Hazards

This pesticide is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Birds killed by carbofuran pose a hazard to hawks and other birds-of-prey; contact your state wildlife agency or the U.S. Fish and Wildlife Service upon finding birds or other animals that may have been poisoned. Cover or incorporate granules in spill areas. Runoff from treated areas may be hazardous to fish in neighboring areas. Do not apply directly to water. Do not contaminate wells, wetlands or any body of water by cleaning of equipment or disposal of waste.

Notice: It is a Federal offense to use any pesticide in a manner that results in the death of a member of an endangered species.

The use of Furadan 15 G may pose a hazard to the following Federally designated endangered/threatened species known to be found in certain areas within the named locations.

Attwater's Greater Prairie Chicken—Texas counties including: Aransas, Austin, Brazoria, Colorado, Galveston, Goliad, Harris, Refugio and Victoria

Aleutian Canada Goose—California counties including Colusa, Merced, Stanislaus and Sutter

Kern Primrose Sphinx Moth—Walker Basin of Kern County, California.

This product may not be used in areas where adverse impact on the Federally designated endangered/threatened species, noted above, is likely. Prior to making applications, the user of this product must determine that no such species are located in or immediately adjacent to the area to be treated. If the user is in doubt whether or not the above named endangered species may be affected, he should contact either the regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or personnel of the State Fish and Game office.

Carbofuran is a chemical which can travel (seep or leach) through soil and can contaminate ground water which may be used as drinking water. Carbofuran has been found in ground water as a result of agricultural use. Users are advised not to apply carbofuran where the water table (ground water) is close to the surface and where the soils are very permeable, i.e., well-drained soils such as loamy sands. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use this product on Long Island, N.Y. Do not use this product on corn or sorghum after 8/31/94.

STORAGE AND DISPOSAL

Pesticide Storage

Do not store in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC. (800) 331-3148.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.


Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FMC.

FMC Corporation
Agricultural Chemical Group
Philadelphia PA 19103

3/93



BEST AVAILABLE COPY

General Use Instructions

If prolonged intimate contact with corn and/or sorghum foliage will result, do not reenter treated fields within 14 days of application without wearing proper protective clothing. For all other foliar application situations, do not reenter fields less than 24 hours following application without wearing proper protective clothing.

Cover or incorporate granules in spill areas.

Do not plant with any crop other than alfalfa, artichokes, bananas, barley, coffee, corn (field pop or sweet), cotton, cranberries, cucurbits (cucumbers, melons, pumpkins, squash), flax, grapes, non-bearing fruit, oats, ornamentals, peanuts, peppers, potatoes, rice, seed crops (Bermuda-grass, spinach), sorghum, strawberries, soybeans, sugar beets, sugarcane, sunflowers, tobacco, or wheat for at least 10 months following use of this product.

Cranberries—Washington Only

Insects Controlled	Rate of Application	Method of Application
Black Vine Weevil	13.3 Pounds per acre	Spread granules uniformly over soil surface using a rotary spreader. Make one application about June 15 to 25 and repeat one month later. Incorporate granules by following each application with sprinkler irrigation.

For use only on cranberry bogs that are sprinkler irrigated and are not flooded for purposes of irrigation or harvest.

Do not apply within 60 days of harvest.

Cucurbits (Cucumber, Melons, Squash, Pumpkins)—Southeastern States Only

Insects Controlled	Rate of Application	Method of Application
Nematodes (Dagger, Lesion and Root-Knot)	1½ pounds per 1,000 linear feet of row	Apply in a 12 to 15 inch band and incorporate into the top 3 inches of soil.

Pine Seedlings in Pine Progeny Tests

Insects Controlled	Rate of Application	Method of Application
Pales Weevil Pitch-Eating Weevil Nantucket Pine Tip Moth Pitch Pine Tip Moth	7 grams (¾ teaspoon) per seedling	Apply at transplanting. Distribute the granules in a 6 inch radius on the soil around each seedling. Cover granules with soil.

Do not use at transplanting if seedlings were treated with Furadan 4F slurry root dip.

Note: FMC will sell this product in conformity to volume limitations agreed to with EPA.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Furadan and —FMC Trademarks

(3669-10/2/92-A)

BEST AVAILABLE COPY

