

Code 3876

Net Weight  
Pounds

# Kolodust® 60

## Fungicide-Miticide

EPA Reg. No. 279-3024      EPA Est. 279-

Active Ingredient:	
Sulfur.....	60%
Inert Ingredients.....	40%
	100%

Sulfur fineness 97.0% through 325 mesh  
Kolodust® - FMC Trademark

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

For Emergency Assistance Call 716-735-3785.

See Other Panels for Additional Precautionary Information



FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans (& Domestic Animals)

##### Caution

Avoid contact with skin, eyes or clothing. Avoid breathing vapors, dust or spray mist. May be harmful if swallowed. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

#### Environmental Hazards

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not reuse container. Dispose of container in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

Some fruits and other plants are susceptible to injury from sulfur under certain climate conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when the temperature is high (90° or above). Do not make sulfur applications at such times.

Do not use within 2 weeks of an oil spray treatment. When growing crops for processing, consult the processor before applying sulfur.

Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with label cautions and directions.

This material may be applied with ground equipment or aircraft.

Where recommended for prevention of Powdery Mildew, apply when disease first appears and repeat at 7 to 10 day intervals throughout the growing season. Shorten the interval during humid rainy weather.

Where recommended for prevention of Brown Rot, apply at pink bud or popcorn stage during bloom and at petal fall, shuck split and shuck fall. Repeat at 10 to 14 day intervals in cover sprays as long as necessary.

Where recommended for control of Mites, apply when insects first appear and repeat as necessary throughout the growing season.



**Allalfa:** For control of Atlantic Mite, Spider Mite and Clover Mite—Use 30 to 50 lbs. per acre.

**Beans:** For the control of Pacific Mite (Red Spider) Use 25 lbs. per acre. For control of Powdery Mildew—Use 40 to 50 lbs. per acre. Certain varieties of beans may be insured by sulfur.

**Corn:** For the control of Spider Mites—Use 30 to 50 lbs. per acre.

**Cotton:** For control of Mites—Apply 10 to 50 lbs. per acre.

**Grapes:** For the control of Powdery Mildew—Use 8 to 20 lbs. per acre. Make first application when shoots are 6 inches to 8 inches long. Make a second application when shoots are 12 inches to 15 inches long and the third about 14 days later. Repeat applications at 10 to 14 day intervals to protect new growth. Avoid applications during hot weather as injury may occur. Concord and other *labrusca* type grapes may be injured by sulfur.

**Hops:** For control of Red Spider Mites—Use 30 to 50 lbs. per acre.

**Peaches:** For the control of Peach Silver Mite, Powdery Mildew, and prevention of Brown Rot of fruit—Apply 40 to 50 lbs. per acre. For prevention of Brown Rot of fruit—Apply when fruit begins to ripen.

**Plums and Prunes:** For prevention of Brown Rot of fruit—Appl. 40 to 50 lbs. per acre. Apply when fruit starts to ripen.

**Sugar Beets:** For control of Spider Mites and Powdery Mildew—Apply 60 to 80 lbs. per acre.

**Tomatoes:** Tomato Russet Mite, Powdery Mildew—Use 40 to 55 lbs. per acre.

### **Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.