

Code

Net Weight  
Pounds

# Kolodust® 60

## Fungicide-Miticide

EPA Reg. No. 279-3024

EPA Est., 279-

**Active Ingredient:**

Sulfur.....	60%
Inert Ingredients: .....	40%
	<u>100%</u>


Sulfur fineness 97.0% through 325 mesh.  
Kolodust®—FMC Trademark

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

For Emergency Assistance Call 718-735-3765.

See Other Panels for Additional Precautionary Information.



# FMC

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

PA 1/81

**ACCEPTED**

FEB 25 1981

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 279-3024

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans (& Domestic Animals)

Avoid contact with skin, eyes or clothing. Avoid breathing vapors, dust or spray mist. May be harmful if swallowed. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

#### Environmental Hazards

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Do not reuse container. Dispose of container in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

Some fruits and other plants are susceptible to injury from sulfur under certain climate conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Sulfur may burn foliage when the temperature is high (80° or above). Do not make sulfur applications at such times.

Do not use within 2 weeks of an oil spray treatment. When growing crops for processing, consult the processor before applying sulfur.

Consult your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information, consistent with label cautions and directions.

This material may be applied with ground equipment or aircraft.

Where recommended for prevention of Powdery Mildew, apply when disease first appears and repeat at 7 to 10 day intervals throughout the growing season. Shorten the interval during humid rainy weather.

Where recommended for prevention of Brown Rot, apply at pink bud or popcorn stage during bloom and at petal fall, shuck split and shuck fall. Repeat at 10 to 14 day intervals in cover sprays as long as necessary.

Where recommended for control of Mites, apply when insects first appear and repeat as necessary throughout the growing season.