

Code 180

Net Contents: 5 Gallons


Liqui-Stik 200  
Growth Regulator

EPA Reg. No. 279-

EPA Est., 279-CA-2

	By Wt.
Active Ingredient:	
1-Naphthaleneacetic Acid, sodium salt* . . .	5.85%
Inert Ingredients: . . . . .	94.15%
	<u>100.00%</u>

\*Equivalent to 5.2% 1-Naphthaleneacetic acid or 200 grams per gallon



KEEP OUT OF REACH OF CHILDREN

CAUTION

FOR EMERGENCY ASSISTANCE CALL 716-735-3765

See other panels for additional precautionary information.

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia, PA 19103  
(PA 10/82)

CODE 180  
EPA Est. 279-C-2

#### PRECAUTIONARY STATEMENTS

##### Environmental Hazards

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

##### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act. To dispose of container, triple rinse and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved State and local procedures.

Liqui-Stik 200 is a hormone-like spray to be used in the following manner on apples and pears for the prevention of pre-harvest fruit drop and the thinning of olives. Do not use in undiluted form:

##### Directions for Apples and Pears

Spray trees thoroughly, wetting all fruit and adjacent leaves. Apply 7 to 14 days before harvest of apples and 5 to 7 days before harvest of pears. This product becomes effective in 2 to 3 days after application and controls drop for 10 days to 4 weeks (less for McIntosh variety) depending on weather conditions. Do not make more than 2 applications and do not delay picking beyond optimum maturity.

Ground Spray: Apples - Thoroughly mix 12 fluid ounces of Liqui-Stik 200 with 250 gallons of water (4.8 fluid ounces per 100 gallons water) to prepare a 20 ppm spray. Application of 630 gallons per acre of this 20 ppm spray provides an equivalent of 48 grams of 1-Naphthaleneacetic acid per acre. Pears - use a 10 ppm spray obtained by mixing 12 fluid ounces with 500 gallons of water. Application of 670 gallons per acre of this 10 ppm spray provides an equivalent of 25 grams of 1-Naphthaleneacetic acid per acre.

CODE 160  
LFP. ESP. 979-CA-2

Aerial Application: Apples - Use 30 fluid ounces in 5 gallons of water. Pears - Use 11 fluid ounces of this product in 5 gallons of water per acre.

Directions for Thinning Olive Trees

Liqui-Stik 200 is effective in thinning olive trees when an excessively heavy fruit set is expected.

Timing is critical. Spray when young fruits are 1/8 to 3/16 inches in diameter, which is from 1 1/2 to 2 1/2 weeks following full bloom. Thinning is effective on all varieties except Sevillano. Without Oil. Apply a solution of 150 ppm. 1-Naphthaleneacetic acid equivalent obtained by adding 9 fluid ounces and a wetting agent (e.g. Dreft 3 to 4 ounces) to 25 gallons of water. With Oil. Apply a solution of 100 ppm. 1-Naphthaleneacetic acid equivalent obtained by mixing thoroughly 6 fluid ounces with 25 gallons of water. Emulsify by agitation with 3 pints of light or medium summer oil. Spray only the heavily loaded trees until the leaves are thoroughly wet. This will require 10 to 30 gallons per tree. CAUTION: Unless you wish to remove all your fruit, do not apply during bloom.

Dealer Should Sell in Original Packages Only

Terms of Sale or Use On purchase of this product, buyer and user agree to the following conditions:

Warranty FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

0.1. 180  
EPA EST. 9279-CA-2

Damages Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

FMC<sup>R</sup> is a registered trademark of FMC Corporation.