

Code _____

_____ lbs. Net Weight

RESTRICTED USE PESTICIDE

For retail sale to, and use only by, Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Pounce^R 25 WP
Insecticide

EPA Reg. No. 279-_____

EPA Est. 279-_____

	<u>By Wt.</u>
Active Ingredients	
*Permethrin**.....	25.0%
Inert Ingredients	75.0%
	<u>100.0%</u>

* (3-Phenoxyphenyl)methyl(+) cis-trans 3-(2,2-dichloro-ethenyl)-2,2-dimethylcyclopropanecarboxylate
**cis/trans ratio: Max. 55% (+) cis and min. 45% (+) trans

U.S. Patent No. 4,024,163

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

DECI 1984

Insecticide,
Fungicide,
Herbicide
EPA Reg. No.
279-3051

For Emergency Assistance Call 716-735-3765

See other panels for additional precautionary information.

FMC.

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

r711A20
ds21

Pounce 25 WP (Draft)
6/84
Page 1 of 7

BEST AVAILABLE COPY

Written warnings cannot be understood by workers. Written warnings must include the following information:

WARNING

Area treated with Pounce 25 WP on (insert date). Do not enter without appropriate protective clothing until sprays have dried (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidents: exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

Storage and Disposal

Pesticide Storage

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Instructions

Pounce 25 WP is a 25% formulation of the insecticide permethrin. Apply Pounce 25 WP when insects appear or feeding is noticed. The higher rate should be used as pest populations increase. Repeat the application as necessary to maintain control. Pounce 25 WP may be applied by both ground and aerial equipment. Use sufficient water to obtain full coverage. With the exception of crops listed below, rotational crops should not be planted within 60 days of last application.

Mix as needed; do not store diluted material.

BEST AVAILABLE COPY

Pounce 25 WP (Draft)
6/84
Page 3 of 7

PM17
PS 5000

Almonds: To control Navel Orangeworm, Peach Twig Borer - Use Pounce 25 WP at a rate of 12.8-25.6 ounces (0.2 to 0.4 pounds active) per acre. Apply when insects appear. Do not make more than two applications during hull split. Do not make more than 5 applications per season. Use sufficient water to obtain full coverage of foliage. Do not apply within seven days of harvest. Do not graze livestock in treated orchards.

Apples: To control Plum Curculio, Redbanded Leafroller, Rosy Apple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Oblique Banded Leafroller and Green Fruitworm - Use Pounce 25 WP at a rate of 6.4-12.8 ounces (0.1 to 0.2 pounds active) per acre with ground equipment only. Apply in 25 to 400 gallons of finished spray per acre when insects appear and repeat as required to maintain control. Do not make more than three applications per season. Do not apply after petal fall.

Broccoli, Brussels Sprouts, Cauliflower: To control Beet Armyworm, Cabbage Looper, Imported Cabbageworm and Diamondback Moth - Use Pounce 25 WP at a rate of 3.2-6.4 ounces (0.05 to 0.1 pound active) per acre. Use sufficient water to obtain full coverage of foliage. Do not make more than 8 applications per season. Do not apply within one day of harvest.

Cabbage: To control Beet Armyworm, Cabbage Looper, Diamondback Moth and Imported Cabbageworm - Use Pounce 25 WP at a rate of 3.2 to 6.4 ounces (0.05 to 0.1 pound active) per acre. Do not make more than 10 applications per season. Do not apply within one day of harvest.

Celery: To control Leafminer, Cutworms and Southern Armyworm - Use Pounce at a rate of 6.4 to 12.8 ounces (0.1 to 0.2 pound active) per acre on a 3 to 5 day schedule, or as needed to maintain control. Use sufficient water to obtain full coverage of foliage. Do not make more than 10 applications per season in California. Do not apply within 3 days of harvest. Do not graze celery forage nor feed trimmings from treated fields to livestock.

BEST AVAILABLE COPY

Pounce 25 WP (Draft)
6/84
Page 4 of 7

17
156018

Cotton: To control Beet Armyworm (California and Arizona only), Bollworm, Cabbage Looper, Cotton Fleahopper, Cotton Leaf Perforator, Lygus Bugs, Pink Bollworm, Tarnished Plant Bug and Tobacco Budworm - Use Pounce 25 WP at a rate of 6.4 to 12.8 (0.1 to 0.2 pound active) per acre. To control Boll Weevil, alternate Pounce 25 WP at 6.4 ounces (0.1 pound active) per acre with Pounce 25 WP at 3.2 ounces (0.05 pound active) per acre every 3-4 days until control is achieved.

Use sufficient water to obtain full coverage of foliage.

Do not make more than 15 applications per season. Do not apply within 14 days of harvest. Do not graze or feed cotton forage.

Head Lettuce: To control Alfalfa Looper and Cabbage Looper - Use Pounce 25 WP at a rate of 3.2 to 12.8 ounces (0.05 to 0.2 pound active) per acre. To control Beet Armyworm, Corn Earworm and Tobacco Budworm - Use Pounce 25 WP at a rate of 6.4 to 12.8 ounces (0.1 to 0.2 pounds active) per acre. Use sufficient water to obtain full coverage of foliage. Do not make more than 15 applications per season.

Peaches: To control Lesser Peach Tree Borer, Peach Twig Borer, Rose Chafer, Plum Curculio, Tarnished Plant Bug, Green Fruitworm, Oriental Fruit Moth - Use Pounce 25 WP at a rate of 3.2-6.4 ounces (0.05-0.1 pound active) per 100 gallons of water. Do not exceed 400 gallons per acre. Apply when insects appear. Make applications with ground equipment, using sufficient water to obtain uniform coverage. Do not apply within 7 days of harvest. Do not make more than 8 applications per season. Do not graze livestock in treated orchards.

Pears (Dormant through Delayed Dormant): To control Pear Psylla - Apply Pounce 25 WP at a rate of 12.8 to 25.6 ounces (0.2 to 0.4 pound active) per acre. Pounce 25 WP may be combined with 2 to 8 gallons of spray oil per acre. Apply during the dormant through delayed dormant growth periods only. Apply in 3 to 20 gallons of finished spray per acre by aircraft and 25 to 400 gallons per acre by ground equipment. Do not make more than two applications per season. Do not graze orchard floor.

Pears (Summer): To control Pear Psylla, Green Fruitworm, Codling Moths - Use Pounce 25 WP at a rate of 12.8 to 25.6 ounces (0.2-0.4 pound active) per acre. Use in sufficient water to achieve runoff and apply by ground equipment. Do not make more than 3 applications per season. Do not apply within 14 days of harvest.

Pounce 25 WP (Draft)
6/84
Page 5 of 7

BEST AVAILABLE COPY

1 m 17
28 7.84

Potatoes: To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Cutworms, European Corn Borer, Potato Flea Beetle, Potato Leafhopper, Potato Tuberworm, Aster Leafhopper and Tarnished Plant Bug. Use Pounce 25 WP at a rate of 6.4 to 12.8 ounces (0.1 to 0.2 pound active) per acre. Use sufficient water to obtain full coverage of foliage. Do not make more than 12 applications per season. Do not apply within 7 days of harvest. Do not graze or feed potato forage.

Soybeans: To control Cabbage Looper, Corn Earworm, Green Cloverworm, Mexican Bean Beetle, Soybean Looper, Velvetbean Caterpillar, Bean Leaf Beetle and Potato Leafhopper - Use Pounce 25 WP at 3.2 to 6.4 ounces (0.05 to 0.1 pound active) per acre. To control Beet Armyworm - Use Pounce 25 WP at a rate of 6.4 to 12.8 oz. (0.1 to 0.2 lbs. active) per acre. Use higher dosage rate as population pressure increases. Use sufficient water to obtain full coverage of foliage. Do not make more than two applications per season. Do not apply within 60 days of harvest. Do not graze or feed soybean forage. Do not plant rotational crops within 60 days of last application.

Spinach: To control Beet Armyworm, Corn Earworm, Cabbage Looper, Cutworms, European Corn Borer, Green Cloverworm, Leafhoppers - Use Pounce 25 WP at a rate of 6.4 ounces (0.1 pound active) per acre. Apply when insects appear. Use sufficient water to obtain full coverage of foliage. Do not make more than 7 applications per season. Do not apply within 7 days of harvest.

Sweet Corn: To control Corn Earworm, European Corn Borer, Fall Armyworm - Use Pounce 25 WP at a rate of 6.4-12.8 ounces (0.1-0.2 pound active) per acre. Apply every 5 days or as needed. Do not apply more than 6 applications. Do not apply within 1 day of harvest of ears or forage or livestock grazing.

Sweet Corn (Florida Only) fresh market use only: To control Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm - Use 6.4-16 ounces (0.1-0.25 pound active) per acre. Apply every 5 days or as needed.

Do not apply more than 3.25 pounds active ingredient per acre per crop season. Do not graze livestock in treated areas or harvest for forage.

r722B27
rk30

BEST AVAILABLE COPY

Pounce 25 WP (Dropt)
6/84
Page 6 of 7

