



**STORAGE AND DISPOSAL**

**Pesticide Storage**

Do not freeze. Do not store below 40°F. Solid crystals are observed, warm material to above 60°F by placing container in warm location. Shake or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC Collect (716) 735-3765.

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal**

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal**

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Empty Containers: Do not rinse container or other empty remaining formulated product. Do not break seals. Return waste to point of purchase.

**IMPORTANT**

**FAILURE TO OBSERVE THE APPLICATION PRECAUTIONS SECTION OF THIS LABEL MAY RESULT IN INJURY TO DESIRABLE VEGETATION**

- Desirable plants including some species of trees, shrubs, flowers, agronomic crops, and fruits and vegetables are sensitive to Command 4 herbicide.
- Foliar contact with spray drift or vapors may cause whitening of sensitive plants. Symptoms are generally temporary in nature but may persist on some plants.

**SPRAYER CLEANUP**

Do not drain or flush equipment on or near desirable trees or other plants, or in areas where their roots may extend or in locations where the chemical may be washed or move into contact with their roots. Do not contaminate any body of water including irrigation water that may be used on other crops. Carefully follow sprayer clean-up instructions noted below to prevent spray tank residues from damaging other crops.

Sprayer equipment should be thoroughly rinsed to remove residues of herbicide that might injure other subsequently sprayed crops. The steps below should be followed for the thorough rinsing of spray equipment following applications of Command 4 EC herbicide or tank mixes of Command 4 EC with other labeled products.

- 1) Drain any remaining spray solution from tank, pump, hoses and boom and discard in an approved manner (See Note that follows).
- 2) Clean tank and fittings by
  - Thoroughly hosing down the inside walls of the spray tank with a quantity of water equal to 1/2 of the total tank capacity and operating the pump to circulate this solution through the sprayer system for 15 minutes.
  - Washing down the outside surfaces of equipment.
  - Removing nozzle tip and screen from end nozzle in each boom section and allowing several gallons of rinsate solution to flush completely through boom (collect rinsate while flushing).
- 3) Thoroughly drain remaining rinsate solution from tank, pump and hoses. Combine with boom flushings and dispose of all rinsates from this first rinsing in an approved manner (see Note that follows).
  - When switching from water dilutions to applications utilizing crop oil or liquid fertilizer as a carrier, a small volume of crop oil or liquid fertilizer should be flushed through the tank, pump, hoses and boom prior to the next use. Dispose of crop oil or liquid fertilizer rinsate in an approved manner (see Note for local, state and Federal guidelines).
- 4) Remove the remaining nozzle tips and screens and the line filter and wash in a pail of warm soapy water, thoroughly hose and replace.
- 5) Hose down the inside walls of the spray tank a second time and circulate this solution using the same procedure as noted in 2) above.

6) If the next use of the sprayer will be for applying a preemergent or preplant incorporated pesticide on any crop for which Command 4 EC is registered, rinsate from this second rinsing may be utilized by diluting with the water for the next pesticide load.

**HOWEVER**

If the next use of the sprayer will be a postemergence applied pesticide on any crop, drain rinsate solution from this second rinsing. Retain rinsate solution for use only with a soil incorporated pesticide to be applied on a crop for which Command 4 EC is registered. Refill tank (after draining second rinsate solution) in accordance with post-emergence product label directions.

**NOTE:** Dispose of excess spray mixture and/or rinsate from first tank rinsing by application to cropland as described on this label. If excess spray mixture and/or rinsate from first rinsing cannot be disposed of according to label instructions, dispose of in compliance with local, state and Federal guidelines. Contact your state pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance.

**GENERAL MIXING INSTRUCTIONS**

Care must be taken when mixing Command 4 EC herbicide. Avoid mixing in areas adjacent to desirable plants (See Section headed "IMPORTANT").

**Command 4 EC Alone:** Mix Command 4 EC with water or liquid fertilizer in the following manner: Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 4 EC, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

**Tank Mixtures:** Fill spray tank one-fourth to one-third full with water; with agitator operating add the recommended amount of ingredients using the following order: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's). Mix thoroughly and fill tank one-half full continuing agitation. Add Command 4 EC herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Where use of a surfactant is recommended, add as the last ingredient to the spray tank. Maintain agitation during filling, mixing and application. When using drift reducing agents, follow specific product label instructions for order of addition to spray tank.

**Fertilizer Spray Mixtures:** Applications of Command 4 EC alone, or with recommended tank mixtures, in conjunction with fertilizer solutions may be used unless directions specifically state otherwise. Small quantities should be tested for compatibility by the following procedure before mixing tankful quantities.

- 1) Put 1 pint of fertilizer solution in a quart jar.
- 2) Add the appropriate amount of herbicide based on the tables below. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).

| Herbicide                    | Rate/Acre | Amount To Be Added Per Pint of Fluid Fertilizer* |
|------------------------------|-----------|--|
| Wettable Powders             | 1/2 pound | 3/4 level teaspoon                               |
| or                           | 1 pound   | 1 1/2 level teaspoons                            |
| Dry Flowables (Dry Granules) | 2 pounds  | 3 level teaspoons                                |
|                              | 3 pounds  | 4 1/2 level teaspoons                            |
| Emulsifiable Concentrates    | 1 pint    | 1/2 level teaspoon                               |
| or                           | 1 quart   | 1 level teaspoon                                 |
| Flowables                    | 2 quarts  | 2 level teaspoons                                |
|                              | 3 quarts  | 3 level teaspoons                                |

\*Based on a spray volume of 25 gpa. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

- 3) Close jar and shake well.
- 4) Watch mixture for several seconds, again in 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed.


If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below.

- 1) **Command 4 EC Alone:** Dilute with 2 parts of water and add slowly to spray tank of fertilizer with agitator running. Maintain agitation during application.
- 2) **Command 4 EC plus Tank Mixtures:** Dilute individual products with two (2) parts of water, then add to the spray tank of fertilizer while maintaining agitation using the following order—slurry of dry formulations (wetable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Continue agitation during application.

# APPLICATION PRECAUTIONS

**GEOGRAPHIC RESTRICTIONS:** Command 4 EC must be applied only as a soil incorporated treatment in the following states, unless there is an approved SLN (24C) for counties within these states:

|          |                |               |
|----------|----------------|---------------|
| Delaware | Michigan       | New York      |
| Illinois | Minnesota      | Ohio          |
| Indiana  | New England    | Pennsylvania  |
| Iowa     | (all 6 states) | Virginia      |
| Kentucky | New Jersey     | West Virginia |
| Maryland |                | Wisconsin     |



\* DOES NOT APPLY TO FALLOW LAND APPLICATIONS OR COUNTIES FOR WHICH AN APPROVED SLN (24C) IS ON FILE

**FOR SOIL INCORPORATED APPLICATIONS:** (Preplant Incorporated)  
See Geographic Restrictions above.

Except as restricted above (see Geographic Restrictions), Command may be used as a soil applied treatment without incorporation, however, additional precautions, beyond those required for a soil incorporated treatment, must be taken as follows:

Do not apply Command 4 EC within 1,500 feet of the following areas: Towns and Subdivisions, Commercial Fruit or Vegetable Production, Commercial Greenhouses or Nurseries.

Prior to application, adjacent properties must be checked, and spraying within 1,500 feet of desirable plants must be avoided.

\* except sweet corn

**Spray Drift Precautions:** Non-target spray drift of Command 4 EC herbicide should be avoided to prevent withering of desirable vegetation. Drift is influenced by many factors which include wind speed, spray pressure, particle size, nozzle type, and boom height.

- Do not apply when weather conditions favor drift if wind speeds exceed 10 miles per hour, a drift reducing additive must be used
- A minimum spray volume of 15 gallons per acre is recommended with appropriate nozzle types and sizes that produce coarser sprays
- The use of agriculturally approved drift reducing additives is recommended for application volumes of 15 to 40 gallons per acre when spraying in the proximity of desirable plants (see list above)
- The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre
- Use minimum nozzle pressure and boom height while maintaining uniform spray pattern.

• Do not exceed 40 psi spray pressure

Selection and proper use of spray equipment is critical in minimizing spray drift. The following table suggests pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types

## Suggested Nozzle Types, Minimum Size and Recommended Pressure Ranges for Minimizing Drift

| Nozzle               | Pressure Range (PSI) | Minimum Flow Rate Within Pressure Range (GPM) | Minimum Nozzle Size |
|----------------------|----------------------|---|---------------------|
| Flat-fan             | 15-30                | 0.3   | #4*                 |
| LP-Flat-fan          | 10-25                | 0.3   | #3                  |
| Even Flat-fan**      | 15-30                | 0.3   | #4                  |
| Flood                | 10-25                | 0.3   | #2.5*               |
| Whirl-chamber        | 5-20                 | 0.3   | #5                  |
| Randrop              | 15-40                | 0.15  | #2                  |
| Wide angle full cone | 15-40                | 0.3   | #5                  |

\* Refers to tip number such as 8004 or LF2.5

\*\* Recommended for banded applications

**ROTATIONAL CROPPING PRECAUTIONS.** Temporary whitening or yellowing of leaves may occur on approved rotational crops where undesirable soil residues of Command 4 EC herbicide exist. The following conditions may result in undesirable soil residues and greater potential for injury to rotational crops

- 1) Exceeding label recommended rates
- 2) Overapplication resulting through use of worn nozzles, excessive overlapping of spray swaths, failing to shut off spray booms when turning (end row areas), or slowing or stopping sprayer.

Refer to Rotational Guidelines and replanting instructions of specific crops for additional crop planting information

## SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

Do not apply Command 4 EC within 1,500 feet of the areas listed below

- Towns and Subdivisions
- Commercial Vegetable Production
- Commercial Fruit Production
- Commercial Nurseries
- Commercial Greenhouses

\*except sweet corn

Plants such as the following may show symptoms of foliar whitening or yellowing if contacted by Command 4 EC herbicide. It is recommended that, prior to application, adjacent properties be checked and that spraying within 100 feet of such plants be avoided

- Trees (Deciduous)**
- Apple (inc. fruit & ornamental types)
  - Ash (Green, White Mountain)
  - Basswood
  - Boxelder
  - Catalpa
  - Cherry (inc. fruit & ornamental types)
  - Cottonwood
  - Elm
  - Ginkgo
  - Hackberry
  - Mulberry
  - Peach
  - Pear (inc. fruit & ornamental types)
  - Pecan
  - Poplar
  - Russian olive
  - Tree-of-Heaven
  - Tulip tree
  - Walnut trees
  - Willow species

- Trees (Evergreen)**
- Fir species
  - Spruce species

- Shrubs & Vines**
- Azalea
  - Burningbush (Winged Euonymus)
  - Grape
  - Honeysuckle
  - Roses
  - Yew

- Agronomic Cereals**
- Ailanth
  - Oats

- Vegetable and Flower plants**

- Others**
- Ferns
  - Herbs
  - Strawberry
  - Raspberry
  - Blackberry

Apply Command 4 EC herbicide only to surfaces that will be soil incorporated. Do not apply Command 4 EC herbicide to non-field areas including fence rows, waterways, ditches, and roadsides

**FOR SOIL APPLIED APPLICATIONS:**  
See Geographic Restrictions above.

# GENERAL APPLICATION INSTRUCTIONS

This Product Must Be Soil Incorporated

Do not apply aerially or through irrigation equipment.

## Ground Applications

**Broadcast or Banded Applications:** Apply Command 4 EC alone or in tank mix combinations by ground equipment using a finished spray volume of 10 to 40 gallons of water per acre. NOTE: The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre. Use nozzles suitable for broadcast boom or banded application of herbicides. Coarse sprays are less likely to drift out of the target area than fine sprays. See "APPLICATION PRECAUTIONS" Section for specific recommendations to reduce spray drift. For Command 4 EC tank mixtures with wettable powder or dry flowable formulations, nozzle screens and strainers should be no finer than 50-mesh.

### Banded Applications—

Calculate the rates and volumes required by using the following formulas

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Rate Per Acre} = \text{Band rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Volume Per Acre} = \text{Band volume per acre}$$

Follow broadcast minimum gallonage requirements adjusted to band width. Do not exceed 40 psi spray pressure.

Refer to Spray Drift Precautions for suggested pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types. For banded applications an even flat-fan nozzle is recommended.

**Soil Incorporated Application:** Apply Command 4 EC to the soil surface and uniformly incorporate. Select equipment suited for proper incorporation.

In States where soil incorporation is required (see Geographical Restrictions), application and immediate incorporation to a maximum depth of 3 inches is required unless the soil is dry. On dry soils, incorporation to a maximum depth of 3 inches must be completed within 8 hours of Command 4 EC herbicide application.

Soil must be in good tilth to allow for thorough mixing of the soil. Application to wet soils will increase the potential for off-site movement of Command 4 EC vapors and may result in poor soil incorporation and unsatisfactory weed control.

## INCORPORATION EQUIPMENT

### ONE PASS INCORPORATION

#### Suggested Implements

For one pass incorporation use C or S shank field cultivators equipped with a leveling device such as a spike or coil line narrow or rolling basket attachment. Field cultivator must have 3 or 4 rows of sweeps spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Equipment design, including any drag attachments, must be adequate to avoid soil ridging which may result in streaked or reduced weed control. The implement should be operated at 5 to 8 mph and set to cut 3 to 4 inches deep to incorporate herbicide 1 to 2 inches.

Combination equipment (three or more tillage devices combined and used as a single tool) should be operated at a depth of 3 to 4 inches and a minimum speed of 6 mph.

PTO-drive seed bed preparation equipment should be set to cut 2 inches deep and operated at 4 mph or less.

### TWO PASS INCORPORATION

#### Suggested Implements

When using two pass incorporation, the second pass should be at an angle to the first pass.

Disc set to cut 3 to 4 inches deep at 4 to 6 mph. Blades should be less than 20 inches in diameter and spaced 9 inches or less apart.

Rolling cultivator set to cut 2 to 3 inches deep at 5 to 8 mph (use on coarse or medium textured soils only).

Field cultivator with 3 to 4 rows of sweeps spaced at 7 inch intervals or less and set to cut 3 inches deep at 5 mph or greater.

### Cultivation After Planting

Soil treated with Command 4 EC may be shallow-cultivated, rotary-hoed or hand-hoed without reducing the weed control activity of Command 4 EC. Do not cultivate deeper than the Command 4 EC treated layer of soil since this may bring untreated soil to the surface and poor weed control may result.

## COMMERCIAL IMPREGNATION AND APPLICATION OF COMMAND 4 EC ON DRY BULK FERTILIZERS

Command herbicide may be impregnated on dry bulk fertilizers. When applied as directed, Command/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of Command applied in water.

The Command fertilizer mixtures must be soil incorporated in all States listed under Geographic Restrictions of the Application Precautions Section of this label.

**Impregnation:** Apply using a minimum of 200 pounds of dry bulk fertilizer per acre and up to a maximum of 450 pounds per acre with the recommended amount of Command 4 EC herbicide per acre. Use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform, fine spray pattern over the tumbling fertilizer for thorough coverage. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Command provides a satisfactory, dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry, flowable mixture. Microcel E (Johns-Manville Products Corporation) is a recommended absorbent powder. Generally less than 2% by weight of Microcel E is required. DO NOT impregnate Command 4 EC onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with Command.

The amount of Command actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with Command 4 EC herbicide should be applied immediately, not stored. Care should be taken to ensure spreading procedures in the field do not place fertilizer treated with Command in areas which cannot be incorporated. If incorporation is required within the Geographic Restrictions of this label.

All state regulations, labeling, etc. of the mixtures are the responsibility of the seller.

For those rates not listed in the following table, calculate the amount of Command to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{2000 \text{ pounds dry fertilizer per acre}}{\text{PINTS of Command per acre (recommended rate for soil texture)}} = \frac{\text{PINTS of Command per ton of fertilizer}}$$

## RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZERS WITH COMMAND 4 EC HERBICIDE

PINTS OF COMMAND 4 EC PER TON OF FERTILIZER

| Fertilizer Rate Lbs. Acre | Command 4 EC Rate Per Acre |                |                        |
|---------------------------|----------------------------|----------------|------------------------|
|                           | 1 1/2 pts (75 lbs A1)      | 2 pt (1 lb A1) | 2 1/2 pts (125 lbs A1) |
| 200                       | 15                         | 20             | 25                     |
| 250                       | 12                         | 16             | 20                     |
| 300                       | 10                         | 13 1/2         | 16 2/3                 |
| 350                       | 8 1/2                      | 11 1/2         | 14                     |
| 400                       | 7 1/2                      | 10             | 12 1/2                 |
| 450                       | 7 1/2                      | 9              | 11                     |

If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

## SPECIAL PRECAUTIONS FOR DRY BULK FERTILIZER


### IMPREGNATION AND APPLICATION WITH COMMAND 4 EC

- All equipment used to apply Command should be thoroughly cleaned immediately following use to ensure no contamination results which could cause injury to non-labeled crops or desirable vegetation. Refer to "Sprayer Cleanup" section for additional details and disposal of rinsates.
- Equipment used to physically transport Command treated fertilizer, including boots, augers, conveyors, bins, etc. should be covered to prevent loss of fine particles and subjected to cleanup procedures previously described.

- Mixing of Command herbicide and dry fertilizer should be conducted in a blender which will ensure that airborne particles and potential for volatilization are avoided
- Bulk containers should be tightly covered while the product is being transported and applied to reduce chances of Command volatilization or product loss
- It is recommended that Dry Bulk Fertilizer impregnation operations using Command not be conducted within 1/4 mile of residential areas or areas where Command symptomology on desirable vegetation would not be tolerated

## COMMAND GEOGRAPHICAL AREAS

FOR RATES, WEED CONTROL, AND CROP ROTATION



- Southern area includes Missouri bootheel, which incorporates the following counties: Butler, Franklin, Mississippi, New Madrid, Pemiscot, Scott, and Stoddard.

## SOYBEANS

### GENERAL INFORMATION


Command 4 EC herbicide may be utilized as a preemergent soil applied treatment for the control of annual, grass and broadleaf weeds in soybeans.

Command 4 EC must be soil incorporated in the following States unless an approved(24 (C)) is on file for counties within that state (check with your local dealer): Delaware, Illinois, Indiana, Iowa, Kentucky, Maryland, Michigan, Minnesota, New England (all 4 states), New Jersey, New York, Ohio, Pennsylvania, Virginia, West Virginia, and Wisconsin.

Command 4 EC herbicide may be tank mixed with other herbicides to broaden the weed control spectrum compared to the products applied alone. Command 4 EC may be tank mixed with Canopy, Dual, Lasso, Lexion, Previc, Provl, Scepter, Sensor, Sonalan, Squadron, Trellan, or Tri-Scept herbicides. Observe all precautions, instructions, and rotational cropping guidelines of each product's label when tank mixing. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with the herbicides listed above, unless use directions specifically state otherwise.

### REPLANTING INSTRUCTIONS

If initial seeding of soybeans fails to produce a stand, soybeans may be replanted in fields treated with Command 4 EC alone or with recommended tank mixtures. Do not retreat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the soybean replant instructions for that product. Do not replant treated fields with any crop intervals which are inconsistent with the Rotational Guidelines on this label.



**COMMAND® 4 EC HERBICIDE  
APPLIED ALONE  
RATES AND WEEDS CONTROLLED**

| 1.0 PINT<br>(0.50 LB. A.I.)                            | 1.5 PINTS<br>(0.75 LB. A.I.)   | 2.0 PINTS<br>(1.0 LB. A.I.)  |
|--|--|--|
| <u>GRASSES:</u><br>None                                | <u>GRASSES:</u><br>Eaglegrass+<br>Cypergrass<br>- Large<br>- Smooth<br>Field Sandbur<br>Foxtail:<br>- Giant<br>- Green<br>- Foxtail<br>- Yellow<br>Goosegrass<br>Panicum:<br>- Common<br>- Fall<br>- Texas<br>Seedling Johnsongrass* | <u>GRASSES:</u><br>All grasses controlled<br>at 1.5 pints <u>plus</u> :<br><br>Barnyardgrass*<br>Broodleaf Signalgrass<br>Cypergrass<br>- Southwestern<br>- Yellow*<br>Seedling Johnsongrass<br>Shattercane*<br>Wild Proso Millet* |
| <u>BROADLEAVES:</u><br>Velvetleaf*<br>Spotted Amaranth | <u>BROADLEAVES:</u><br>Velvetleaf*<br>Spotted Amaranth<br><u>plus</u> :<br>Callisotea<br>Lanceleaflet<br>Florida Spider<br>Purslane<br>Venice Mallow<br>Common Ragweed*<br>Giantweed*<br>Pennsylvania Smartweed*                     | <u>BROADLEAVES:</u><br>All broadleaves controlled<br>at 1.5 pints <u>plus</u> :<br><br>Common Ragweed<br>Giantweed<br>Purslane<br>Pennsylvania Smartweed<br>Venice Mallow<br>Giantweed*  |


\* Partially controlled  
 + Higher rates are recommended for fields that have previously been in  
 crop rotation programs. In these heavy weed pressure is anticipated.  
 Control is based on higher rates based on lighter to heavier soil types.

NOTE: When using Command 4 EC on broadleaves incorporated more than 30 days prior  
 to planting, the highest label rate should be used.

**ROTATIONAL CROP GUIDELINES**

Rotate to crops as listed below, otherwise crop injury may occur

| 1.0 PINT<br>(0.50 LB. A.I.)  | 1.5 PINTS<br>(0.75 LB. A.I.)                                    | 2.0 PINTS<br>(1.0 LB. A.I.)   |
|--|---|---|
| <u>ANYTIME</u><br>Soybeans<br>Peanuts<br>Peanut<br>Peanut  | <u>ANYTIME</u><br>Soybeans<br>Peanuts                           | <u>ANYTIME</u><br>Soybeans<br>Peanuts   |
| <u>9 MONTHS</u><br>Corn<br>- Field<br>- Pop<br>- Seed<br>- Sweet<br>Cotton<br>Cucurbits<br>Dry Beans<br>Peanuts<br>Peppers<br>Potatoes<br>Rice<br>Snap Beans<br>Sorghum<br>Sweet Beans<br>Sweet Potatoes<br>Tobacco<br>Tomatoes (Transplanted) | <u>9 MONTHS</u><br>Same as 1.0<br>pint. <u>plus</u> :<br>Peanut | <u>9 MONTHS</u><br>Same as 1.5<br>pints. <u>except</u><br>Seed Corn and<br>Sweet Corn require<br>a 12 month rotation. |
| <u>12 MONTHS</u><br>Cabbage<br>Cabbage (Fall)<br>Wheat   | <u>12 MONTHS</u><br>Same as 1.0<br>pint.                        | <u>12 MONTHS</u><br>Same as 1.5<br>pints. <u>plus</u> :<br>Seed Corn<br>Sweet Corn                                    |
| <u>15 MONTHS</u><br>All crops  | <u>15 MONTHS</u><br>All crops                                   | <u>15 MONTHS</u><br>All crops   |



Cover crops, however, may be planted anytime but stand reductions  
 may occur in some areas. Do not graze or harvest for food or feed cover  
 crops planted less than 9 months after Command 4 EC treatment. Do  
 not allow livestock to graze on treated soybean vines or feed treated  
 vines or vine trash to livestock.

# Southern Area—See Map

## COMMAND 4 EC HERBICIDE APPLIED ALONE RATES AND WEEDS CONTROLLED

| 1.0 PINT<br>(0.50 LB. A.I.)  | 1.5 PINTS<br>(0.75 LB. A.I.)  | 2.0 PINTS<br>(1.0 LB. A.I.)   | 2.5 PINTS<br>(1.25 LB. A.I.)   |
|--|---|---|--|
| <b>GRASSES:</b><br>None  | <b>GRASSES:</b><br>Barnyardgrass+<br>Broadleaf Signalgrass+<br>Crabgrass+<br>- Lacta<br>- Smooth<br>Foxtail+<br>- Giant<br>- Green<br>Clovegrass+<br>Panicum+<br>- Cotton<br>- Fall<br>- Texas<br>Seedling Johnsongrass+<br>Field Bindweed* | <b>GRASSES:</b><br>All grasses<br>controlled at<br>1.5 pints <u>plus</u> :<br>Sugarcass<br>- Southwestern<br>- Hoop<br>Field Bindweed<br>Barnyardgrass*<br>Red Rice*<br>Johnsongrass*                                       | <b>GRASSES:</b><br>All grasses<br>controlled at<br>2.0 pints <u>plus</u> :<br>Foxtail<br>- Purple<br>- Pectus<br>- Yellow<br>Crabgrass+<br>Wild Penna Millers*   |
| <b>BROADLEAVES:</b><br>All broadleaf<br>broadleaves+<br>Sourwood+* | <b>BROADLEAVES:</b><br>Field Bindweed<br>Spurred Amaranth<br>plus:<br>Lambquarters+<br>Prickly Soda+<br>Purslane+<br>Redvein+<br>Venice Malabar<br>Common Ragweed*<br>Jimsonweed*<br>Pennsylvania<br>Smartweed*                             | <b>BROADLEAVES:</b><br>All broadleaves<br>controlled at<br>1.5 pints <u>plus</u> :<br>Cocklebur+<br>Common Ragweed+<br>Dayflower<br>Florida Pusley<br>Jimsonweed<br>Tropic Cotton<br>Wild Poinsettias<br>Yucca*<br>Redvine* | <b>BROADLEAVES:</b><br>All broadleaves<br>controlled at<br>2.0 pints <u>plus</u> :<br>Yucca<br>Morningglory<br>- Pitted<br>Pennsylvania<br>Smartweed<br>Prostrate Scurf<br>Black Nightshade*<br>Bigweed<br>- Redroot*<br>- Smooth* |

- \* Partially controlled.
- + Higher rates are recommended for fields that have previously been in no-till or strip-till programs, or where heavy weed pressure is anticipated.
- + Rates listed to higher rates based on lighter to heavier soil types.
- + Use in April to June only. For high pressure and heavy soils use 2.5 pints.
- + Use only where applications will not require soil incorporation (See Geographic Restrictions). For high weed pressure and heavy soils use 2.5 pints.

NOTE: When using Command 4 EC preplant or preemergence treatments, mark with 10 days before to 14 days after the broadcast application of the herbicide.

### ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below. Otherwise crop injury may occur.

| 1.0 PINT<br>(0.50 LB. A.I.)   | 1.5 TO 2.0 PINTS<br>(0.75 TO 1.0 LB. A.I.)               | 2.5 PINTS<br>(1.25 LB. A.I.)  |
|---|--|---|
| <b>ANYTIME</b><br>Soybeans<br>Peanuts<br>Rice   | <b>ANYTIME</b><br>Soybeans<br>Peanuts                    | <b>ANYTIME</b><br>Soybeans  |
| <b>9 MONTHS</b><br>Corn<br>- Field<br>- Pop<br>- Seed<br>- Sweet<br>Cotton<br>Cucumbers<br>Dry Beans<br>Peanuts<br>Pumpkins<br>Rice<br>Shed Beans<br>Sorghum<br>Sweet Beans<br>Sweet Potatoes<br>Tomato<br>TOMATOES - Transplanted. | <b>9 MONTHS</b><br>Same as 1.0 pint, <u>plus</u><br>Rice | <b>9 MONTHS</b><br>Same as 1.5 to 2.0 pints, <u>plus</u><br>Peanuts |
| <b>12 MONTHS</b><br>All crops   | <b>12 MONTHS</b><br>All crops                            | <b>12 MONTHS</b><br>All crops                                       |

**BEST AVAILABLE COPY**

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

# COMMAND 4 EC HERBICIDE APPLIED IN COMBINATIONS FOR SOYBEANS

Command 4 EC herbicide may be tank mixed with other soybean herbicides to broaden the weed control spectrum compared to the products applied alone. In the sections that follow, FMC recommends a combination of products and application rates to broaden weed control spectrum. FMC makes no claims as to weed control for any combination of rates or products other than as recommended in this label. Additional recommendations for rate combinations with other herbicides not included in the following sections may become available subsequent to the date of publication for this label. Please check with your dealer for more recent FMC technical bulletins and label supplements. When apply mixtures of Command 4 EC with other herbicides, observe all precautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixes.

## COMMAND 4 EC Herbicide plus Canopy<sup>®</sup> Herbicide— Northern & Southern Areas

Tank mix combination for broadened weed control spectrum

| Soil Texture  | Broadcast Rates Per Acre* |                  |           |
|---|---------------------------|------------------|-----------|
|   | Command                   | Canopy           |           |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 1 1/2 to 2 pints          | 2 to 2 1/2 pints | 4 to 6 oz |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 1 1/2 to 2 pints          | 2 to 2 1/2 pints | 5 to 8 oz |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 1 1/2 to 2 pints          | 2 to 2 1/2 pints | 6 to 8 oz |

\* Select lower to higher rates within the range to correspond to the coarser to finer soil types within the textural group.

\*\* Rate within the range should correspond to anticipated weed pressure.

\*\*\* Do not apply to soils less than 1% Organic Matter.

### WEEDS CONTROLLED

#### Grass Weeds

All species listed under Command Alone Weeds Controlled.

#### Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:

Cocklebur<sup>†</sup>  
Pigweed (Redroot, Smooth)  
Hemp sesbania  
Morningglory\*\* (annual ivy-leaf, entire-leaf, pitted, tall)  
Sicklepod\*\*  
Copperleaf hophornbeam\*\*

<sup>†</sup> Large seeded weeds germinating deep in the soil, such as morningglory, sicklepod and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a postemergence herbicide for season-long control.

\*\* Partially Controlled.

When applying mixtures of Command 4 EC plus Canopy, observe all cautions and limitations appearing on the labels of each product. Special attention must be paid to sections regarding planting depth, prostrate pesticide interaction, previous pesticides, and sensitive soybean varieties on the Canopy label. Observe Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixtures.

## COMMAND 4 EC herbicide plus Dual<sup>®</sup> 8 E Herbicide— Northern & Southern Areas

Tank-mix combination for broadened weed control spectrum

| Soil Texture  | Broadcast Rates Per Acre |                  |                  |
|---|--------------------------|------------------|------------------|
|   | Command                  | Dual 8 E         |                  |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 1 1/2 to 2 pints |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 2 to 2 1/2 pints |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 2 to 2 1/2 pints |

### WEEDS CONTROLLED

#### Grass Weeds

All species listed under Command Alone Weeds Controlled plus the following:  
Red Rice<sup>†</sup>  
Witchgrass  
Nutsedge

#### Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:  
Pigweed  
Carpetweed  
Black Nightshade

\* Use higher rates in areas with heavy pressure.

When applying mixtures of Command 4 EC plus Dual, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixtures.

## COMMAND 4 EC Herbicide plus Lasso<sup>®</sup> or Lasso<sup>®</sup> Micro-Tech—Northern & Southern Areas

A tank-mix combination of Command 4 EC herbicide and Lasso or Lasso Micro-Tech herbicide will broaden the weed control spectrum compared to either herbicide used alone.

| Soil Texture  | Broadcast Rates Per Acre |                  |                   |
|---|--------------------------|------------------|-------------------|
|   | Command                  | Lasso            | Lasso Micro-Tech  |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 2 to 2 1/2 quarts |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 2 1/2 to 3 quarts |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 2 1/2 to 3 quarts |

\* NOTE: Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., sand, loamy sand, sandy loam) within a textural group.

### WEEDS CONTROLLED

#### Grass Weeds

All species listed under Command Alone Weeds Controlled.

#### Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:  
Carpetweed

#### Pigweed

When applying mixtures of Command 4 EC herbicide plus Lasso or Lasso Micro-Tech, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixtures.



**COMMAND 4 EC herbicide plus Metribuzin (Sencor\* or Lexone\*)—Northern Area Only**

A tank mix combination of Command 4 EC herbicide and metribuzin will broaden the weed control spectrum compared to either herbicide used alone.

| Soil Texture**   | Broadcast Rates Per Acre* |                         |
|--|---------------------------|-------------------------|
|  | Command                   | Sencor DF*<br>Lexone DF |
| Coarse (light) Soils: (loamy sand**, sandy loam)                   | 1½ to 2 pints             | ¼ to ½ lb               |
| Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam) | 1½ to 2 pints             | ¼ to ½ lb               |
| Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay) | 1½ to 2 pints             | ¼ to ½ lb               |

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

\*\* For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.

\*\*\* Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

**WEEDS CONTROLLED:**

Grass Weeds

All species listed under Command Alone Weeds Controlled,

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled, plus the following:  
Carpetweed  
Pigweed (Redroot, Smooth)  
Wild mustard

\* Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixtures.

**COMMAND 4 EC herbicide plus Metribuzin (Sencor\* or Lexone\*)—Southern Area Only**

A tank mix combination of Command 4 EC herbicide and metribuzin will broaden the weed control spectrum compared to either herbicide used alone.

| Soil Texture**   | Broadcast Rates Per Acre* |                         |
|--|---------------------------|-------------------------|
|  | Command                   | Sencor DF*<br>Lexone DF |
| Coarse (light) Soils: (loamy sand**, sandy loam)                   | 2 to 2½ pints             | ¼ to ½ lb               |
| Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam) | 2 to 2½ pints             | ¼ to ½ lb               |
| Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay) | 2 to 2½ pints             | ¼ to ½ lb               |

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

\*\* For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.

\*\*\* Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur. Use 4½ to 7 pound metribuzin on silty clay and heavy clay soils of the Mississippi Delta.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled.

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled, plus the following:  
Carpetweed  
Hemp sesbania (coffee weed)  
Pigweed (Redroot, Smooth)  
Wild mustard  
Sicklepod\*\*

\* Reduced Competition

\*\* Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Preview\*—Northern Area Only**

A tank-mix combination of Command 4 EC herbicide and Preview herbicide will broaden the weed control spectrum compared to either herbicide used alone.

| Soil Texture   | Broadcast Rates Per Acre |           |
|--|--------------------------|-----------|
|  | Command                  | Preview*  |
| Coarse (light) Soils: (loamy sand, sandy loam)                     | 1½ to 2 pints            | 5 to 6 oz |
| Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam) | 1½ to 2 pints            | 5 to 7 oz |
| Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay) | 1½ to 2 pints            | 6 to 8 oz |

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

Do not use Command 4 EC in combination with Preview on soils with less than 0.5% organic matter, or on soils with pH greater than 6.8 or on soils having a calcareous surface area as Preview injury to rotational crops may occur.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled.

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:  
Carpetweed  
Spurge spotted  
Copperleaf, hophornbeam  
Mustards  
Pigweed  
Cocklebur\*  
Sunflower\*  
Burdock  
Giant Ragweed\*  
Morningglory\* (Annual, Tall, leafy, Pitted, Entireleaf)

\* NOTE: Control of morningglory, sunflower, cocklebur and giant ragweed may be erratic ranging from poor to excellent depending upon soil temperature, time of weed seed germination, depth of weed seed in the soil, and the amount and timing of soil moisture. Control may be improved with timely tillage.

When applying mixtures of Command 4 EC plus Preview, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Prowl<sup>®</sup> C Herbicide—  
Northern & Southern Areas**

Tank-mix combination for broadened weed control spectrum

Broadcast Rates Per Acre

| Soil Texture  | Command       |               | Prowl 4 EC*   |
|---|---------------|---------------|---------------|
|   | North         | South         |               |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 1½ to 2 pints | 2 to 2½ pints | 1 to 1½ pints |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 1½ to 2 pints | 2 to 2½ pints | 2 pints       |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 1½ to 2 pints | 2 to 2½ pints | 2 to 2½ pints |

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled.

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:  
Pigweed (Redroot, Smooth)  
Carpenterweed

When applying mixtures of Command 4 EC plus Prowl, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Scepter<sup>®</sup> Herbicide—  
Northern Area Only**

See Scepter label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

Broadcast Rates Per Acre

| Soil Texture  | Command       |             | Scepter* |
|---|---------------|-------------|----------|
|   | Command       | Scepter*    |          |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 1½ to 2 pints | ½ to ¾ pint |          |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 1½ to 2 pints | ½ to ¾ pint |          |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 1½ to 2 pints | ½ to ¾ pint |          |

\* Do not exceed 1/3 pint in Northern Area (as defined on the Scepter label). Use higher rate of Scepter in areas with heavy weed pressure and anticipated insecticide.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled.

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:  
Cocklebur  
Pigweed spp.\*  
Giant Ragweed\*

\* Partially controlled.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than two inches, or make additional Command 4 EC or Command Scepter tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

**DO NOT GRAZE OR FEED TREATED FORAGE, HAY OR STRAW TO LIVESTOCK**

When applying mixtures of Command 4 EC plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Scepter Herbicide—  
Southern Area Only**

See Scepter label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

Broadcast Rates Per Acre

| Soil Texture  | Command       |             | Scepter* |
|---|---------------|-------------|----------|
|   | Command       | Scepter*    |          |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 2 to 2½ pints | ½ to ¾ pint |          |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 2 to 2½ pints | ½ to ¾ pint |          |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 2 to 2½ pints | ½ to ¾ pint |          |

\* Use higher rates when heavier weed pressure is anticipated.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled.

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:

Cocklebur  
Pigweed spp.\*  
Wild poinsettia

\* Partially controlled.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than two inches, or make additional Command 4 EC or Command-Scepter tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

**DO NOT GRAZE OR FEED TREATED FORAGE, HAY OR STRAW TO LIVESTOCK**

When applying mixtures of Command 4 EC plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Sonalan<sup>®</sup> EC Herbicide—  
Northern Area Only**

Tank-mix combination for broadened weed control spectrum

Broadcast Rates Per Acre\*

| Soil Texture  | Command       |                | Sonalan EC |
|---|---------------|----------------|------------|
|   | Command       | Sonalan EC     |            |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 1½ to 2 pints | 1¼ to 2 pints  |            |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 1½ to 2 pints | 1½ to 2½ pints |            |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 1½ to 2 pints | 2¼ to 3 pints  |            |

\* Select lower to higher rates within the range to correspond to the coarser to finer soil types within the textural group. Rates within the range should correspond to anticipated weed pressure.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled.

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:

Carpenterweed\*  
Pigweed (Redroot, Smooth)

When applying mixtures of Command 4 EC plus Sonalan, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Squadron**  
Northern Area Only

See *Squadron* label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Squadron will broaden the weed control spectrum of either product used alone

| Soil Texture   | Broadcast Rates Per Acre |                      |
|--|--------------------------|----------------------|
|  | Command                  | Squadron*            |
| Coarse (light) Soils: (loamy sand, sandy loam)                     | 1 1/2 to 2 pints         | 1 1/2 to 2 1/4 pints |
| Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam) | 1 1/2 to 2 pints         | 1 1/2 to 2 1/4 pints |
| Fine (heavy) Soils: (silt clay, clay loam, silty clay loam, clay)  | 1 1/2 to 2 pints         | 1 1/2 to 2 1/4 pints |

\* Do not exceed 1 1/2 pints in Northern Area (as defined on the Squadron label). Use higher rate of Squadron in areas of heavy cocklebur infestation.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following

Cocklebur  
Pigweed spp

Giant Ragweed\*

\* Partially controlled

When applying mixtures of Command 4 EC plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Squadron --**  
Southern Area Only

See *Squadron* label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Squadron will broaden the weed control spectrum of either product used alone

| Soil Texture  | Broadcast Rates Per Acre |                  |
|---|--------------------------|------------------|
|   | Command                  | Squadron*        |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 2 to 2 1/2 pints         | 2 1/4 to 3 pints |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 2 to 2 1/2 pints         | 2 1/4 to 3 pints |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 2 to 2 1/2 pints         | 2 1/4 to 3 pints |

\* Use higher rate of Squadron in areas of heavy pigweed and cocklebur infestation.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following

Cocklebur  
Pigweed spp  
Wild poinsettia

When applying mixtures of Command 4 EC plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Trellan EC (Trifluralin)**  
Northern & Southern Areas

A tank mix combination of Command 4 EC herbicide and Trellan EC herbicide will broaden the weed control spectrum compared to either herbicide used alone

| Soil Texture   | Broadcast Rates Per Acre |                  |                  |
|--|--------------------------|------------------|------------------|
|  | Command North            | Command South    | Trellan EC*      |
| Coarse (light) Soils: (loamy sand, sandy loam)                     | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 1 pint           |
| Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam) | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 1 to 1 1/2 pints |
| Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay) | 1 1/2 to 2 pints         | 2 to 2 1/2 pints | 1 to 1 1/2 pints |

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following

Carpetweed  
Chickweed  
Field Bindweed  
Pigweed (Redroot, Smooth)

When applying mixtures of Command 4 EC plus Trellan, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Tri-Scept Herbicide**  
Northern Area Only

See *Tri-Scept* label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Tri-Scept applied soil incorporated will broaden the weed control spectrum of either product used alone

| Soil Texture  | Broadcast Rates Per Acre |                      |
|---|--------------------------|----------------------|
|   | Command                  | Tri-Scept            |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 1 1/2 to 2 pints         | 1 1/8 to 1 1/4 pints |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 1 1/2 to 2 pints         | 1 1/8 to 1 1/4 pints |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 1 1/2 to 2 pints         | 1 1/8 to 1 1/4 pints |

\* Do not exceed 1 1/8 pints in Northern Area (as defined on the Tri-Scept label). Use higher rates of Tri-Scept in areas of heavy cocklebur infestation.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following

Cocklebur  
Pigweed spp

Giant Ragweed\*

\* Partially controlled

When applying mixtures of Command 4 EC plus Tri-Scept observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC Herbicide plus Tri-Sect Herbicide—  
Southern Area Only**

See Tri-Sect label for appropriate geographic restrictions

A tank mix combination of Command 4 EC and Tri-Sect applied soil incorporated will broaden the weed control spectrum of either product used alone

| Soil Texture  | Broadcast Rates Per Acre |                |
|---|--------------------------|----------------|
|   | Command                  | Tri-Sect       |
| Coarse (light) Soils (loamy sand, sandy loam)                     | 2 to 2½ pints            | 1½ to 2½ pints |
| Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam) | 2 to 2½ pints            | 1½ to 2½ pints |
| Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay) | 2 to 2½ pints            | 1½ to 2½ pints |

\* Use higher rate of Tri-Sect in areas of heavy pigweed and cocklebur infestation

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Alone Weeds Controlled

Broadleaf Weeds

All species listed under Command Alone Weeds Controlled plus the following:  
Cocklebur  
Pigweed spp.  
Wild poinsettia

When applying mixtures of Command 4 EC plus Tri-Sect, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting instructions of each product's label when using tank mixtures

**COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY PREEMERGENCE TREATMENTS**

Preemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of additional weed species listed on the preemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information that appear on the product labels. DO NOT apply compounds containing metribuzin after a soil incorporated Command 4 EC plus Lexone® or Sencor® tank mix treatment, unless soil applications are used where the pounds of active ingredient do not exceed the total allowed

DO NOT apply Canopy, Preview, Pursuit Plus, Scepter, Squadron, or Tri-Sect herbicides after Command 4 EC plus any of those products in a tank mix treatment, as these other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

**COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY POSTEMERGENCE TREATMENTS**

Postemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of emerged weeds as listed on the postemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information appearing on the product labels.

DO NOT apply Classic, Pursuit, or Scepter as a postemergence treatment after Command 4 EC plus Canopy, Preview, Pursuit Plus, Scepter, Squadron, or Tri-Sect in a tank mix treatment, unless the other product labels specifically recommend such a treatment, as these other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

**COMMAND 4 EC HERBICIDE FOLLOWED BY ACIFLUORFEN (BLAZER® 2L OR TACKLE®) HERBICIDE**

Following an application of Command 4 EC or tank mixture with Command 4 EC, a postemergence application of acifluorfen will control the following emerged broadleaf weeds

Broadleaf Weeds

- Cocklebur
- Hemp sesbania
- Pigweed (redroot)
- Black nightshade
- Wild mustard
- Purple Moonflower
- Pitted Morning glory

Apply acifluorfen at a rate of 1 pint per acre with 25% crop oil concentrate to actively growing weeds at no more than the 4 leaf growth stage (do not count cotyledonary leaves but only the fully developed true leaves). This timing generally correlates to soybean growth stages of the first to third trifoliate leaves

**NOTE:** Application of Command 4 EC herbicide generally retards the development rate of weeds which may extend the period where acifluorfen can be effectively applied

Read and follow all precautions, restrictions, and warnings on all product labels

**COMMAND 4 EC Herbicide followed by OPTION Herbicide**  
Following an application of Command 4 EC or tank mixture with Command 4 EC, a postemergence application of Option herbicide will control the following emerged annual and perennial grasses at the stages of growth that follow:

| Annual Grasses    | Growth Stage | Application Rates |           | Crop Oil Concentrate<br>1 qt./79/gal.<br>or 1 pt./49/gal. |
|-------------------|--------------|-------------------|-----------|---|
|                   |              | Fls./A            | Fl. Or./A |   |
| Volunteer Corn    | 5-10"        | 1 1/2             | 1 1/2     | DO NOT ADD OIL  |
| Wild Proso Millet | 5-10"        | 1 1/2             | 1 1/2     |   |
| Shattercane       | 5-10"        | 1 1/2             | 1 1/2     |   |
| Johnsongrass      | 1-5"         | 1 1/2             | 1 1/2     |   |
| Jungle Rice       | "            | "                 | "         |   |
| Sprangletop       | "            | "                 | "         |   |
| Wild Oats         | "            | "                 | "         |   |
| Wires on muleth   | "            | "                 | "         |   |
| Yellow foxtail    | "            | "                 | "         |   |
| Perennial Grasses | "            | "                 | "         |   |
| Wild Oats         | 10-10"       | 1 1/2             | 1 1/2     | DO NOT ADD OIL  |

\* Use crop oil concentrate for annual grasses that have grown beyond recommended stages of growth  
† When controlling mixed populations of grassy weeds, use the higher rate

Read and follow all precautionary statements and restrictions, directions for use, and all other information appearing on the Option label

**COMMAND 4 EC HERBICIDE FOLLOWED BY SCEPTER® HERBICIDE**

Following an application of Command 4 EC or tank mixtures with Command 4 EC, a postemergence application of Scepter herbicide will control the following emerged broadleaf weeds

Broadleaf Weeds

- Cocklebur, common
- Pigweed
- Wild poinsettia

Apply Scepter after crop emergence but before weeds exceed a height of 12 inches (6 inches for wild poinsettia). Apply at a broadcast rate of 1/3 pints per acre. Apply when weeds are actively growing. DO NOT apply Scepter postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes. The total amount of Scepter should not exceed 1/3 pints per acre per season in northern areas or 2/3 pint per acre per season in southern areas as defined on the Scepter label. Squadron and Tri-Scept both include the same active ingredient as Scepter.

For postemergence applications, the addition of a nonionic surfactant or crop oil concentrate is required. The nonionic surfactant should contain at least 80% active ingredient and should be applied at a rate of 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate stated on the COC label.

Read and follow all precautionary statements and restrictions, directions for use and all other information appearing on the Scepter label

**PUMPKINS**

**GENERAL INFORMATION**

Command 4 EC herbicide may be utilized as a preemergent soil applied treatment for the control of annual grass and broadleaf weeds in pumpkins.

Command 4 EC must be soil incorporated in the following states unless an approved (24C) is on file for counties within that state (check with your local dealer): Delaware, Illinois, Indiana, Iowa, Kentucky, Maryland, Michigan, Minnesota, New England (all 6 states), New Jersey, New York, Ohio, Pennsylvania, West Virginia, and Wisconsin.

Command 4 EC herbicide may be tank mixed with other pumpkin herbicides to broaden the weed control spectrum compared to the products applied alone. When applying mixtures of Command 4 EC herbicide plus other pumpkin herbicides, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixtures. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with the herbicides listed above unless use directions specifically state otherwise.

**SPECIAL PRECAUTION**

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause leaf whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

## REPLANTING INSTRUCTIONS

If initial seeding of pumpkins fails to produce a stand, pumpkins may be replanted in fields treated with Command 4 EC alone. Do not retreat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the pumpkin replant instructions for that product. Do not replant treated fields with any crop

at intervals which are inconsistent with the Rotational Guidelines on this label.

### COMMAND 4 EC APPLIED ALONE RATES AND WEEDS CONTROLLED

Do not exceed a rate of 2 pints per acre. Weed controlled (by rate) are as listed under Soybean for Northern and Southern areas.

### ROTATIONAL CROP GUIDELINES

Rotate to crops as listed under Soybean Rotational Crop Guidelines, otherwise crop injury may occur. Do not allow livestock to graze on treated pumpkin vines or feed treated vines or vine trash to livestock.

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated pumpkin vines or feed treated vines or vine trash to livestock.

## SUCCULENT PEAS

### GENERAL INFORMATION

Command 4 EC herbicide may be utilized as a preemergent soil applied treatment for the control of annual grass and broadleaf weeds in peas.

Command 4 EC must be soil incorporated in the following states unless an approved (24C) is on file for counties within that state (check with your local dealer): Delaware, Illinois, Indiana, Iowa, Kentucky, Maryland, Michigan, Minnesota, New England (all 6 states), New Jersey, New York, Ohio, Pennsylvania, West Virginia, and Wisconsin.

Command 4 EC herbicide may be tank mixed with other herbicides registered for use in peas to broaden the weed control spectrum compared to the products applied alone. When applying mixtures of Command 4 EC plus other succulent pea herbicides, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Guidelines and Replanting Instructions of each product's label when using tank mixes. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with other succulent pea herbicides unless use directions specifically state otherwise.

### SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

## REPLANTING INSTRUCTIONS

If initial seeding of peas fails to produce a stand, peas may be replanted in fields treated with Command 4 EC alone (or with recommended tank mixtures). Do not retreat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the replant instructions for that product. Do not replant treated fields with any crop

at intervals which are inconsistent with the Rotational Guidelines on the product booklet.

### COMMAND 4 EC APPLIED ALONE RATES AND WEEDS CONTROLLED

FOR PEAS ONLY. Apply only at 1 pint (0.5 LB. A.I.) per acre.

RATE AND WEEDS CONTROLLED (All Soil Types)  
1 PT. (0.5 LB. A.I.)

#### GRASSES

Crabgrass (Large Smooth)  
Panicum (Fall Common)  
Foxtail (Green Giant Robust)  
Goosegrass  
Field Sandspur  
Barnyardgrass  
Seedling Johnsongrass\*

#### BROADLEAVES

Velvetleaf  
Spurred Anoda  
Verice-Mallow  
Lambsquarters  
Prickly Sida  
Galinsoga spp.

\* Partially Controlled

## ROTATIONAL CROP GUIDELINES

Rotate to crops as listed under Soybean Rotational Crop Guidelines, otherwise crop injury may occur. Do not allow livestock to graze on treated pea vines or feed treated vines or vine trash to livestock.

Cover crops may be planted at anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated pea vines or feed treated vines or vine trash to livestock.

## FALLOW LAND

### GENERAL INFORMATION

Command 4 EC is a herbicide which may be utilized as a preemergence surface applied treatment to control many annual grass and broadleaf weeds on fallow land. Command 4 EC may be tank mixed with other herbicides registered for fallow cropland use (e.g., atrazine, chlor-sulfuron, etc.) to broaden the weed control spectrum compared to the products applied alone.

### SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants prior to making applications. Read and strictly follow all precautions and application instructions on this label.

Additional Precaution for Fallow Land Applications only: Do not apply within 500 feet of Emerged Winter Wheat.

ROTATIONAL CROP/GRAZING AND FEEDING RESTRICTIONS. Do not plant wheat sooner than nine (9) months after a late summer or fall application. Do not allow livestock to graze on treated fields.

### APPLICATION RATES:

Use Command 4 EC herbicide alone at a rate of 1 1/2 to 2 pints (24 to 32 ounces) per acre in a surface applied broadcast application for pre-emergent control of weeds. Use the higher rate when heavier weed infestations are anticipated or for longer residual control.

In areas where the winter wheat-fallow-winter wheat cropping system is practiced, make application after wheat harvest. For optimum control, make applications after August 15 but before October 31. Command 4 EC can also be used in areas where spring wheat is planted following an 18 month fallow period.

Treatments should be made only to fields where good weed control has been maintained in the wheat crop through winter harvest. Treatment recommendations are only designed to control weeds that have recently emerged and those that germinate throughout the fall, winter and spring period. Do not apply in situations where dense weed growth exists.

### WEEDS CONTROLLED

#### Grass Weeds

Downy bromegrass  
Jointed goatgrass  
Volunteer rye  
Volunteer wheat

#### Broadleaf Weeds

Common lambsquarters  
Common purslane  
Kochia  
Prickly lettuce  
Tranly mustard  
Tumble mustard  
Wild buckwheat

## COMMAND 4 EC APPLIED IN COMBINATIONS

Tank mix combinations of Command 4 EC herbicide with atrazine will broaden the weed control spectrum and length of control compared to the herbicides used alone.

### COMMAND plus ATRAZINE

Command 4 EC at 1 to 1 1/2 pints (16 to 24 ounces) per acre plus 1 to 2 pints of atrazine 4L or equivalent rates of atrazine 90 DF and 80 W. Weeds controlled in addition to those listed under Command 4 EC alone will depend on atrazine rate used (see atrazine label). Use the higher rate of Command 4 EC when heavier weed infestations are anticipated or for longer residual control.

There is significant risk of atrazine carryover if not used according to local Experiment Station and label recommendations. Consult your local Experiment Station for recommended atrazine rates for use in fallow in your area.

Refer to the atrazine label for planting, cropping and other restrictions. Follow all precautions, statements, and restrictions on this and the atrazine label.

### Mixing Instructions:

Fill spray tank 1/2 to 3/4 full with water and add recommended amount of atrazine with agitator operating. Mix thoroughly, and fill tank 3/4 full continuing agitation. Add Command 4 EC herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Maintain agitation during mixing and application.

### Application Instructions:

Follow the General Application Instructions as for Command 4 EC alone.

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions.

**Warranty:** FMC makes no warranty expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

COMMAND<sup>®</sup>, and FMC<sup>®</sup>—Trademarks of FMC Corporation  
CANOPY<sup>®</sup>, LEXONE<sup>®</sup> and PREVIEW<sup>®</sup>—Trademarks of E. I. duPont de Nemours and Co., Inc.  
OPTION<sup>®</sup>—Trademark of Hoechst-Roussel Agri-Vet Company  
SONALAN<sup>®</sup>, TREFLAN<sup>®</sup>—Trademarks of Elianco Products Company  
LASSO<sup>®</sup>—Trademark of Monsanto Company  
SENCOR<sup>®</sup>—Trademark of Parent Company of Farbentabriken Bayer GmbH, Leverkusen  
PROWL<sup>®</sup>, SCOPER<sup>®</sup>, SQUADRON<sup>®</sup>, TRISCEPT<sup>®</sup>—Trademarks of American Cyanamid Company  
DUAL<sup>®</sup>—Trademark of Ciba-Geigy Corporation  
BLAZER<sup>®</sup>—Trademark of BASF AG  
TACKLE<sup>®</sup>—Trademark of Rhone-Poulenc, Inc.  
© 1987 FMC Corporation

All rights reserved  
618-1188 A1