

Code 618

Net Contents

Command 4 EC

Herbicide

For Agricultural or Commercial Use Only
NOT FOR SALE OR USE IN CALIFORNIA

EPA Reg. No. 279-3053

EPA Est. 279-

Active Ingredient:	By Wt.
2-(2-Chlorophenyl)methyl-4, 4-dimethyl-3-isoxazolidinone	47.1%
Inert Ingredients:	52.9%
	100 0%

Contains xylene range aromatic solvents
Contains 4 pounds of active ingredient per gallon
U.S. Patent No. 4,405,357

KEEP OUT OF REACH OF CHILDREN WARNING

FIRST AID

If in eyes: Flush immediately with plenty of water. Get medical attention as soon as possible.

If swallowed: Drink promptly large quantities of milk or water. Avoid alcohol. Do not induce vomiting. Call a physician.

If inhaled: Remove to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention as soon as possible.

If on skin: Remove contaminated clothing. Wash skin with plenty of soap and water.

For Emergency Assistance Call Collect: (716) 735-3765

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

Causes eye injury (corneal opacity) that is temporary. Do not get in eyes or on clothing. Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing vapor or spray mist. Wear goggles or face shield when measuring, mixing or loading. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.


BEST AVAILABLE COPY

FMC

FMC Corporation
Agricultural Chemical Group
Philadelphia PA 19103

Dated on
1/4/84 - SAL

Page 1 of 13



Environmental Hazards

Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

See other panels for additional precautionary information.

TABLE OF CONTENTS

Read all instructions on pages 1-4 before referring to specific crop use.

	Page
Ingredients	1
First Aid	1
Precautionary Statements	1
Directions for Use	1
Storage and Disposal	1
Sprayer Cleanup	1
General Mixing Instructions	2
Application Precautions	2
General Application Instructions	2
Command Geographical Areas	2
SOYBEANS	2
General Information	2
Replanting Instructions	2
Pests and Weeds Controlled	2
Rotational Crop Guidelines	2
Command Applied in Combination	2
Tank Mixes	2
or Followed by Preemergence or Postemergence Treatments	2
PUMPKINS	3
SUCCELLENT PEAS	3
FALLOW LAND	3

SPECIAL PRECAUTION

Off-site movement of spray drift or vapor from Command 4 EC herbicide can cause leaf whitening or yellowing of surrounding crops. Prior to making applications, read and study the following items and application instructions on the label.

Physical/Chemical Hazards

The nature of storage and handling.

DIRECTIONS FOR USE

Information on methods of application, rates, timing, and sequence without labeling.

Do not apply this product to areas containing, or near, sensitive plants.

STORAGE AND DISPOSAL

Pesticide Storage

Do not freeze. If product is below 40°F. If solid crystals are observed, warm it to room temperature (60°F) by placing container in warm location. Stir well or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticide, fertilizer, water, food or feed by storage or disposal.

In case of spill, avoid contact. Isolate area and keep out animals and unprotected persons. Contain spills. Call FMC Collect (716) 735-3765.

To contain spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Returnable Refillable Sealed Containers

Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

IMPORTANT

FAILURE TO OBSERVE THE APPLICATION PRECAUTIONS SECTION OF THIS LABEL MAY RESULT IN INJURY TO DESIRABLE VEGETATION

- Desirable plants including some species of trees, shrubs, flowers, agronomic crops, and fruits and vegetables are sensitive to Command 4 EC herbicide.
- Foliar contact with spray, drift or vapors may cause whitening of sensitive plants. Symptoms are generally temporary in nature but may persist on some plants.

SPRAYER CLEANUP

Do not spray or flush equipment in or near or on trees or other plants, or in areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. Do not contaminate any body of water including irrigation water that may be used on other crops. Carefully follow sprayer clean up instructions noted below to prevent contaminating other crops from damage.

Sprayer equipment should be thoroughly flushed to remove residues of herbicide that may remain after the previous spray chores. The steps below should be followed in the order listed to clean sprayer equipment following application of Command 4 EC herbicide or tank mixes of Command 4 EC with other selected products.

1. Rinse any remaining spray solution from tank pump, hoses and nozzles and discard in an approved manner (See Note that follows).
2. Drain tank and flush by:
 - Thoroughly hose tank with a quantity of water equal to the total tank capacity and operating the pump to circulate the water through the sprayer system for 15 minutes.
 - Washing down the outside surfaces of equipment.
 - Removing nozzle tips and rinsing from end nozzle, in each three section and allowing several gallons of rinsate to flush complete. Thoroughly hose end rinsate while flushing.
3. Thoroughly drain remaining residue solution from tank, pump and hoses. Combine with bloom buildings and dispose of rinsates from first rinsing in an approved manner (See Note that follows).
 - When switching from one tank mix to another, flush tank with a quantity of water equal to the total tank capacity and operating the pump to circulate the water. A quantity of water equal to the total tank capacity should be flushed through the tank pump, hoses and bloom buildings. If application of Command 4 EC has just been completed in an adjacent tank, no rinsing is required to flush tank and pump.
4. Remove the remaining nozzle tips and screens and the filter screen and wash in a quantity of water equal to the total tank capacity.

Note: If, after the initial flushing of the spray tank, a second tank must be used, the same procedure is noted in #3 above.

5. If the tank used for rinsing is to be used for applying a postemergence herbicide, do not use the same tank to apply a crop for which Command 4 EC is registered. Another tank from the second rinsing may be utilized by flushing with the water for the next pesticide load.

HOWEVER

If the head use of the sprayer will be a postemergence applied per label on any crop, drain rinsate solution from this second rinsing. Return rinsate solution for use only with a second incorporated pesticide to be applied on a crop for which Command 4 EC is registered. Use tank (after draining second rinsate solution) in accordance with post-emergence product label directions.

NOTE: Dispose of excess spray mixture and/or rinsate from first tank rinsing by application and incorporation to cropland as described on this label. If excess spray mixture and/or rinsate from first rinsing cannot be disposed of according to label instructions, dispose of in compliance with local, state and Federal guidelines. Contact your state pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance.

GENERAL MIXING INSTRUCTIONS

Care must be taken when mixing Command 4 EC herbicide. Avoid mixing in areas adjacent to desirable plants (See Section headed "IMPORTANT").

Command 4 EC Alone: Mix Command 4 EC with water or liquid fertilizer in the following manner. Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 4 EC, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

Tank Mixtures: Fill spray tank one-fourth to one-third full with water with agitator operating; add the recommended amount of ingredients using the following order: dry formulations (e.g., wettable powders, dry flowables) first; liquid suspensions (e.g., flowables) next and finally, liquids (e.g., ECs). Mix thoroughly and fill tank one-half full continuous agitation. Add Command 4 EC herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Where use of a surfactant is recommended, add as the last ingredient to the spray tank. Maintain agitation during filling, mixing and application. When using drift-reducing agents, follow specific product label instructions for order of addition to spray tank.

Fertilizer Spray Mixtures: Applications of Command 4 EC alone or with recommended tank mixtures in conjunction with fertilizer solutions may be used unless use directions specifically state otherwise. Small quantities should be tested for compatibility by the following procedure in a small tankful quantity.

1. Fill a small tank with water.

2. Add the appropriate amount of herbicide based on the label directions. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first; liquid suspensions (e.g., flowables) next and finally, liquids (e.g., ECs).

Herbicide	Rate/Acre	Amount To Be Added Per Pint of Fluid Fertilizer
Wettable Powders	1 pint/1 acre	1 level teaspoon
Flowables	1/2 pint/1 acre	1 level teaspoon
Liquids	1/4 pint/1 acre	2 level teaspoons
Dry Flowables	3/4 pint/1 acre	1/4 level teaspoon
Dry Granules	3/4 pint/1 acre	1/4 level teaspoon
Emulsifiable Concentrates	1 pint/1 acre	1 level tablespoon
Liquids	1/2 pint/1 acre	2 level teaspoons
Flowables	1/4 pint/1 acre	3 level teaspoons


Empty Tank Spraying: Command 4 EC may be sprayed from an empty tank. Do not use a tank that has been used previously for Command 4 EC.

3. Clean, rinse and shake well.

4. Watch mixture for signs of separation. After 5 minutes, and again after 30 minutes, if the mixture has not completely recombined, add one quart of water to tank. If the mixture remains separated, do not use. If mixture becomes completely recombined, add one quart of water to tank.

If the mixture is incompatible, please notify the building manufacturer or the firm that supplied the tank.

1. **Command 4 EC Alone:** Fill tank with 21 pt. of water or fertilizer, add spray tank of herbicide with 1/2 pint per acre. Mix thoroughly during application.




- 2) **Command 4 EC plus Tank Mixtures:** Dilute individual products with two (2) parts of water, then add to the spray tank of fertilizer, while maintaining agitation, using the following order—slurry of dry formulations (wettable powders, dry flowables) first, diluted liquid formulations (EC s, flowables) second. Continue agitation during application.

APPLICATION PRECAUTIONS

Geographic Restrictions Command 4 EC must be applied only as a soil incorporated treatment in the following states, unless there is an approved SLN (24C) for counties within these states

Connecticut	Maryland	Ohio
Delaware	Massachusetts	Pennsylvania
Illinois	Michigan	Rhode Island
Indiana	Minnesota	Vermont
Iowa	New Hampshire	Virginia
Kentucky	New Jersey	West Virginia
Maine	New York	Wisconsin



*Does Not Apply to Fallow Land Applications or Counties for which an Approved SLN (24C) is on File

For Soil Incorporated Applications:
See Geographic Restrictions above

Do not apply Command 4 EC within 1,000 feet of the areas listed below

Towns and Subdivisions
Commercial Vegetable Production*
Commercial Fruit Production
Commercial Nurseries
Commercial Greenhouses
*except sweet corn

Plants such as the following may show symptoms of foliar whitening or yellowing if contacted by Command 4 EC herbicide. It is recommended that, prior to application, adjacent properties be checked and that spraying within 100 feet of such desirable plants be avoided.

Trees (Deciduous)	Trees (Evergreen)
Apple (inc. fruit & ornamental types)	Fir species
Ash (Green, White, Mountain)	Spruce species
Basswood	
Boxelder	Shrubs & Vines
Catalpa	Azalea
Cherry (inc. fruit & ornamental types)	Burningbush (Winged Euonymus)
Cottonwood	Grape
Elm	Honeysuckle
Ginkgo	Roses
Hawthberry	Yew
Mulberry	
Peach	Agronomic Crops
Pear (inc. fruit & ornamental types)	Wheat
Pecan	Oats
Poplar	
Russian olive	Vegetable and Flower plants
Tree of Heaven	
Tulip tree	Others
Walnut trees	Ferns
Willow species	Herbs
	Blueberry
	Raspberry
	Blackberry

Apply Command 4 EC herbicide only to soil incorporated treatments. Do not apply Command 4 EC herbicide to fallow land or areas including fencerows, willow trees, alfalfa, and other crops.

For Preemergent Applications. See Geographic Restrictions above

Except as restricted above (see Geographic Restrictions), Command may be used as a preemergent soil applied treatment without incorporation, however, additional precautions, beyond those required for a soil incorporated treatment, must be taken as follows:

Do not apply Command 4 EC within 1,500 feet of the following areas: Towns and Subdivisions, Commercial Fruit or Vegetable Production, Commercial Greenhouses or Nurseries.

Prior to application, adjacent properties must be checked and spraying within 1,500 feet of desirable plants must be avoided except sweet corn.

SPRAY DRIFT PRECAUTIONS

Non-target spray drift of Command 4 EC herbicide should be avoided to prevent whitening of desirable vegetation. Drift is influenced by many factors which include wind speed, spray pressure, particle size, nozzle type, and boom height.

- Do not apply when weather conditions favor drift. If wind speeds exceed 10 miles per hour, a drift reducing additive must be used.
- A minimum spray volume of 10 gallons per acre is recommended with appropriate nozzle types and sizes that produce coarser sprays.
- The use of agriculturally approved drift reducing additives is recommended for application volumes of 15 to 40 gallons per acre when spraying in the proximity of desirable plants (see list above).
- The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre.
- Use minimum nozzle pressure and boom height while maintaining uniform spray pattern.
- Do not exceed 40 psi spray pressure.
- Selection and proper use of spray equipment is critical in minimizing spray drift. The following table suggests pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types.

Suggested Nozzle Types, Minimum Size and Recommended Pressure Ranges for Minimizing Drift

Nozzle	Pressure Range (PSI)	Minimum Flow Rate Within Pressure Range (GPM)	Minimum Nozzle Size
Flat-fan	15-30	0.3	#4*
LP-flat-fan	10-25	0.3	#3
Even flat-fan**	15-30	0.3	#4
Flood	10-25	0.3	#2.5*
Whirl chamber	5-20	0.3	#5
Raindrop	15-40	0.15	#2
Wide angle full cone	15-40	0.3	#1

*Refers to EC number 10, 11, 12, 13 or 14.

**Recommended for banded application.

ROTATIONAL CROPPING PRECAUTIONS: Temporary whitening or yellowing of leaves may occur on unrotated rotational crops where undesirable soil residues of Command 4 EC herbicide exist. These soil conditions may result in undesirable soil residue dues and greater potential liability to rotation crops.

- 1) Exceeding legal maximum drift rates.
- 2) Overlapping or straying of drift from low boom angles or overlapping of spray swathes failing to shut off spray boom when turning end row areas. See use of stepped sprayer.

Refer to Rotational Guidelines and replanting instructions of specific crops for additional crop planting information.

SPECIAL PRECAUTION

If site movement of spray drift occurs, the herbicide can cause foliar whitening or yellowing of sensitive crops. Prior to making applications, it is important to know the direction and distance of spray drift potential.

GENERAL APPLICATION INSTRUCTIONS

Do not apply aerially or through irrigation equipment.

Ground Applications

Broadcast or Banded Applications: Broadcast or banded applications are recommended for most situations. Broadcast applications should be made with a ground sprayer equipped with a ground boom or a ground sprayer unit.

Note: The use of ground sprayers is not recommended for broadcast applications.

BEST AVAILABLE COPY

agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre. Use nozzles suitable for broadcast or banded application of herbicides. Command spray can be applied to a maximum of the band times that the spray is applied. AERIAL ATTITUDE: Use At least 10 feet above ground level and 100 feet from the spray drift. For Command 4 EC tank mixtures with wettable powders or dry herbicides, application distance between plant rows should be no greater than 10 inches.

Banded Application: Calculate the rates and volume required by using the following formula:

Band width in inches	Band length in feet	Band rate per acre
Row width in inches	Acre	
Band width in inches	Broadcast Volume Per Acre	Band volume per acre

Follow broadcast minimum gallooning requirements adjusted to band width. Do not exceed 40 psi spray pressure.

Refer to Spray Drift Precautions for suggested pressures, flow rates and nozzle sizes for drift reduction using various nozzle types. For banded applications an even flat-fan nozzle is recommended.

Soil Incorporated Application: Apply Command 4 EC to the soil surface and uniformly incorporate. Select equipment suited for proper incorporation. Shallow incorporation normally results in better seed control than a deeper incorporation.

In states where soil incorporation is required (see Geographical Restrictions) application and immediate incorporation to a maximum depth of 3 inches is required unless the soil is dry. On dry soils incorporation to a maximum depth of 3 inches must be completed within 6 hours of Command 4 EC herbicide application.

Soil must be in good tilth to allow for thorough mixing of the soil. Application to wet soils will increase the potential for off-site movement of Command 4 EC herbicide vapors and may result in poor soil incorporation and unsatisfactory weed control.

INCORPORATION EQUIPMENT

ONE PASS INCORPORATION

Suggested Implements

For one pass incorporation use C or S share field cultivators equipped with a leveling device such as a spike or coil-tine harrow or culturing basket attachment. Field cultivator must have 3 or 4 rows of sweeps spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Equipment design including any drag attachments must be adequate to avoid soil binding which may result in streaks and reduce weed control. The implement should be operated at 6 to 7 mph and 1 to 1.5 ft. 3 ft. or 4 ft. deep depending on soil texture.

Cultivator equipment may require a tillage device or not. If so, it is used at a single pass, it should be operated at a depth of 2 to 4 inches and a minimum speed of 6 mph.

PTO drive seed bed preparation equipment should be set to cut 2 inches deep and operated at 4 mph or less.

TWO PASS INCORPORATION

Suggested Implements

When using two passes incorporate the second pass at a 90° angle to the first pass.

Disc set to cut 3 to 4 inches deep at 4 to 6 mph. Blades should be less than 20 inches in diameter and spaced 9 inches or less apart.

Rolling cultivator set to cut 2 to 3 inches deep at 6 to 8 mph, rule on coarse or medium textured soils only.

Fans cut tines with 3 to 4 rows of sweeps spaced at 7 to 10 inches or less and set to cut 3 inches deep at 5 mph or greater.

Cultivation After Planting

Soil treated with Command 4 EC may be shallowly cultivated by hand or hand-hoe without requiring the wheel control activity of Command 4 EC. Do not cultivate deeper than the Command 4 EC treated layer of soil since this may bring untreated soil to the surface and poor weed control may result.

COMMERCIAL IMPREGNATION AND APPLICATION OF COMMAND 4 EC ON DRY BULK FERTILIZERS

Command herbicide may be impregnated on dry bulk fertilizers. When applied directly, Command dry bulk fertilizer mixture should weigh 20% less than that provided by the same amount of Command 4 EC per ton of water.

The Command fertilizer mixture must be stored in bags. See bag storage listed under General. Furthermore, the Application Rate section of this label.

Impregnation: Apply using a minimum of 100 pounds of dry bulk fertilizer per acre and up to a maximum of 400 pounds per acre with the recommended amount of Command 4 EC herbicide per acre. Use a standard rotary drum mixer or a similar type of coater/binder designed with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform fine spray pattern over the tumbling fertilizer to thorough coverage. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Command provides a satisfactory dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry flowing mixture. Microcel E (Dolica Marine Product Corporation) is a recommended absorptive powder. Generally less than 2% by weight of Microcel E is required. DO NOT impregnate Command 4 EC onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with Command.

The amount of Command actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with Command 4 EC herbicide should be applied immediately, not stored. Care should be taken to ensure spreading procedures in the field do not place fertilizer treated with Command in areas which cannot be incorporated, if incorporation is required within the Geographic Restrictions of this label.

All state regulations, labeling, etc., of the mixtures are the responsibility of the seller.

For those rates not listed in the following table, calculate the amount of Command to be impregnated on a ton of dry bulk fertilizer using the following formula:

2000 pounds dry fertilizer per acre	PINTS of Command (recommended rate for soil texture) per acre	PINTS of Command per ton of fertilizer
11.75 lbs A.I.	1 pt	1 pt
11 lbs A.I.	2 pt	2 pt
11.25 lbs A.I.	2.5 pt	2.5 pt

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZERS WITH COMMAND 4 EC HERBICIDE

PINTS OF COMMAND 4 EC PER TON OF FERTILIZER

Fertilizer Rate lbs/Acre	Command 4 EC Rate Per Acre 1 pt. (75 lbs A.I.)	Command 4 EC Rate Per Acre 2 pt. (1 lb A.I.)	Command 4 EC Rate Per Acre 2.5 pt. (1.25 lbs A.I.)
200	15	20	25
250	12	16	20
300	10	13	16
350	8	11	14
400	7	10	12
450	6	9	11


Fertilizer materials are excess, very dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will not distribute during application. Crop injury and/or plant control may occur where the impregnated fertilizer is not uniformly applied.

SPECIAL PRECAUTIONS FOR DRY BULK FERTILIZER IMPREGNATION AND APPLICATION WITH COMMAND 4 EC

- All equipment used to apply Command should be thoroughly cleaned immediately following use to remove any contaminant of residue which could cause injury to non-target vegetation or desirable vegetation. Refer to Sprayer Cleaning section for additional details and disposal information.
- Equipment used to physically transport Command treated fertilizer, including boat, trailer, conveyor bins, etc. should be covered to prevent loss of the product and subjected to cleaning procedure previously described.
- Mixing of Command herbicide and dry fertilizer should be conducted in a tank or wheelbarrow. It is difficult to attain an even application in an auger.
- Bulk containers should be tightly closed while the product is transported and applied in bulk. Check for damage to bags or bulk product bins.
- It is recommended that dry bulk fertilizer application operations using Command 4 EC be conducted with a crew of no less than four people, where Command symptoms of any nearby vegetation will not be tolerated.

COMMAND GEOGRAPHICAL AREAS

E. Rates, Weed Control, and Crop Rotation



Southern area includes Missouri border, which incorporates the following counties: Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott, and Stoddard.

SOYBEANS

GENERAL INFORMATION

1 and 4 EC herbicide may be utilized as a preemergent soil applied or soil incorporated agent for the control of annual grass and broadleaf weeds in soybeans.

Command 4 EC must be soil incorporated in the following states unless an approved 24(C) is on file for counties within that state (check with your local dealer; Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, Wisconsin). Command 4 EC herbicide may be tank mixed with other herbicides to broaden the weed control spectrum compared to the products applied alone. Command 4 EC may be tank mixed with Canopy® (Dual® Lasso), Lurex® (Lexone® Preview®), Prowl®, Scepter®, Sencor®, Sonalan®, Squadron®, Triallan®, Tri-Scept®, or other registered herbicides. Observe all precautions, instructions,

and rotational cropping guidelines of each product's label when tank mixing. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with the herbicides listed above, unless use directions specifically state otherwise.

REPLANTING INSTRUCTIONS

Initial seeding of soybeans fails to produce a stand, soybeans may be replanted in fields treated with Command 4 EC alone (or with recommended tank mixtures). Do not reseed field with a second application of Command 4 EC. When tank mixed with a labeled product refer to the soybean replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the 'Rotational Guidelines' on this label.

Northern Area—See Map

COMMAND 4 EC HERBICIDE APPLIED ALONE

RATES AND WEEDS CONTROLLED^t

	1.0 PT. (0.50 LB A.I.)	1.5 PT. (0.75 LB A.I.)	2.0 PT. (1.0 LB A.I.)
GRASSES	GRASSES	GRASSES	GRASSES
Barnyardgrass	Barnyardgrass	Barnyardgrass	Barnyardgrass
Cogon Grass	Cogon Grass	Cogon Grass	Cogon Grass
Lambsquarters	Lambsquarters	Lambsquarters	Lambsquarters
Smooth Pigweed	Smooth Pigweed	Smooth Pigweed	Smooth Pigweed
Fleabane	Fleabane	Fleabane	Fleabane
Common Purslane	Common Purslane	Common Purslane	Common Purslane
Common Glasswort	Common Glasswort	Common Glasswort	Common Glasswort
Common Amaranth	Common Amaranth	Common Amaranth	Common Amaranth
Common Bindweed	Common Bindweed	Common Bindweed	Common Bindweed
Common Waterlettuce	Common Waterlettuce	Common Waterlettuce	Common Waterlettuce
Black Seeded Plantain			

BROADLEAVES

Velvetleaf
Spurred Anoda

plus
Galinsoga
Lambsquarter
Prickly Sida
Purslane
Venice Mallow
Common Ragweed^{*}
Jimsonweed
Smartweed
-Pennsylvania
-Lady's thumb*

* Partially Controlled

BROADLEAVES

Velvetleaf
Spurred Anoda

plus
Galinsoga
Lambsquarter
Prickly Sida
Purslane
Venice Mallow
Common Ragweed^{*}
Jimsonweed
Smartweed
-Pennsylvania
-Lady's thumb*

BROADLEAVES

All broadleaves
controlled at 1.5
pints plus
Common Ragweed
Jimsonweed
Kochia
Tropic Croton
Cocklebur^{*}
Smartweed
-Pennsylvania
-Lady's thumb*

¹Higher label rates (a minimum of 2 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types.

²For heavy weed pressure or heavy soils use 1.25 pints per acre to control Velvetleaf or Spurred Anoda.

NOTE: When using Command 4 EC more than 30 days prior to planting, the highest label rate should be used.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur.

1.0 PT. (0.50 LB A.I.)	1.5 PT. (0.75 LB A.I.)	2.0 PT. (1.0 LB A.I.)	For 1st A.I.
<u>ANYTIME</u>	<u>ANYTIME</u>	<u>ANYTIME</u>	
Soybeans	Soybeans	Soybeans	
Pumpkins	Pumpkins	Pumpkins	
Peas			
<u>9 MONTHS</u>	<u>9 MONTHS</u>	<u>9 MONTHS</u>	
Corn	Same as 1.0 pint, plus Peas	Same as 1.5 pints, except Seed Corn and Sweet Corn require a 12 month rotation	
—Field			
—Pop			
—Seed			
—Sweet			
Cotton			
Cucurbits			
Dr. Beans			
Peanuts			
Peppers			
Potatoes			
Rice			
Snap Beans			
Sorghum			
Sugar Beets			
Sweet Potatoes			
Tobacco			
Tomatoes (Transplanted)			
<u>12 MONTHS</u>	<u>12 MONTHS</u>	<u>12 MONTHS</u>	
Cabbage	Same as 1.0 pint	Same as 1.5 pints plus	
Tomatoes (all)		Seed Corn ‘Sweet’ Corn	
Wheat			
<u>15 MONTHS</u>	<u>16 MONTHS</u>	<u>16 MONTHS</u>	
All crops	All crops	All crops	

Cover crops however, may be planted anytime but stand reduction may occur in some areas. Do not graze or harvest for feed or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

BEST AVAILABLE COPY

**Southern Area—See Map
COMMAND® 4 EC HERBICIDE
APPLIED ALONE**

RATES AND WEEDS CONTROLLED^t

	10 PT. (0.50 LB A.I.)	15 PT. (0.75 LB A.I.)	20 PT. (1.0 LB A.I.)	25 PT. (1.25 LB A.I.)
GRASSES	GRASSES	GRASSES	GRASSES	GRASSES
Suppression of some annual grasses	Barnyardgrass Bentgrass Signalgrass Crabgrass - Large - Smooth Foxtail - Giant - Green Goosegrass Panicum Common Fall - Texas Seedling Johnsongrass Field Sandbur [†]	Bentgrass Controlled at 1.5 pints plus Cupgrass - Eastern - Western - Wavy Field Sandbur Bermudagrass Red Rice [†] Itchgrass ^{††}	Amaranth - Intermediate - Upinyard Fatgrass Paspalum Rabbitfoot Vinegar St. Augustine Weed & Grass Millet	

BROADLEAVES BROADLEAVES

Velvetleaf	Velvetleaf [†]
Spurred Anoda	Spurred Anoda [†] plus
Lambquarters	Cocklebur
Pickly Sida	Common Ragweed
Purslane	Dayflower
Reed	Florida pusley
Venice Mallow	Jimsonweed
Common Ragweed [†]	Tropic Croton
Jimsonweed [†]	Wild Poinsettia ^{††}
Pennsylvania Smartweed [†]	Kochia [†]
	Redvine [†]
	Florida
	Beggarweed

* Partially Controlled

^t Higher label rated (a minimum of 2 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, a minimum application rate of 2.5 pints per acre for cocklebur or common ragweed control is recommended.

* Use in AR LA MS TX OK on, —for high pressure and heavy soils use 2.5 pms

** For heavy weed pressure or heavy soils use 1.25 pints per acre to control Velvetleaf or Spurred Anoda.

*** For control of Itchgrass, do not incorporate Command 4 EC. For high weed pressure and heavy soils use 2.5 pints

NOTE: When using Command 4 EC more than 30 days prior to planting, the highest label rate should be used

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur

1.0 PT. (0.50 LB. A.I.)	1.5 to 2.0 PT. (0.75 to 1.0 LB. A.I.)	2.5 PT. (1.25 LB. A.I.)
ANYTIME	ANYTIME	ANYTIME
Soybeans Pumpkins Peas	Soybeans Pumpkins	Soybeans

9 MONTHS

Corn	9 MONTHS	9 MONTHS
—Field —Pop —Seed —Sweet	Same as 1.0 pint plus Peas	Same as 1.5 to 2.0 pints plus Pumpkins

Cotton

Cucurbits

Dry Beans

Peanuts

Potatoes

Pumpkins

Rice

Snap Beans

Sorghum

Sugar Beets

Sweet Potatoes

Tobacco

Tomatoes

(Transplanted)

12 MONTHS

A crops

NOT AVAILABLE CROP[†]

Cover crops however may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock

**COMMAND® 4 EC HERBICIDE
APPLIED IN COMBINATIONS FOR SOYBEANS**

Command 4 EC herbicide may be tank mixed with other soybean herbicides to broaden the weed control spectrum, compared to the product applied alone. In the sections that follow, FMC recommends a combination of products and application rates to broaden weed control spectrum. FMC makes no claims as to weed control for any combination of rates or products other than as recommended in this label. Additional recommendations for rate combinations with other herbicides not included in the following sections may become available subsequent to the date of publication for this label. Please check with your dealer, more recent FMC technical bulletins and label supplements. When applying mixtures of Command 4 EC with other herbicides, observe all precautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and repainting instructions of each product's label when using tank mixes.

**Command 4 EC Herbicide plus Canopy® Herbicide—
Northern & Southern Areas**

Look this chart out for broadened weed control spectrum.

Broadcast Rates Per Acre[†]

Soil Texture	Command		
	North	South	Canopy
[†] 1" to 6" Organic Matter			
Coarse (light) Soils (loamy sand, sandy loam)	1 to 2 pints	2 to 2 pints	4 to 6 oz
Medium Soils (loam, silty soil, loamy sandy clay, sandy clay loam)	1 to 2 pints	2 to 2 pints	8 to 8 oz
Fine (heavy) Soils (clay, clay loam, silty clay loam)	1 to 2 pints	2 to 2 pints	16 to 16 oz

[†] Broadcast rates in pints per acre correspond to the rates in Table 1 with the following:

1 pt. = 1 lb. of product. 1 lb. = 1 quart. 1 quart = 16 oz.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Cocklebur[†]
Pigweed (Redroot, Smooth, Hemp, Sesbania)
Morningglory (annual ivy, leaf, entire leaf, pitted leaf, sicklepod)
Copperleaf (hophornbeam)
Plantain, Common

[†] For best weed control, apply Command 4 EC with Canopy® Herbicide. Both herbicides will control most weeds. Some Morningglory species require a cutback application of Canopy® Herbicide to obtain control.

When applying mixtures of Command 4 EC plus Canopy® Herbicide, the broadcast rates are determined by the rate of each product. Special attention must be paid to sections requiring planting-depth, phosphate, pesticide interaction, previous pesticides, and selected soybean varieties on the Canopy® label. Observe application precautions, rotational guidelines and repainting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Dual® 8 E Herbicide—
Northern & Southern Areas**

Look this chart out for broadened weed control spectrum.

Broadcast Rates Per Acre[†]

Texture	Command		
	North	South	Dwarf
[†] 1" to 6" Organic Matter			
Coarse (light) Soils (loamy sand, sandy loam)	1 to 2 pints	2 to 2 pints	4 to 6 oz
Medium Soils (loam, silty soil, loamy sandy clay, sandy clay loam)	1 to 2 pints	2 to 2 pints	8 to 8 oz
Fine (heavy) Soils (clay, clay loam, silty clay loam)	1 to 2 pints	2 to 2 pints	16 to 16 oz

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following

Red Rice
Witchgrass
Nutsedge

* Use higher rates in areas with heavy grass.

When applying mixtures of Command 4 EC plus Lasso or Lasso Micro-Tech herbicide will broaden the weed control spectrum compared to either product used alone

Command 4 EC Herbicide plus Lasso® or Lasso® Micro-Tech—Northern & Southern Areas

A tank-mix combination of Command 4 EC herbicide and Lasso or Lasso Micro-Tech herbicide will broaden the weed control spectrum compared to either product used alone

Product

Broadcast Rates Per Acre*

Soil Texture	Command North	South	Lasso® Lasso Micro-Tech
Coarse (light) Soils (loamy sand, sandy loam)	1½ to 2 pints	2 to 2½ pints	2 to 2½ quarts
Medium Soils (loam, silt, silty loam, sandy clay, sandy clay loam)	1 to 2 pints	2 to 2½ pints	2 to 3 quarts
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	2 to 3 quarts

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., sand, loamy sand, sandy loam) within a textural group

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following
Carpetweed
Pigweed
Black Nightshade

When applying mixtures of Command 4 EC herbicide plus Lasso or Lasso Micro-Tech, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational crop grazing and feeding restrictions and replanting instructions of each product's label when using tank mixtures

COMMAND 4 EC HERBICIDE PLUS LOROX®

A preemergence tank mix combination of Command 4 EC herbicide and Lorox herbicide will broaden the weed control spectrum compared to either product alone

The tank mix combination may not be incorporated. Where Command 4 EC must be incorporated (see Geographical Restrictions), apply Lorox as an overlay treatment.

Broadcast Rates Per Acre*

Soil Texture	Command	Lorox DF†
Coarse (light) Soils (loamy sand, sandy loam)	1 to 2 pints	2 to 2½ pints
Medium Soils (loam, silt, silty loam, sandy clay, sandy clay loam)	1 to 2 pints	2 to 2½ pints
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1 to 2 pints	2 to 2½ pints

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silty loam, sandy clay, sandy clay loam) within a textural group

† Plant seed at least 1½ inches deep or Lorox may result. Refer to Lorox label for other planting and application precautions

Deeper than 1½ inches, Lorox may result. Refer to Lorox label for other planting and application precautions

One pint of Lorox DF equals one pint of Lorox DF, use equivalent rates

WEEDS CONTROLLED

Glassy Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following

Pigweed
Carpetweed
Chickweed

When applying mixtures of Command plus Lorox observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop Grazing and Feeding Restrictions and Replanting Instructions sections of this label when using tank mixtures

COMMAND 4 EC HERBICIDE PLUS LOROX PLUS

A preemergence tank mix combination of Command 4 EC herbicide and Lorox herbicide will broaden the weed control spectrum compared to either product alone

The tank mix combination may not be incorporated. Where Command 4 EC must be incorporated (see Geographical Restrictions), apply Lorox as an overlay treatment.

Broadcast Rates Per Acre*

Soil Texture	Command		Lorox DF†
	North	South	Pints
Coarse (light) Soils (loamy sand, sandy loam)	1 to 2 pints	2 to 2½ pints	9 to 12 oz
Medium Soils (loam, silt, silty loam, sandy clay, sandy clay loam)	1 to 2 pints	2 to 2½ pints	6 to 8 oz
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1 to 2 pints	2 to 2½ pints	11 to 17 oz

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silty loam, sandy clay, sandy clay loam) within a textural group

Deep or Lorox more than 1½ inches may result. Refer to Lorox label for other planting and application precautions

Deeper than 1½ inches, Lorox may result. Refer to Lorox label for other planting and application precautions

WEEDS CONTROLLED

Glassy Weeds

All species listed under Command Applied Alone, Weeds Controlled section

Plus the following

Broadleaf Weeds

All species listed under Command Applied Alone, Weeds Controlled section

Plus the following

Carpetweed, Cocklebur, Mustard, Pigweed, Redroot, smooth Black Nightshade, Morningglory (annual), Purslane, Turnip leaf, Turnip leafy, Giant Ragweed

† Partially Controlling

Large seeds treated with Lorox may result. Refer to Lorox label for other planting and application precautions

When applying mixtures of Command plus Lorox observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop Grazing and Feeding Restrictions sections of this label when using tank mixtures

BEST AVAILABLE COPY

Command 4 EC herbicide plus Metribuzin (Sencor[®] or Lexone[®])—Northern Area Only

A tank mix combination of Command 4 EC herbicide and metribuzin will broaden the weed control spectrum compared to either ~~Product~~ used alone.

Soil Texture**	Broadcast Rates Per Acre [†]	
	Command	Sencor DF [*] Lexone DF
Coarse (light) Soils (loamy sand ^{**} , sandy loam)	1½ to 2 pints	½ to 1½ lb
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	½ to 1½ lb
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	½ to 1½ lb

- Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- For pints Sencor 4 or Lexone 4L multiply pounds DF by 1.5.
- Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Carpetweed
Pigweed^{*} (Redroot, Smartweed)
Wild mustard

* Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC herbicide plus Metribuzin (Sencor[®] or Lexone[®])—Southern Area Only

A tank mix combination of Command 4 EC herbicide and metribuzin will broaden the weed control spectrum compared to either ~~Product~~ used alone.

Soil Texture**	Broadcast Rates Per Acre	
	Command	Sencor DF [*] Lexone DF
Coarse (light) Soils (loamy sand ^{**} , sandy loam)	2 to 2½ pints	½ to 1½ lb
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	2 to 2½ pints	½ to 1½ lb
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	2 to 2½ pints	½ to 1½ lb

- Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- For pints Sencor 4 or Lexone 4L multiply pounds DF by 1.5.
- Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur. Use Sencor 4L and metribuzin on silty clay and heavy clay soils of the Mississippi Delta.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Carpetweed
Hemp sesbania (coffeeweed)
Pigweed (Redroot, Smartweed)
Wild mustard
Sicklepod^{*}

* Reduced Competition

* Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Preview[®]—Northern Area Only

A tank-mix combination of Command 4 EC herbicide and Preview herbicide will broaden the weed control spectrum compared to either ~~Product~~ used alone.

Product

Broadcast Rates Per Acre

Soil Texture	Command	Preview [*]
Coarse (light) Soils (loamy sand, sandy loam)	1½ to 2 pints	½ to 1½ lb
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	½ to 1½ lb
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	½ to 1½ lb

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

Do not use Command 4 EC in combination with Preview on soils with less than 0.5% organic matter, or on soils with pH greater than 7.5 or on soils having a calcareous surface area or a pH of 7.5 or higher as Preview may injure soybeans.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Carpetweed
Common spotted
Copperleaf, nophornbeam
Mustards
Pigweed
Cocklebur
Sunflower^{*}
Burcupimbush
Giant Ragweed^{*}
Morningglory (Amaranthus)
Ivyleaf^{*}, Pitted Erigeron^{*}

* NOTE: Control of morningglory, sunflower, cocklebur and giant ragweed may be erratic depending on soil type, temperature, depth of weed seed in the soil, and the amount and timing of soil moisture. Control may be improved with timely cultivation.

When applying mixtures of Command 4 EC plus Preview, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

BEST AVAILABLE COPY

Command 4 EC Herbicide plus Prowl® 4 EC Herbicide—

Northern & Southern Areas

A tank mix combination of Command 4 EC and Prowl will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre		
	Command North	South	Prowl 4 EC*
Coarse (light) Soils (loamy sand, sandy loam)	1½ to 2 pints	2 to 2½ pints	1 to 1½ pints
Medium Soils (loam, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	2 to 2½ pints	2 pints
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	2 to 2½ pints

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g. loam, silt loam, sand, clay, sandy clay loam) within a textural group.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Pigweed (Redroot, Smooth)
Carpetweed

When applying mixtures of Command 4 EC plus Prowl, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Scepter® Herbicide— Northern Area Only

See Scepter® label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

Note: The Scepter label does not allow rotation to Corn, Wheat, Barley, or Oats prior to 18 months after last Scepter application when Scepter is applied preemergent or preplant incorporated within the Northern Use Area as defined on the Scepter label. See Scepter label for appropriate geographic restrictions.

Soil Texture	Broadcast Rate Per Acre	
	Command	Scepter®
Light Soils	1½ to 2 pints	1 to 1½ pints

* Use higher rates when heavier weed pressure is anticipated.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Buckwheat
Cocklebur
Pigweed spp.
Common Purslane
Giant Hogweed

* Partially controlled.

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than two inches, or make additional Command 4 EC or Command Scepter tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

When applying mixtures of Command 4 EC plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide p Southern Area Only

See Scepter label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Scepter®
All Soil Types	2 to 2½ pints	½ to 1 pint

Scepter Herbicide—

Geographic restrictions

4 EC and Scepter will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Scepter®
All Soil Types	2 to 2½ pints	½ to 1 pint

* Use higher rates when heavier weed pressure is anticipated.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Note: S anticipated.

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

Buckwheat
Cocklebur
Pigweed spp.
Wild poinsettia

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than two inches, or make additional Command 4 EC or Command Scepter tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

In other conditions, soybeans can be replanted deeper than two inches, or make additional Command 4 EC or Command Scepter tank mix applications.

Important: For optimum tank mix applications, it is recommended that soybeans be replanted no deeper than two inches.

When applying mixtures of Command 4 EC plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

COMMAND 4 EC Herbicide ; Northern Area Only

A tank mix combination of Command 4 EC will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rate Per Acre	
	Command	Scepter®
Coarse (light) Soils (loamy sand, sandy loam)	1 to 1½ pints	1 to 1½ pints

Soil Texture	Broadcast Rate Per Acre	
	Command	Scepter®
Medium Soils (loam, silt loam, silty loam, sandy clay, sandy clay loam)	1 to 2 pints	1½ to 2 pints

Soil Texture	Broadcast Rate Per Acre	
	Command	Scepter®
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	2 to 3 pints

* Select lower to higher rates within the ranges noted based on the coarser soil types within the texture of a given soil. Rates within the range of a given soil type may be used for lighter or heavier weed pressure.

Note: Scepter label indicates that Scepter is partially controlled by Buckwheat, Cocklebur, Pigweed spp., and Wild poinsettia.

BEST AVAILABLE COPY

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following

Carpetweed

Pigweed (Redroot, Smooth)

When applying mixtures of Command 4 EC plus Sonalan observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Squadron—Northern Area Only

See Squadron label for appropriate geographic restrictions

A tank-mix combination of Command 4 EC and Squadron will broaden the weed control spectrum of either product used alone

Soil Texture	Broadcast Rates Per Acre	
	Command	Squadron*
All Soil Types	1½ to 2 pints	1½ to 2½ pints

* Do not exceed 1½ pints in Northern Area (as defined on the Squadron label). Use higher rate of Squadron if areas of heavy rockbur or shale.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following

Black Nightshade
Cocklebur
Pigweed spp (Common Sunflower)

Giant Ragweed*

* Partially controlled

When applying mixtures of Command 4 EC plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Squadron—Southern Area Only

See Squadron label for appropriate geographic restrictions

A tank-mix combination of Command 4 EC and Squadron will broaden the weed control spectrum of either product used alone

Soil Texture	Broadcast Rates Per Acre	
	Command	Squadron*
All Soil Types	2 to 2½ pints	2 to 3 pints

* Use higher rate of Squadron if areas of heavy rockbur or shale.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following

Black Nightshade
Cocklebur
Pigweed spp (Common Sunflower)
Giant Ragweed*

When applying mixtures of Command 4 EC plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Trellan® EC (Trifluralin)—Northern & Southern Areas

A tank-mix combination of Command 4 EC herbicide and Trellan® EC herbicide will broaden the weed control spectrum compared to either product used alone.

Soil Texture	Broadcast Rates Per Acre		
	Command North	South	Trellan® EC
Coarse (light) Soils (loamy sand, sandy loam)	1½ to 2 pints	2 to 2½ pints	1 to 1½ pints
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	2 to 2½ pints	1 to 1½ pints
Fine (heavy) Soils (silt, clay, clay loam, silty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	1 to 1½ pints

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g. loam, silt loam, sandy clay, sandy clay loam) within a texture group.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following

Carpetweed
Chickweed
Field Bindweed
Pigweed (Redroot, Smooth)

When applying mixtures of Command 4 EC plus Trellan® EC observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Tri-Scept Herbicide—Northern Area Only

See Tri-Scept® label for appropriate geographic restrictions

A tank-mix combination of Command 4 EC and Tri-Scept applied to incorporated soil will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Tri-Scept®
All Soil Types	1 to 1½ pints	1 to 1½ pints

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g. loam, silt loam, sandy clay, sandy clay loam) within a texture group.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following

Black Nightshade
Cocklebur
Pigweed spp (Common Sunflower)
Giant Ragweed*

* Partially controlled

When applying mixtures of Command 4 EC plus Tri-Scept® herbicide observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Tri-Scept—Southern Area Only

See Tri-Scept® label for appropriate geographic restrictions

A tank-mix combination of Command 4 EC and Tri-Scept applied to incorporated soil will broaden the weed control spectrum of either product used alone.

BEST AVAILABLE

847

Broadleaf Weed	Per Acre	
	Command	Tri-Scepter
All broadleaf weeds	1/2 to 2 pints	

* Use the rate of one product per acre for broadleaf weed control in soybeans and other oilseed rotations.

WEEDS CONTROLLED

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:

- Cocklebur Black nightshade
- Pigweed spp. Common sunflower
- Wild poinsettia

When applying mixtures of Command 4 EC plus Tri-Scepter observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY PREEMERGENCE TREATMENTS

Preemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of additional weed species listed on the preemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information that appear on the product labels. **DO NOT** apply compounds containing metribuzin after a soil incorporated Command 4 EC plus Lexone® or Sencor® tank mix treatment unless split applications are used where the pounds of active ingredient do not exceed the total allowed.

DO NOT apply Canopy® Preview® Pursuit® Plus® Scepter® Squadron® or Tri-Scepter® herbicides after Command 4 EC plus any of those products in a tank mix treatment as these other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY POSTEMERGENCE TREATMENTS

Postemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of emerged weeds as listed on the postemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information that appear on the product labels.

DO NOT apply Canopy® Preview® Pursuit® Plus® Scepter® as a postemergence treatment after Command 4 EC plus Canopy® Preview®, Pursuit®, Plus®, Scepter®, Squadron®, or Tri-Scepter® in a tank mix treatment unless the other product labels specifically recommend such a treatment as these other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

COMMAND 4 EC HERBICIDE FOLLOWED BY ACIFLUORFEN (BLAZER® 2L OR TACKLE®) HERBICIDE

Following an application of Command 4 EC or tank mixture with Command 4 EC, a postemergence application of acifluorfen will control the following emerged broadleaf weeds:

Type of Weeds

Cocklebur
Hemp-stinknet
Fogwillow-weed
Black nightshade
Wild mustard
Purple Morning glory
Pitted Morning glory

Apply acifluorfen at a rate of 1 pint per acre with 25% crop oil concentrate to actively growing weeds at no more than the 4 leaf growth stage (do not count cotyledons). Count only the true leaves developed from the true leaf axils. Count the true leaf axils of all stages of the first to third internode of the true leaf.

NOTE: Application of Command 4 EC herbicide generally retards the development rate of weeds which may extend the period where acifluorfen can be effectively applied.

Read and follow all preadmixing restrictions and warnings on all product labels.

COMMAND 4 EC HERBICIDE FOLLOWED BY SCEPTER® HERBICIDE

Following an application of Command 4 EC or tank mixture with Command 4 EC, a postemergence application of Scepter herbicide will control the following emerged broadleaf weeds:

Broadleaf Weeds

Cocklebur common

Pigweed

Wild poinsettia

Apply Scepter after crop emergence but before weeds exceed 6 to 12 inches tall for wild poinsettia. Apply at a broadcast rate of 1 pint per acre. Apply when weeds are actively growing. DO NOT apply Scepter postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes. The total amount of Scepter should not exceed 1 pint per acre per season in northern areas or 1/2 pint per acre per season in southern areas as defined on the Scepter label. Squadron and Tri-Scepter both include the same active ingredient as Scepter.

For postemergence applications, the addition of a nonionic surfactant or crop oil concentrate is required. The nonionic surfactant should contain at least 80% active ingredient and should be applied at a rate of 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate stated on the COC label.

Read and follow all precautionary statements and restrictions, directions for use and all other information appearing on the Scepter label.

COMMAND 4 EC HERBICIDE APPLICATION—MINIMUM TILLAGE OR NO-TILLAGE SOYBEANS

Command 4 EC herbicide may be used alone or in combination with Sencor® Lexone® Lorox® or Lorox Plus herbicides, plus Roundup® Roundup® herbicide for preemergence (to crop and weeds) or postemergence (to weeds) but prior to crop emergence control of broadleaf and grass weeds where soybeans will be planted into a cover crop in previous crop residues (e.g., corn, small grain stubble).

WEEDS CONTROLLED

When used as directed, the paraquat in the tank mix controls most emerged annual weeds and suppresses many perennial weeds. Roundup® Roundup® herbicide controls emerged annual and perennial weeds when applied as directed on the Roundup® label. Command 4 EC or Command 4 EC plus Sencor® Lexone® Lorox® provides preemergence control of the weeds listed in the Command 4 EC Applied Alone and Command 4 EC Applied in Combinations sections of this label.

APPLICATION AND MIXING INSTRUCTIONS

Paraquat Combinations: Tank mix Command 4 EC alone or with Sencor® Lexone® Lorox® Lorox Plus observe mixing instructions in the General Mixing Instructions portion of this label, as well as individual product label use directions. Add 1 to 2 pints paraquat (based on product containing 2 pints active ingredient per gallon) per acre. Use lower rate for annual weeds less than 4 inches tall and the higher rate for weeds 4 to 6 inches tall. As the last ingredient, add a suitable surfactant (refer to paraquat label).

Roundup Combinations: Tank mix Command 4 EC alone or with Sencor® Lexone® Lorox® Lorox Plus observe mixing instructions in the General Mixing Instructions portion of this label, as well as individual product label use directions. Add 1 quart of Roundup per acre for control of emerged annual weeds or 2 to 4 quarts per acre for control of emerged perennial and annual weeds.

Apply in 20 to 60 gallons of water per acre. Use the higher rate for dense stubble or vegetation.

PUMPKINS

GENERAL INFORMATION

Command 4 EC herbicide may be utilized as a pre-emergent soil application for the control of annual grass and broadleaf weeds in pumpkins. Command 4 EC must be soil incorporated in the full 12-in state of incorporation (approved 24-C) or on the full 12-inches with a tiller blade (till with a chisel plow, disk, harrow, or similar tool). Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, Wisconsin.

Note: When utilizing Command 4 EC as a soil incorporated treatment, incorporate Command to a depth of 1 inch of loose soil and place the seed below the chemical barrier when planting.

Command 4 EC can be applied to the surface of the soil with a pump-up sprayer, 1/2 to 1 pint per acre. Water up to 1/2 inch of rain or irrigation is recommended. When applying, do not spray on Command 4 EC treated or other pump-up herbicides, as they will reduce the effectiveness of Command 4 EC. If rainfall or irrigation occurs within 12 hours of application, a repeat application may be required. Do not apply more than 1/2 pint per acre in a single application. Water up to 1/2 inch of rain or irrigation is recommended for Command 4 EC which is applied alone or when tank mixed with the herbicides listed above. Do not use different spray equipment.

BEST AVAILABLE COPY

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

REPLANTING INSTRUCTIONS

If initial seeding of pumpkins fails to produce a stand, pumpkins may be replanted in fields treated with Command 4 EC alone. Do not re-treat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the pumpkin replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Guidelines on this label.

COMMAND 4 EC APPLIED ALONE RATES AND WEEDS CONTROLLED

Do not exceed a rate of 2 pints per acre. Weeds controlled (by rate) are as listed under Soybean for Northern and Southern areas.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed under Soybean Rotational Crop Guidelines otherwise crop injury may occur.

Cover crops however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for feed or seed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated pumpkin vines or feed treated vines or vine trash to livestock.

SUCCULENT PEAS

GENERAL INFORMATION

Command 4 EC herbicide may be utilized as preemergent soil applied treatment for the control of annual grass and broadleaf weeds in peas.

Command 4 EC must be soil incorporated in the following states unless an approved 24(C) is on file for counties within that state (check with your local dealer). Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, Wisconsin.

Note: When utilizing Command 4 EC as a soil incorporated treatment, incorporate Command to a depth of 1 inch or less and place the seed below the chemical barrier when planting.

Command 4 EC herbicide may be tank mixed with other herbicides registered for use in peas to broaden the weed control spectrum compared to the product's individual use. When tank mixing rates of Command 4 EC plus other significant herbicides observed by tank mix and limitations as per the tank mix label and label of Command 4 EC application precautions, rotational guidelines and replanting intervals on each product's label, when using tank mixes. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone or when tank mixed with others. Excellent pea herbicides unless use directions specifically state otherwise.

SPECIAL PRECAUTION

Off site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

REPLANTING INSTRUCTIONS

If initial seeding of pea fails to produce a stand, do not re-treat field in fields treated with Command 4 EC alone (with recommended tank mixtures). Do not re-treat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the tank mix instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Guidelines on this label.

COMMAND 4 EC HERBICIDE APPLIED ALONE

FOR PEAS ONLY All rates of 1 pt per acre or 1/2 pt

RATE AND WEEDS CONTROLLED (PT 10.5 LB A)

(for all soil types)

GRASSES

Crabgrass (Large)

Panicum (Faint)

Foxtail (Green)

 Robust

Goosegrass

Field Sandbar*

Barnyardgrass*

Seeding Johnson grass

*Partial Control

 Cyperus (Sedge)

 Lamium

 Mimulus (Mallow)

 Oxalis (Wood Sorrel)

 Pigweed (Amaranth)

 Poa (Smooth)

 Rumex (Smartweed)

 Salsify (Goosefoot)

 Spartina (Cattail)

 Suaeda (Saltgrass)

 Spergula (Spurred)

 Thlaspi (Mustard)

 Tragopogon (Hawkbit)

 Vulpia (Wild Oat)

 Xanthium (Pigweed)

 Zizaniopsis (Barnyardgrass)

 Zizaniopsis (Cattail)

 Zizaniopsis (Goosefoot)

 Zizaniopsis (Smartweed)

 Zizaniopsis (Spartina)

 Zizaniopsis (Suaeda)

 Zizaniopsis (Thlaspi)

 Zizaniopsis (Tragopogon)

 Zizaniopsis (Vulpia)

 Zizaniopsis (Xanthium)

 Zizaniopsis (Zizaniopsis)

Rotate to crops as listed under Soybean Rotational Crop Guidelines otherwise crop injury may occur.

Cover crops may be planted at any time after treatment. In some areas, the use of cover crops may reduce stand reductions less than 9 months after Command 4 EC treatment. Do not allow live stock to graze on treated fields or feed treated vines or vine trash to livestock.

FALLOW LAND

GENERAL INFORMATION

Command 4 EC is a herbicide which may be utilized as a pre-emergent surface applied treatment to control many annual grass and broadleaf weeds on fallow land. Command 4 EC may be tank mixed with other herbicides registered for fallow cropland use (e.g. atrazine).

To broaden the weed control spectrum compare to the products applied alone.

SPECIAL PRECAUTION

Off site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

Additional Precaution For Fallow Land Applications Only

Do not apply Command 4 EC within 500 feet of Emerged Winter Wheat or within 1,000 feet of the areas listed below:

To ns and Subdivisions

Commercial Vegetable Production

Commercial Fruit Production

Commercial Nurseries

Commercial Greenhouses

* except sweet corn

Supplemental Spray Drift Precautions For Fallow Land Application: Non-target spray drift of Command 4 EC herbicide should be avoided to prevent whitening of desirable vegetation. Specific recommendations and requirements for minimizing spray drift are included the Application Precautions section of this label. For Fallow Land applications only:

- The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 5 to 15 gallons per acre and is recommended for application volumes of 15 to 40 gallons per acre when spraying in the proximity of desirable plants (see list under Application Precautions).

ROTATIONAL CROP GRAZING AND FEEDING RESTRICTIONS

Do not plant winter wheat or winter rye within 100 feet of the application. Do not allow animals to graze on treated land.

APPLICATION RATES

Use Command 4 EC herbicide alone at a rate of 1 pt per acre or 32 ounces per acre in a surface applied treatment and tank mix with an emergent cereal grain.

Where Gleam is registered and was applied at 1/4 to 1/2 ounce per acre for weed control in the growing wheat crop, a rate of 1/2 pint (16 ounces) per acre of Command 4 EC is recommended; however the additional weed control benefit from Gleam may not last throughout the following spring (see Gleam label). Use the higher rate of Command 4 EC where heavy seed infestations are anticipated or for longer residual control.

In areas where it is winter wheat followed winter rye cropping system is practiced, make application after wheat harvest. For example, if the application date is July 15, the application date for Command 4 EC can be delayed until August 15 when the grain is at the soft dough stage (16 to 18 months after sowing).

Treatment of a pasture may be required when the area has been maintained under a wheat crop through weed control and incorporation of the area, sprayed with Command 4 EC when the crop is emerged and the seed is no longer present throughout the growing period. Do not graze in situations where there will be a delay.

WEEDS CONTROLLED

Grass Weeds	Broadleaf Weeds
Crabgrass (Large)	Bindweed
Panicum (Faint)	Common Purslane
Foxtail (Green)	Common Sunflower
Robust	Convolvulus
Goosegrass	Corn
Field Sandbar*	Cotton
Barnyardgrass*	Dandelion
Seeding Johnson grass	Glechoma (Ground Ivy)
*Partial Control	Groundsel
	Hemp
	Horseradish
	Jack-in-the-Pulpit
	Lamium
	Mimulus (Mallow)
	Oxalis (Wood Sorrel)
	Pigweed (Amaranth)
	Poa (Smooth)
	Polygonum (Smartweed)
	Rumex (Smartweed)
	Salsify (Goosefoot)
	Spartina (Cattail)
	Suaeda (Saltgrass)
	Spergula (Spurred)
	Thlaspi (Mustard)
	Tragopogon (Hawkbit)
	Vulpia (Wild Oat)
	Xanthium (Pigweed)
	Zizaniopsis (Barnyardgrass)
	Zizaniopsis (Cattail)
	Zizaniopsis (Goosefoot)
	Zizaniopsis (Smartweed)
	Zizaniopsis (Spartina)
	Zizaniopsis (Saltgrass)
	Zizaniopsis (Thlaspi)
	Zizaniopsis (Tragopogon)
	Zizaniopsis (Vulpia)
	Zizaniopsis (Xanthium)
	Zizaniopsis (Zizaniopsis)

COMMAND 4 EC APPLIED IN COMBINATIONS

For more information on Command 4 EC tank mix applications to broaden the weed control spectrum please contact the manufacturer or distributor of the other products used alone.

BEST AVAILABLE COPY

-12-

COMMAND plus ATRAZINE

Command 4 EC at 1 to 1 pints (7.5 to 15 fluid ounces) per acre and 1 to 2 pints of atrazine 4E or equivalent rates of atrazine 4E, 1 to 1 to 1.5 Weeds controlled include, but not limited to, Palmer amaranth, pigweed, waterhemp, and other annual weeds. Control will depend on atrazine rate used (see atrazine label) and the proper rate of Command 4 EC when ready to apply information can be required or for longer residual control.

There is significant risk of atrazine damage to crops. Check with your local Experiment Station and/or the nearest Extension Service or Research Experiment Station for recommendations on atrazine use before you plant in your area.

Refer to the atrazine label for planting, cropping and other restrictions. Follow all precautions, statements, and restrictions on the atrazine label.

Mixing Instructions.

Fill spray tank 1/2 to 2/3 full with water and add Command 4 EC and atrazine with regular operation. Mix thoroughly and then add herbicide timing addition. Add Command 4 EC to tank to time of mixing and then timing addition. Complete filling the spray tank with water. Shut off application during mixing and application.

Application Instructions

Follow the Command Acreage directed by the label for Command 4 EC.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use. On purchase of this product buyer and user agree to the following conditions:

Warranty. FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations. Follow directions carefully. Timing and method of application, weather and crop condition, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product. FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with directions or established sale practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Labels, trademarks and registered trademarks:

COMMAND and **FMC**—Trademark of FMC Corporation

CANOPY, LEXONE and PREVIEW—Trademarks of E.I. duPont de Nemours and Co., Inc.

OPTION—Trademark of Hoechst-Roussel Agri-Vet Company

SONALAN, TREFLAN—Trademarks of Elanco Products Company

LASSO—Trademark of Monsanto Company

SENCOR—Trademark of Parent Company of Farbenfabriken Bayer GmbH, Leverkusen

PROWL, SCEPTER, SOUNDROUN, TRI-SCPT—Trademarks of American Cyanamid Company

DEALER—Trademark of Ciba-Geigy Corporation

BLAZER—Trademark of BASF AG

TACKLE—Trademark of Rhone-Poulenc, Inc.

1987 FMC Corporation

All rights reserved

Product Bulletin

COMMAND® 4 EC HERBICIDE PREEMERGENCE APPLICATION SOYBEANS

Code 618
Command 4 EC Herbicide
EPA Reg. No. 279-3053



Supplemental Label

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Soybeans—Southern Area*



* Southern area includes Missouri bootheel which incorporates the following counties: Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott and Stoddard.

General Information:

Command 4 EC herbicide may be utilized as a preemergent soil applied or as a soil incorporated treatment for the control of annual grass and broadleaf weeds in soybeans. Command 4 EC herbicide may be tank mixed with other herbicides to broaden the weed control spectrum compared to the products applied alone. Observe all precautions, instructions, and rotational cropping guidelines of each products label when tank mixing.

IMPORTANT

This supplemental labeling contains directions for use of this product which do not appear on the package label. Follow instructions carefully. Read and carefully observe the precautionary statements and all other information appearing on the product label. This label must be in the possession of the user at the time of application.



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

Command® 4 EC Herbicide Preemergence Application

Southern Area

Application Precautions for Preemergent Use

When utilizing Command 4 EC as a preemergent soil applied treatment without incorporation, however, additional precautions, beyond those required for a preplant incorporated treatment, must be taken as follows:

Do not apply Command 4 EC within 1,500 feet of the following areas:

- Towns and Subdivisions**
- Commercial Fruit or Vegetable* Production**
- Commercial Greenhouses or Nurseries**

Prior to application, adjacent properties must be checked, and spraying within 1,500 feet of desirable plants must be avoided.


***except sweet corn**

Spray Drift Precautions:

Non-target spray drift of Command 4 EC herbicide should be avoided to prevent whitening of desirable vegetation. Drift is influenced by many factors which include wind speed, spray pressure, particle size, nozzle type and boom height.

Refer to package label for specific recommendations and requirements regarding:

- Precautionary Statements**
- Storage and Disposal**
- Sprayer Cleanup**
- General Mixing Instructions**
- Application Precautions**
- Special Precautions**
- Drift Control Supplement**



Command® 4 EC Herbicide Applied Alone

Southern Area

Rates and Weeds Controlled

1.5 pints*
(0.75 lb a.i.)

2.0 pints
(1.0 lb. a.i.)

2.5 pints
(1.25 lbs. a.i.)

Grasses

Barnyardgrass
Broadleaf Signalgrass
Crabgrass (Large, Smooth)
Field Sandbur*
Foxtail (Green, Giant)
Goosegrass
Panicum (Fall, Texas, Common)
Seedling Johnsongrass

Grasses

All grasses controlled at 1.5 pints
plus:
Cupgrass (Woolly, Southwestern)
Field Sandbur
Bermudagrass*
Red Rice*
Itchgrass†

Grasses

All grasses controlled at 2.0 pints
plus:
Foxtail (Purple, Robust, Yellow)
Shattercane*
Wild Proso Millet*

Broadleaves

Lambsquarters
Prickly Sida
Purslane
Redweed
Spurred Anoda
Velvetleaf
Venice Mallow
Common Ragweed*
Jimsonweed*
Pennsylvania Smartweed*

Broadleaves

All broadleaves controlled at 1.5 pints *plus:*
Cocklebur*
Common Ragweed*
Dayflower
Florida Pusley
Jimsonweed
Tropic Croton
Wild Poinsettia#
Kochia*
Redvine*

Broadleaves

All broadleaves controlled at 2.0 pints *plus:*
Kochia
Morningglory (Pitted)
Pennsylvania Smartweed
Prostrate Spurge
Black Nightshade*
Pigweed (Redroot*, Smooth*)


*Partially Controlled

* Higher labeled rates are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types.

#Use in AR/LA/MS/TX/OK only—for high pressure or heavy soils use 2.5 pints

†For Control of Itchgrass, do not incorporate Command 4 EC. For high weed pressure and heavy soils use 2.5 pints.

Note: When using Command 4 EC more than 30 days prior to planting, the highest label rate should be used.



COMMAND® 4 EC HERBICIDE PREEMERGENCE APPLICATION

Rotational Guidelines

Southern Area

(Soil applied or soil incorporated treatments)

Rotate to crops as listed below, otherwise crop injury may occur.

1.5 to 2.0 pints
(0.75 to 1.0 lb. a.i.) **2.5 pints**
(1.25 lb. a.i.)

Anytime
Soybeans
Pumpkins

9 Months
Corn (Field, Pop, Seed, Sweet)
Cotton
Cucurbits
Dry Beans
Peanuts
Peas
Peppers
Potatoes
Rice
Snap Beans
Sorghum
Sugar Beets
Sweet Potatoes
Tobacco
Tomatoes (Transplanted)

9 Months
Corn (Field, Pop, Seed, Sweet)
Cotton
Cucurbits
Dry Beans
Peanuts
Peas
Peppers
Potatoes
Pumpkins
Rice
Snap Beans
Sorghum
Sugar Beets
Sweet Potatoes
Tobacco
Tomatoes (Transplanted)

12 Months
All crops

Cover crops may be planted anytime, but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

