

Code 618

Net Contents

# Command 4 EC

## Herbicide

For Agricultural or Commercial Use Only  
NOT FOR SALE OR USE IN CALIFORNIA

EPA Reg. No. 279-3053

EPA Est. 279-

<b>Active Ingredient:</b>	<b>By Wt.</b>
2-(2-Chlorophenyl)methyl-4, 4-dimethyl-3-isoxazolidinone.....	47.1%
<b>Inert Ingredients:</b> .....	52.9%
	100.0%

Contains xylene range aromatic solvents  
Contains 4 pounds of active ingredient per gallon  
U.S. Patent No. 4,405,357

**ACCEPTED**

1-9

Under contract to the U.S. Environmental Protection Agency  
Registration No. 279-3053  
EPA Reg. No. 279-3053

### KEEP OUT OF REACH OF CHILDREN WARNING

#### FIRST AID

**If in eyes:** Flush immediately with plenty of water. Get medical attention as soon as possible.

**If swallowed:** Drink promptly large quantities of milk or water. Avoid alcohol. Do not induce vomiting. Call a physician.

**If inhaled:** Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention as soon as possible.

**If on skin:** Remove contaminated clothing. Wash skin with plenty of soap and water.

For Emergency Assistance Call Collect: (716) 735-3765

#### PRECAUTIONARY STATEMENTS

##### Hazards to Humans (and Domestic Animals)

###### Warning

Causes eye injury (corneal opacity) that is temporary. Do not get in eyes or on clothing. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing vapor or spray mist. Wear goggles or face shield when measuring, mixing or loading. Avoid contact with skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**BEST AVAILABLE COPY**



FMC Corporation  
Agricultural Chemical Group  
Philadelphia PA 19103

#### Environmental Hazards

Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water by the leaching of equipment or disposal of wastes. Apply this product only as specified on this label.

See other panels for additional precautionary information.

TABLE OF CONTENTS	
Read all information on pages 1-4 before referring to specific crop use	
Ingredients	Page 1
First Aid	1
Precautionary Statements	1
Directions for Use	1
Storage and Disposal	1
Sprayer Cleanup	2
General Mixing Instructions	2
Application Precautions	2
General Application Instructions	2
Command Geographical Areas	2
SOYBEANS	2
General Information	2
Replanting Instructions	2
Rates and Weeds Control	2
Rotational Crop Guidelines	2
Command Applied in Combination	2
Tank Mixes	2
or Followed by Preemergence or Postemergence Treatments	2
PUMPKINS	2
SUCCULENT PEAS	2
FALLOW LAND	2

#### SPECIAL PRECAUTION

Off-site movement of spray drift is a potential concern. If herbicide can cause leaf whiplash or any wind-blown spray, Prior to making application, refer to the label for drift control and apply it in accordance with the label.

#### Physical-Chemical Hazards

See other panels for additional information.

#### DIRECTIONS FOR USE

Read label and follow directions for use. Do not use in combination with other herbicides.

Do not use in combination with other herbicides.

based on  
1/4/89 - SAL

page 13

## STORAGE AND DISPOSAL

### Pesticide Storage

Do not freeze. If the contents freeze, do not thaw crystals. Do not observe warm-up in the field. After use, place containers in warm, well-ventilated area. Empty containers periodically to residue-free solid.

Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, re-secure and seal tightly. Do not put concentrate or dilute mixture in food or drink containers. Do not contaminate other pesticide containers with food or feed by storage or disposal.

In case of spill, avoid contact with bare area and keep out animals and unprotected persons. Contain spills. Call FMC Collect (716) 735-3765.

To contain spill: Dams around the area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify content.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### Container Disposal

Triple rinse for edibles only. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### Returnable Refillable Sealed Containers

Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

1. Fill the spray tank with water. Fill the spray tank a second time over the plant. Do not observe the same procedure as noted in #1 above.

2. If the contents of the spray tank will be applied to a pest emergency crop, drain the spray tank into a solution from this second rinsing. Retain this solution for use only with a solution registered pesticide to be applied on a crop for which Command 4 EC is registered. Do not dilute with the water for the next pesticide load.

### HOWEVER

If the next use of the sprayer will be a pest emergency applied pesticide on any crop, drain the solution from this second rinsing. Retain this solution for use only with a solution registered pesticide to be applied on a crop for which Command 4 EC is registered. Do not dilute (after draining second rinsate solution) in accordance with pest emergency product label directions.

**NOTE:** Dispose of excess spray mixture and/or rinsate from first rinsing by application and incorporation to cropland as described on the label. If excess spray mixture and/or rinsate from first rinsing cannot be disposed of according to label instructions, dispose of in compliance with local, state and Federal guidelines. Contact your state pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance.

## GENERAL MIXING INSTRUCTIONS

Care must be taken when mixing Command 4 EC herbicide. Avoid mixing in areas adjacent to desirable plants. (See Section headed "IMPORTANT").

**Command 4 EC Alone:** Mix Command 4 EC with water or liquid fertilizer in the following manner: Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 4 EC, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

**Tank Mixtures:** Fill spray tank one-fourth to one-third full with water with agitator operating, add the recommended amount of ingredients using the following order: dry formulations (e.g., wettable powders, dry flowables, first; liquid suspensions (e.g., flowables) next and finally liquids (e.g., ECs). Mix thoroughly and fill tank one-half full, continue agitation. Add Command 4 EC herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Where use of a surfactant is recommended, add as the last ingredient to the spray tank. Maintain agitation during filling, mixing and application. When using drift reducing agents, follow specific product label instructions for order of addition to spray tank.

**Fertilizer Spray Mixtures:** Applications of Command 4 EC alone or with recommended tank mixtures, in conjunction with fertilizer solutions may be used unless use directions specifically state otherwise. Spray quantities should be limited for compatibility by the following procedure:

1. Fill the spray tank with water to 1/4 full.
2. Add the appropriate amount of herbicide based on the label rate. If more than one product is to be used, add each separately, using the following sequence: dry formulations (e.g., wettable powders, dry flowables, first; liquid suspensions (e.g., flowables) next and finally liquids (e.g., ECs).

Herbicide	Rate/Acre	Amount To Be Added Per Pint of Fluid Fertilizer*
Wettable Powders	1.0 pint	1 level teaspoon
Flowables	1.0 pint	1 level teaspoon
EC's	2.0 pints	3 level teaspoons
Dry Flowables (Dry Granules)	3.0 pints	4 level teaspoons
Emulsifiable Concentrate	1.0 pint	1 level teaspoon
Flowables	1.0 pint	1 level teaspoon
Flowables	1.0 pint	2 level teaspoons
Flowables	1.0 pint	3 level teaspoons

\*Based on the spray volume of 100 gallons per acre. Do not exceed 100 gallons per acre.

3. Close tank and shake well.
4. Water mixture for use in 30 minutes and apply after 30 minutes. If herbicide is registered for use as a pre-emergent, it can be removed to the soil surface 10 days before application. Do not be concerned if the herbicide is applied 30 days before application to weeds and grasses.

Pre-emergent herbicides should be applied to soil before the weeds and grasses emerge.

1. **Command 4 EC Alone:** Fill with a part of water and add 1/4 of spray tank of fertilizer with agitator running. Maintain agitation and apply.

## IMPORTANT

FAILURE TO OBSERVE THE APPLICATION PRECAUTIONS SECTION OF THIS LABEL MAY RESULT IN INJURY TO DESIRABLE VEGETATION.

- Desirable plants including some species of trees, shrubs, flowers, agronomic crops, and fruits and vegetables are sensitive to Command 4 EC herbicide.
- Foliar contact with spray drift or vapors may cause whitening of sensitive plants. Such symptoms are generally temporary in nature, but may persist on some plants.

## SPRAYER CLEANUP

Do not drain or flush equipment in or near desirable trees or other plants of the area, where the herbicide may, either in solution where the chemical may be washed or moved into contact with their roots. Do not contaminate any body of water or underground water that may be used on other crops. Carry out a wash sprayer clean up instructions noted below to prevent spray tank residue from damaging other crops.

Sprayer equipment should be thoroughly rinsed with water. Residue of herbicide that may be present on other spray nozzles. The spray nozzle should be tested for the herbicide residue. Do not use equipment following applications of Command 4 EC herbicide with the mixed of Command 4 EC with other labeled products.

1. Drain any remaining spray solution from tank, pump, hoses, and boom, and discard in an approved manner. (See Note that follows.)
2. Clean tank and hoses.
  - Thoroughly flush with water. Fill the spray tank with a quantity of water equal to 1/2 the tank capacity and operate the pump or agitator for 15 minutes through the sprayer system for 15 minutes.
  - Wash nozzles with water. Clean all equipment.
  - Remove the nozzle and hose from the tank and boom sections and wash the inner surfaces of rinsate solution through completely through the nozzle and hose.
3. Thoroughly drain remaining spray solution from tank, pump, and hoses. Combine with the following and dispose of a separate time this first rinsing as an application of water only.
  - When washing from within the tank, application of a spray nozzle should be done with the tank pump, hoses, and boom present. The nozzle should be tested for the herbicide residue. Do not use equipment following applications of Command 4 EC herbicide with the mixed of Command 4 EC with other labeled products.
4. Remove the remaining residue from the nozzle and the boom, and wash it up as follows:
  - Wash with water.





agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre. Use nozzles suitable for broadcast application of herbicide. Do not use a spray nozzle that is not suitable for the target area. Do not use a spray nozzle that is not suitable for the target area. Do not use a spray nozzle that is not suitable for the target area. Do not use a spray nozzle that is not suitable for the target area.

Band Application: Use the rates and volume recommendations and the following formula:

Band width inches	Broadcast Rate Per Acre	Band rate per acre
Band width inches	Broadcast Volume Per Acre	Band volume per acre

Follow broadcast minimum gallonage requirements adjusted to band width. Do not exceed 40 psi spray pressure.

Refer to Spray Drift Precautions for suggested pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types. For banded applications an even flat-fan nozzle is recommended.

**Soil Incorporated Application:** Apply Command 4 EC to the soil surface and uniformly incorporate. Select equipment suited for proper incorporation. **Shallow incorporation normally results in better weed control than a deeper incorporation.**

In states where soil incorporation is required (see Geographical Restrictions) application and immediate incorporation to a maximum depth of 3 inches is required unless the soil is dry. On dry soils, incorporation to a maximum depth of 3 inches must be completed within 8 hours of Command 4 EC herbicide application.

Soil must be in good tilth to allow for thorough mixing of the soil. Application to wet soils will increase the potential for off-site movement of Command 4 EC herbicide vapors and may result in poor soil incorporation and unsatisfactory weed control.

## INCORPORATION EQUIPMENT

### ONE PASS INCORPORATION

#### Suggested Implements

For one pass incorporation use C or S shank field cultivators equipped with a leveling device such as a spike or coil line harrow or rolling basket attachment. Field cultivator must have 3 or 4 rows of sweeps spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Equipment design, including any drag attachments, must be designed to allow soil depth which may result in streaked or reduced weed control. The implement should be operated at 4 to 6 mph and set to cut 3 to 4 inches deep at the maximum herbicide rate. Do not use.

Combination equipment (three or more tillage devices combined and used as a single tool) should be operated at a depth of 3 to 4 inches and a minimum speed of 6 mph.

TO drive seed bed preparation equipment should be set to cut 2 inches deep and operated at 4 mph or less.

### TWO PASS INCORPORATION

#### Suggested Implements

When using two pass incorporation, the second pass should be at an angle to the first pass.

Disc set to cut 3 to 4 inches deep at 4 to 6 mph. Blades should be less than 20 inches in diameter and spaced 9 inches or less apart.

Rolling cultivator set to cut 2 to 3 inches deep at 6 to 8 mph (use on coarse or medium textured soils only).

Field cultivator with 3 to 4 rows of sweeps spaced at 7 inches or less and set to cut 3 inches deep at 4 mph or greater.

#### Cultivation After Planting

Soil treated with Command 4 EC may be shallow cultivated, rotary hoed or handhoed without reducing the weed control activity of Command 4 EC. Do not cultivate deeper than the Command 4 EC treated layer of soil since this may bring untreated soil to the surface and poor weed control may result.

## COMMERCIAL IMPREGNATION AND APPLICATION OF COMMAND 4 EC ON DRY BULK FERTILIZERS

Command 4 EC may be impregnated on dry bulk fertilizers. When applied as a dry bulk fertilizer, Command 4 EC must be applied to wet soil. The application rate provided by the same label of Command 4 EC must be used.

The Command 4 EC fertilizer mixture must be so labeled on a state or federal registration label. For further information, contact the manufacturer of the fertilizer.

**Impregnation:** Apply using a minimum of 200 pounds of dry bulk fertilizer per acre and up to a maximum of 400 pounds per acre with the recommended amount of Command 4 EC herbicide per acre. Use a coated rotary drum mixer or a similar type of coated blender equipped with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform fine spray pattern over the tumbling fertilizer for thorough coverage. The physical properties of fertilizers vary particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Command provides a satisfactory dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry flowable mixture. Microcel E (Johns-Manville Products Corporation) is a recommended absorbent powder. Generally less than 2% by weight of Microcel E is required. **DO NOT impregnate Command 4 EC onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the herbicide.** Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with Command.

The amount of Command actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with Command 4 EC herbicide should be applied immediately, not stored. Care should be taken to ensure spreading procedures in the field do not place fertilizer treated with Command in areas which cannot be incorporated, if incorporation is required within the Geographical Restrictions of this label.

All state regulations, labeling, etc., of the mixtures are the responsibility of the seller.

For those rates not listed in the following table, calculate the amount of Command to be impregnated on a ton of dry bulk fertilizer using the following formula:

2000 pounds dry fertilizer per acre	PINTS of Command per acre (recommended rate for soil texture)	PINTS of Command per ton of fertilizer
--	--	--

## RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZERS WITH COMMAND 4 EC HERBICIDE

### PINTS OF COMMAND 4 EC PER TON OF FERTILIZER

Fertilizer Rate Lbs. Acre	Command 4 EC Rate Per Acre		
	1 1/2 pts (75 lbs A/E)	2 pt (1 lb A/E)	2 1/2 pts (125 lbs A/E)
200	15	20	25
250	12	16	20
300	10	13	16
350	8	11	14
400	7	10	12
450	6	9	11


If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation. As dusty fertilizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

## SPECIAL PRECAUTIONS FOR DRY BULK FERTILIZER IMPREGNATION AND APPLICATION WITH COMMAND 4 EC

- All equipment used to apply Command should be thoroughly cleaned immediately following use to ensure no contamination results which could cause injury to non-target crops or desirable vegetation. Refer to "Sprayer Clean-up" section for additional details and disposal instructions.
- Equipment used to physically transport Command treated fertilizer, including boots, aprons, conveyors, bins, etc., should be covered to prevent loss of fine particles and subjected to clean-up procedures previously described.
- Mixing of Command herbicide and dry fertilizer should be conducted in a blender which will ensure that airborne particles and potential volatilization are avoided.
- Bulk containers should be tightly covered when being stored, transported and applied to reduce chances of Command volatilization or product loss.
- It is recommended that Dry Bulk Fertilizer impregnation operations using Command not be conducted within 1/4 mile of residential areas or areas where Command symbolology or desirable vegetation will not be tolerated.

**BEST AVAILABLE**

**COMMAND GEOGRAPHICAL AREAS**  
**C. Rates, Weed Control, and Crop Rotation**



Southern area includes Missouri, together with incorporates the following counties: Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott, and Stoddard.

**SOYBEANS**

**GENERAL INFORMATION**

Command 4 EC herbicide may be utilized as a preemergent soil applied or soil incorporated treatment for the control of annual grass and broadleaf weeds in soybeans.

Command 4 EC must be soil incorporated in the following states unless approved 24(C) is on file for counties within that state (check with your local dealer): Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, Wisconsin.

Command 4 EC herbicide may be tank mixed with other herbicides to broaden the weed control spectrum compared to the products applied alone. Command 4 EC may be tank mixed with Canopy, Duan, Lasso, Lurex, Lurex Plus, Pexone, Preview, Prowl, Scepter, Sencor, Sonalan, Squadron, Trellan, Tri-Scept, or other registered herbicides. Observe all precautions, instructions,

and rotational cropping guidelines of each product's label when tank mixing. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone or when tank mixed with the herbicides listed above, unless use directions specifically state otherwise.

**REPLANTING INSTRUCTIONS**

If initial seeding of soybeans fails to produce a stand, soybeans may be replanted in fields treated with Command 4 EC alone (or with recommended tank mixtures). Do not retreat fields with a second application of Command 4 EC. When tank mixing with a labeled product, refer to the soybean replant instructions for that product. Do not replant fields with any crop at intervals which are inconsistent with the Rotational Guidelines on this label.

**Northern Area—See Map**

**COMMAND 4 EC HERBICIDE APPLIED ALONE**

**RATES AND WEEDS CONTROLLED†**

1.0 PT. (0.50 LB. A.I.)	1.5 PT. (0.75 LB. A.I.)	2.0 PT. (1.0 LB. A.I.)
<b>GRASSES</b>	<b>GRASSES</b>	<b>GRASSES</b>
Compression fescue crab grasses	Barnyardgrass Crabgrass Large crabgrass Field bindweed Fingerweed Fount grass Rabbit grass Yellow foxtail Panicum grass Poa grass Ryegrass Spike grass	All grasses controlled at 1.5 pints plus Panicum grass Barnyardgrass Field bindweed Fingerweed Fount grass Rabbit grass Yellow foxtail Panicum grass Poa grass Ryegrass Spike grass Wild Horse Mallow Black Seeded Plantain

**BROADLEAVES**

Velvetleaf  
Spurred Anoda

**BROADLEAVES**

Velvetleaf  
Spurred Anoda  
plus  
Galinsoga  
Lambsquarter  
Prickly Sida  
Purslane  
Venice Mallow  
Common Ragweed\*  
Jimsonweed\*  
Smartweed  
- Pennsylvania\*  
- Ladysthumb\*

**BROADLEAVES**

All broadleaves  
controlled at 1.5  
pints plus  
Common Ragweed  
Jimsonweed  
Kochia  
Tropic Croton  
Cocklebur\*  
Smartweed  
- Pennsylvania  
- Ladysthumb

\* Part 1, Controlled

† Higher label rates (a minimum of 2 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types.

^ For heavy weed pressure or heavy soils use 1.25 pints per acre to control Velvetleaf or Spurred Anoda.

For Soil Applied Preemergence Applications Only: Command 4 EC may be applied at a rate of 2.1-2 pints (1.25 pounds A.I.) in states where soil incorporation is not required (refer to Application Precautions Section of this label). This higher rate will improve the level of weed control achieved, and in addition will control the following species not listed at the lower rates above:

2.5 PINTS (1.25 LB. A.I.)  
APPLIED PREEMERGENCE

Weeds Controlled (in addition to those listed above)

Grasses	Broadleaves
Shattercane Witchgrass*	Fitted Morningglory Prostrate Spurge Black Nightshade* Ligweed Bedstraw Sowthistle
Quackgrass*	Horseweed Wild Radish Beggarticks Marestail Swamp Smartweed Common Milkweed* Giant Ragweed*

*Resistant to 4 EC*

NOTE: When using Command 4 EC more than 30 days prior to planting, the highest label rate should be used.

**ROTATIONAL CROP GUIDELINES**

Rotate to crops as listed below, otherwise crop injury may occur.

1.0 PT. (0.50 LB. A.I.)	1.5 PT. (0.75 LB. A.I.)	2.0-2.5 Pints (1.0-1.25 LB. A.I.)
<b>ANYTIME</b> Soybeans Pumpkins Peas	<b>ANYTIME</b> Soybeans Pumpkins	<b>ANYTIME</b> Soybeans Pumpkins
<b>9 MONTHS</b> Corn -- Field -- Pop -- Sweet -- Sweet	<b>9 MONTHS</b> Same as 1.0 pint, plus Peas	<b>9 MONTHS</b> Same as 1.5 pints except Seed Corn and Sweet Corn require a 12 month rotation
Cotton Cucurbits Dry Beans Peanuts Peppers Potatoes Rice Snap Beans Sorghum Sugar Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)		
<b>12 MONTHS</b> Cabbage Tomatoes (all) Wheat	<b>12 MONTHS</b> Same as 1.0 pint	<b>12 MONTHS</b> Same as 1.5 pints plus Seed Corn Sweet Corn
<b>16 MONTHS</b> All crops	<b>16 MONTHS</b> All crops	<b>16 MONTHS</b> All crops

**BEST AVAILABLE COPY**

Cover crops, however, may be planted anytime but stand reduction may occur in some areas. Do not graze or harvest for feed or bedding crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated pasture areas or feed treated vines or vine trash to livestock.

**Southern Area—See Map**  
**COMMAND 4 EC HERBICIDE**  
**APPLIED ALONE**

**RATES AND WEEDS CONTROLLED\***

1.0 PT. (0.50 LB. A.I.)	1.5 PT. (0.75 LB. A.I.)	2.0 PT. (1.0 LB. A.I.)	2.5 PT. (1.25 LB. A.I.)
<b>GRASSES</b> Suppression of some annual grasses	<b>GRASSES</b> Barnyardgrass Broadleaf signalgrass Crabgrass Large Smooth Foxtail Giant -Green Goosegrass Panicum Common Labi Texas Seeding Johnsongrass Field Bindweed	<b>GRASSES</b> All grasses controlled at 1.5 pints plus Cocklebur Soybean -Woody Field bindweed Bermudagrass Red Rice Itchgrass***	<b>GRASSES</b> All grasses controlled at 2.0 pints plus Field bindweed Bermudagrass Red Rice Itchgrass***
<b>BROADLEAVES</b> Velvetleaf Spurred Anoda	<b>BROADLEAVES</b> Velvetleaf Spurred Anoda Plus Lambsquarters Prickly Sida Furflanc Redweed Venice Mallow Common Ragweed* Jimsonweed* Pennsylvania Smartweed*	<b>BROADLEAVES</b> All broadleaves controlled at 1.5 pints plus Cocklebur Common Ragweed Dayflower Florida Pusley Jimsonweed Tropic Croton Wild Poinsettia** Kochia* Redvine* Florida Beggarweed	<b>BROADLEAVES</b> All broadleaves controlled at 2.0 pints plus Kochia Morningglory -Pitted Pennsylvania Smartweed Prostrate Spurge Black Nightshade* Pigweed -Redroot* -Smooth* Balloonvine* Jungle Rice* Joint Vetch* Curly Dock

\* Partially Controlled

Higher label rate (a minimum of 2 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, a minimum application rate of 2 1/2 pints per acre for cocklebur or common ragweed control is recommended.

\*\* Use in AR LAMS T. OK only for high pressure and heavy soils use 2.5 pints

For heavy weed pressure or heavy soils use 1.25 pints per acre to control Velvetleaf or Spurred Anoda.

\*\*\* For control of Itchgrass, do not incorporate Command 4 EC. For high weed pressure and heavy soils use 2.5 pints

NOTE: When using Command 4 EC more than 30 days prior to planting, the highest label rate should be used

**ROTATIONAL CROP GUIDELINES**

Rotate to crops as listed below; otherwise crop injury may occur

1.0 PT. (0.50 LB. A.I.)	1.5 to 2.0 PT. (0.75 to 1.0 LB. A.I.)	2.5 PT. (1.25 LB. A.I.)
<b>ANYTIME</b> Soybeans Pumpkins Peas	<b>ANYTIME</b> Soybeans Pumpkins	<b>ANYTIME</b> Soybeans
<b>9 MONTHS</b> Corn -Field -Pop -Seed -Sweet Cotton Cucurbits Dry Beans Peanuts Potatoes Pumpkins Rice Snap Beans Sorghum Sugar Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)	<b>9 MONTHS</b> Same as 1.0 pint plus Peas	<b>9 MONTHS</b> Same as 1.5 to 2.0 pints plus Pumpkins
<b>12 MONTHS</b> All crops	<b>12 MONTHS</b> All crops	<b>12 MONTHS</b> All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock

**BEST AVAILABLE COPY**

**COMMAND 4 EC HERBICIDE**  
**APPLIED IN COMBINATIONS FOR SOYBEANS**

Command 4 EC herbicide may be tank mixed with other herbicides for control of weeds. For the weed control spectrum, compare the product application rate to the label. Use the following FMC recommendations and number of products and application rates to broaden weed control spectrum. FMC makes no claims as to weed control for any combination of rates or products other than as recommended in this label. Additional recommendations for rate combinations with other herbicides, not included in the following sections may become available subsequent to the date of publication for this label. Please check with your dealer for more recent FMC technical bulletins and label supplements. When applying mixtures of Command 4 EC with other herbicides, observe all precautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixes.

**Command 4 EC Herbicide plus Canopy Herbicide—**  
**Northern & Southern Areas**

Tank mix combination for broadleaf weed control spectrum

Soil Texture	Broadcast Rate Per Acre*		
	Command North	Command South	Canopy
(1% to 5% Organic Matter)			
Coarse (light) Soils (loamy sand, sandy loam)	1 to 2 pints	2 to 2 1/2 pints	4 to 6 oz
Medium Soils (loam, silty loam, sandy clay, sandy clay loam)	1 to 2 pints	2 to 2 1/2 pints	5 to 8 oz
Fine (heavy) Soils (clay, clay loam, silty clay loam)	1 to 1 1/2 pints	2 to 2 1/2 pints	6 to 8 oz

\* Use a lower to higher rate with the higher to lower rate to the coarse to fine soil types with the broadcast application.

\*\* Tank with the label rate of both products applied with a previous application of a herbicide other than Command 4 EC.

**WEEDS CONTROLLED**

Grass Weeds	Broadleaf Weeds
All species listed under Command Applied Alone Weeds Controlled section	All species listed under Command Applied Alone Weeds Controlled section plus the following: Cocklebur** Pigweed (Redroot, Smooth, King, Setaforma) Morningglory* (annual, biennial, perennial, pitted, tall) Sicklepod* Copperleaf hophornbeam*

\* Partially Controlled

\*\* Large seeded weeds such as hairy vetch, soybean, sunflower, and other species may require a cultural or mechanical control method in addition to herbicide control.

When applying mixtures of Command 4 EC plus Canopy, observe all precautions and limitations appearing on the label of each product. Special attention must be paid to sections regarding planting depth, phosphate, pesticide interaction, previous pesticides, and sensitive soybean varieties on the Canopy label. Observe application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Dual 8 E Herbicide—**  
**Northern & Southern Areas**

Tank mix combination for broadleaf weed control spectrum

Soil Texture	Broadcast Rate Per Acre*		
	Command North	Command South	Dual 8 E
Coarse (light) Soils (loamy sand, sandy loam)	1 to 2 pints	2 to 2 1/2 pints	1 to 2 pints
Medium Soils (loam, silty loam, sandy clay, sandy clay loam)	1 to 2 pints	2 to 2 1/2 pints	2 to 2 1/2 pints
Fine (heavy) Soils (clay, clay loam, silty clay loam)	1 to 1 1/2 pints	2 to 2 1/2 pints	2 to 2 1/2 pints

**WEEDS CONTROLLED**

**Grass Weeds\***

All species listed under Command Applied Alone Weeds Controlled section plus the following

- Red Rice\*
- Witchgrass
- Nutsedge

\* Use higher rate in areas with heavy populations

When applying mixtures of Command 4 EC plus Lasso, observe all cautions and limitations appearing on the labels of each product. Observe all Application Precautions, Rotational Crop Guidelines, and Replanting Instructions of each product label when using tank mixtures.

**Command 4 EC Herbicide plus Lasso\* or Lasso\* Micro-Tech—Northern & Southern Areas**

A tank-mix combination of Command 4 EC herbicide and Lasso or Lasso Micro-Tech herbicide will broaden the weed control spectrum compared to either ~~herbicide~~ used alone

**Product Broadcast Rates Per Acre**

Soil Texture	Command		
	North	South	Lasso/Lasso Micro-Tech
Coarse (light) Soils (loamy, silty, sandy loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	2 to 2 1/2 quarts
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	2 to 3 quarts
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1 1/2 to 2 pints	2 to 2 1/2 pints	2 to 3 quarts

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, loamy sand, sandy loam) within a textural group.

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**

All species listed under Command Applied Alone Weeds Controlled section plus the following

- Carpetweed
- Pigweed
- Black Nightshade

When applying mixtures of Command 4 EC herbicide plus Lasso or Lasso Micro-Tech, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product label when using tank mixtures.

**COMMAND 4 EC HERBICIDE PLUS LOROX\***

A preemergence tank mix combination of Command 4 EC herbicide and Lorox herbicide will broaden the weed control spectrum compared to either product alone.

The tank mix combination may not be incorporated. Where Command 4 EC must be incorporated (see Geographical Restrictions), apply Lorox as an overlay treatment.

**Product Broadcast Rates Per Acre\***

Soil Texture	Command 4 EC		Lorox DF†
	North	South	
Coarse (light) Soils (loamy, silty, sandy loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	4 1/2 to 5 pints
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	1 1/2 to 2 pints
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1 1/2 to 2 pints	2 to 2 1/2 pints	1 1/2 to 2 pints

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

† Plant seed at least 1 inch deep.

Deep soil incorporation in soybeans may result. Refer to Lorox label for other planting and application precautions. one pint of Lorox DF equals one pound of Lorox DF. Use equivalent rates.

Do not use on sand, gravelly soil, or exposed subsoil or soils with less than 1% organic matter. Use only on sandy sand with over 1% organic matter.

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**

All species listed under Command Applied Alone Weeds Controlled section plus the following

- Carpetweed
- Chickweed

**Pigweed**

- (Redroot pigweed)
- (Smooth pigweed)
- Wild mustard
- Wild radish

When applying mixtures of Command plus Lorox, observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop Grazing and Feeding Restrictions and Replanting Instructions sections of this label when using tank mixtures.

**COMMAND 4 EC HERBICIDE PLUS LOROX PLUS**

A preemergence tank mix combination of Command 4 EC herbicide and Lorox herbicide will broaden the weed control spectrum compared to either product alone.

The tank mix combination may not be incorporated. Where Command 4 EC must be incorporated (see Geographical Restrictions), apply Lorox as an overlay treatment.

**Product Broadcast Rates Per Acre\***

Soil Texture	Command 4 EC		Lorox Plus
	North	South	
Coarse (light) Soils (loamy, silty, sandy loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	9 to 12 pints
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	10 to 12 pints
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1 1/2 to 2 pints	2 to 2 1/2 pints	11 to 14 pints

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

† Plant seed at least 1 inch deep. Deep soil incorporation in soybeans may result. Refer to Lorox label for other planting and application precautions.

Do not use on sand, gravelly soil, or exposed subsoil or soils with less than 1% organic matter. Use only on sandy sand with over 1% organic matter.

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**

All species listed under Command Applied Alone Weeds Controlled section plus the following

- Carpetweed
- Conklebore
- Mustards
- Pigweed (redroot, smooth)
- Black Nightshade
- Hoary Mallow (annual)
- Spotted Tail
- Grass Pigweeds

\* Partially Controlling

† Plant seed at least 1 inch deep. Deep soil incorporation in soybeans may result. Refer to Lorox label for other planting and application precautions. one pint of Lorox Plus equals one pound of Lorox Plus. Use equivalent rates.

When applying mixtures of Command plus Lorox, observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop Grazing and Feeding Restrictions and Replanting Instructions sections of this label when using tank mixtures.

**BEST AVAILABLE COPY**

**Command 4 EC herbicide plus Metribuzin (Sencor\* or Lexone\*)—Northern Area Only**

A tank mix combination of Command 4 EC herbicide and metribuzin will broaden the weed control spectrum compared to either ~~herbicide~~ used alone

Soil Texture**	Broadcast Rates Per Acre*	
	Command	Sencor DF* Lexone DF
Coarse (light) Soils (loamy sand**, sandy loam)	1½ to 2 pints	¼ to ½ lb
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints*	¼ to ½ lb
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	¼ to ½ lb

- \* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- \*\* For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5
- Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter, on sand or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur.

**NOTE:** Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

**WEEDS CONTROLLED**

**Grass Weeds**  
All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**  
All species listed under Command Applied Alone Weeds Controlled section, plus the following:  
Carpetweed  
Pigweed\* (Redroot, Smooth)  
Wild mustard

\* Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**WEEDS CONTROLLED**

**Grass Weeds**  
All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**  
All species listed under Command Applied Alone Weeds Controlled section, plus the following:  
Carpetweed  
Hemp sesbania, coffee-weed  
Pigweed (Redroot, Smooth)  
Wild mustard  
Sicklepod\*

- \* Reduced Competition
- Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Preview\*—Northern Area Only**

A tank mix combination of Command 4 EC herbicide and Preview herbicide will broaden the weed control spectrum compared to either ~~herbicide~~ used alone

Soil Texture	Broadcast Rates Per Acre	
	Command	Preview*
Coarse (light) Soils (loamy sand, sandy loam)	1½ to 2 pints	6 to 10 oz
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	6 to 7 oz
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	6 to 10 oz

- \* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

Do not use Command 4 EC in combination with Preview on soils with less than 0.5% organic matter, or on soils with pH greater than 7.5 or on soils having a calcareous surface area. Preview injury to soybeans may occur.

**WEEDS CONTROLLED**

**Grass Weeds**  
All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**  
All species listed under Command Applied Alone Weeds Controlled section, plus the following:  
Carpetweed  
Spurge, spotted  
Copperleaf, hophornbeam  
Mustards  
Pigweed  
Cocklebur  
Sunflower\*  
Burdock  
Giant Ragweed  
Morningglory\* (Annual, Tall)  
Ivyleaf Pitted Entireleaf\*

**NOTE:** Control of morningglory, sunflower, cocklebur and giant ragweed may be erratic, ranging from poor to excellent depending upon soil temperature, time of weed seed germination, depth of weed seed in the soil, and the amount and timing of soil moisture. Control may be improved with timely cultivation.

When applying mixtures of Command 4 EC plus Preview, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC herbicide plus Metribuzin (Sencor\* or Lexone\*)—Southern Area Only**

A tank mix combination of Command 4 EC herbicide and metribuzin will broaden the weed control spectrum compared to either ~~herbicide~~ used alone

Soil Texture**	Broadcast Rates Per Acre	
	Command	Sencor DF* Lexone DF
Coarse (light) Soils (loamy sand**, sandy loam)	2 to 2½ pints	¼ to ½ lb
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	2 to 2½ pints	¼ to ½ lb
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	2 to 2½ pints	¼ to ½ lb

- \* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- \*\* For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5
- Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter, on sand or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur. Use 2 to 1½ pints metribuzin on silty clay and heavy clay soils of the Mississippi Delta.

**NOTE:** Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

**BEST AVAILABLE COPY**



**Command 4 EC Herbicide plus Prowl® 4 EC Herbicide—  
Northern & Southern Areas**

A tank-mix combination of Command 4 EC and Prowl will broaden the weed control spectrum of either product used alone.

Broadcast Rates Per Acre

Soil Texture	Command		Prowl 4 EC*
	North	South	
Coarse (light) Soils (loamy sand, sandy loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	1 to 1 1/2 pints
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	2 pints
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1 1/2 to 2 pints	2 to 2 1/2 pints	2 to 2 1/2 pints

\* Select lower to higher rates within the range noted based on lighter to heavier soil types (e.g. loam, silt loam, sandy clay, sandy, clay loam) within a textural group.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section.

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:  
Pigweed (Redroot Smooth)  
Carpetweed

When applying mixtures of Command 4 EC plus Prowl observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Scepter® Herbicide—  
Northern Area Only**

See Scepter label for appropriate geographic restrictions.

A tank-mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

**Note:** The Scepter label does not allow rotation to Corn, Wheat, Barley, or Oats prior to 18 months after last Scepter application when Scepter is applied pre-plant or pre-plant and initiated within the Northern Area as defined on the Scepter label. See Scepter label for appropriate geographic restrictions.

Soil Texture	Broadcast Rate Per Acre	
	Command	Scepter†
Coarse (light) Soils (loamy sand, sandy loam)	1 to 1 1/2 pints	1 to 1 1/2 pints

\* Use higher rates when heavy weed pressure is anticipated.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section.

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:  
Pigweed (Redroot Smooth)  
Cocklebur  
Pigweed spp.  
Grass leaved Galium  
Grass leaved Galium

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than two inches, or make additional Command 4 EC or Command 4 Scepter tank mix application.

**NOTE:** Adequate soil moisture is important for optimum tank mix application.

When applying mixtures of Command 4 EC plus Scepter observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Scepter Herbicide—  
Southern Area Only**

See Scepter label for appropriate geographic restrictions.

A tank-mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Scepter†
All Soil Types*	2 to 2 1/2 pints	1 1/2 to 2 pints

\* Use higher rates when heavy weed pressure is anticipated.

**WEEDS CONTROLLED**

Grass Weeds

All species listed under Command Applied Alone Weeds Controlled section.

Broadleaf Weeds

All species listed under Command Applied Alone Weeds Controlled section plus the following:  
Cocklebur  
Pigweed spp.  
Wild poinsettia

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than two inches, or make additional Command 4 EC or Command 4 Scepter tank mix application.

**NOTE:** Adequate soil moisture is important for optimum tank mix application.

When applying mixtures of Command 4 EC plus Scepter observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Sonalan® EC Herbicide—  
Northern Area Only**

A tank-mix combination of Command 4 EC and Sonalan will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rate Per Acre	
	Command	Sonalan E
Coarse (light) Soils (loamy sand, sandy loam)	1 to 1 1/2 pints	1 to 1 1/2 pints
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1 to 1 1/2 pints	1 1/2 to 2 pints
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1 to 1 1/2 pints	2 to 3 pints

\* Select lower to higher rates within the range noted based on lighter to heavier soil types within the textural group.

† Rates within the range plus or minus 10% are applied weed pressure.

**BEST AVAILABLE COPY**

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**

All species listed under Command Applied Alone Weeds Controlled section plus the following:  
 Carpetweed  
 Pigweed (Redroot Smooth)

When applying mixtures of Command 4 EC plus Sontalan, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines, and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Squadron—Northern Area Only**

See Squadron label for appropriate geographic restrictions. A tank-mix combination of Command 4 EC and Squadron will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Squadron*
All Soil Types	1 1/2 to 2 pints	1 1/2 to 2 1/2 pints

\* Do not exceed 1 1/2 pints in Northern Area as defined on the Squadron label. Use higher rate of Squadron in areas of heavy rock or thin topsoil.

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**

All species listed under Command Applied Alone Weeds Controlled section plus the following: *Black Nightshade*, *Cockspur*, *Pigweed spp. Common*, *flower*, *Grant Ragweed*\*

\* Partially controlled

When applying mixtures of Command 4 EC plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines, and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Squadron—Southern Area Only**

See Squadron label for appropriate geographic restrictions. A tank mix combination of Command 4 EC and Squadron will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Squadron*
All Soil Types	2 to 2 1/2 pints	2 to 3 pints

\* Use higher rate of product in areas of heavy rock or thin topsoil.

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**

All species listed under Command Applied Alone Weeds Controlled section plus the following: *Black Nightshade*, *Cockspur*, *Pigweed spp. Common*, *flower*, *Grant Ragweed*\*

When applying mixtures of Command 4 EC plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines, and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Trellan® EC (Trifluralin)—Northern & Southern Areas**

A tank mix combination of Command 4 EC herbicide and Trellan EC herbicide will broaden the weed control spectrum compared to either product used alone.

Soil Texture	Broadcast Rates Per Acre		
	Command	Trellan EC*	Command + Trellan EC†
Coarse (light) Soils (loamy sand, sandy loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	1 pint
Medium Soils (loam, silty silt loam, sandy clay, sandy clay loam)	1 1/2 to 2 pints	2 to 2 1/2 pints	1 to 1 1/2 pints
Fine (heavy) Soils (silt, clay, clay loam, silty clay loam, clay)	1 1/2 to 2 pints	2 to 2 1/2 pints	1 to 1 1/2 pints

\* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g. loam, silt loam, sandy clay, sandy clay loam) within a texture group.

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**

All species listed under Command Applied Alone Weeds Controlled section plus the following:  
 Carpetweed  
 Chickweed  
 Field Birdweed  
 Pigweed (Redroot Smooth)

When applying mixtures of Command 4 EC plus Trellan, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines, and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Tri-Scept Herbicide—Northern Area Only**

See Tri-Scept label for appropriate geographic restrictions. A tank-mix combination of Command 4 EC and Tri-Scept herbicide will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Tri-Scept
All Soil Types	1 1/2 to 2 pints	1 1/2 to 2 pints

\* Do not exceed 1 1/2 pints in Northern Area as defined on the Tri-Scept label. Use higher rate of Tri-Scept in areas of heavy rock or thin topsoil.

**WEEDS CONTROLLED**

**Grass Weeds**

All species listed under Command Applied Alone Weeds Controlled section

**Broadleaf Weeds**


All species listed under Command Applied Alone Weeds Controlled section plus the following:  
 Carpetweed  
 Chickweed  
 Field Birdweed  
 Pigweed (Redroot Smooth)

\* Partially controlled

When applying mixtures of Command 4 EC plus Tri-Scept observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines, and replanting instructions of each product's label when using tank mixtures.

**Command 4 EC Herbicide plus Tri-Scept—Southern Area Only**

See Tri-Scept label for appropriate geographic restrictions. A tank mix combination of Command 4 EC and Tri-Scept herbicide will broaden the weed control spectrum of either product used alone.



Use Rate	Broadcast EC	Per Acre
	Container	Tri-Scept®
Annual Weeds	2.1 to 2.5 pints	1 to 2 pints

For more information, see the separate label for each product.

**WEEDS CONTROLLED**

- |   |  |
|---|--|
| <p><b>Grass Weeds:</b><br/>All species listed under Command Applied Alone Weeds Controlled section.</p> | <p><b>Broadleaf Weeds:</b><br/>All species listed under Command Applied Alone Weeds Controlled section plus the following:<br/>Cocklebur <i>Black nightshade</i><br/>Pigweed spp. <i>Common Sulfur</i><br/>Wild poinsettia</p> |
|---|--|

When applying mixtures of Command 4 EC plus Tri-Scept observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY PREEMERGENCE TREATMENTS**

Preemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of additional weed species listed on the preemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information that appear on the product labels. **DO NOT** apply compounds containing metolachlor after a soil incorporated Command 4 EC plus Lexone® or Sencor® tank mix treatment, unless split applications are used where the pounds of active ingredient do not exceed the total allowed.

**DO NOT** apply Canopy, Preview, Pursuit Plus, Scepter, Squadron or Tri-Scept herbicides after Command 4 EC plus any of these products in a tank mix treatment, as these other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

**COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY POSTEMERGENCE TREATMENTS**

Postemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of emerged weeds, as listed on the postemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information that appear on the product labels.

**DO NOT** apply Canopy, Pursuit Plus, Scepter, Squadron or Tri-Scept herbicides after Command 4 EC plus any of these products in a tank mix treatment, as these other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

**COMMAND 4 EC HERBICIDE FOLLOWED BY ACIFLUORFEN (BLAZER 2L OR TACKLE) HERBICIDE**

Following an application of Command 4 EC or tank mixture with Command 4 EC, a postemergence application of acifluorfen will control the following emerged broadleaf weeds:

- Broadleaf Weeds
- Cocklebur
- Flowering Spurge
- Pigweed spp.
- Black nightshade
- Wild poinsettia
- Purple Marshmallow
- Pitted Marshmallow

Apply acifluorfen at a rate of 1.0 to 1.5 pints per acre with 25 to 50 gallons concentrate to actively growing weeds that are more than 4 inches growth stage into not count to height in the case of 1 foot or taller weeds. Do not spray for mowers. This herbicide generally controls weeds that are in the first or second stage of the first to third trifoliate leaves.

**NOTE:** Application of Command 4 EC herbicide, alone or in tank mix, to soybeans that have emerged will not control weeds that are in the first or second trifoliate leaf stage.

Read and follow all precautionary statements, cautions and warnings on all product labels.

**COMMAND 4 EC HERBICIDE FOLLOWED BY SCEPTER® HERBICIDE**

Following an application of Command 4 EC or tank mixture with Command 4 EC, a postemergence application of Scepter herbicide will control the following emerged broadleaf weeds:

- Broadleaf Weeds
- Cocklebur (common)
- Pigweed
- Wild poinsettia

Apply Scepter after crop emergence but before weeds exceed a height of 12 inches to inches for wild poinsettia. Apply at a broadcast rate of 1.0 to 1.5 pints per acre. Apply when weeds are actively growing. **DO NOT** apply Scepter postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes. The total amount of Scepter should not exceed 1.0 pint per acre per season in northern areas, or 1.0 pint per acre per season in southern areas as defined on the Scepter label. Squadron and Tri-Scepter include the same active ingredient as Scepter.

For postemergence applications, the addition of a nonionic surfactant or crop oil concentrate is required. The nonionic surfactant should contain at least 80% active ingredient and should be applied at a rate of 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate stated on the COC label.

Read and follow all precautionary statements and restrictions, directions for use and all other information appearing on the Scepter label.

**COMMAND 4 EC HERBICIDE APPLICATION—MINIMUM TILLAGE OR NO-TILLAGE SOYBEANS**

Command 4 EC herbicide may be used alone or in combination with Sencor®, Lexone®, Lorox®, or Lora® herbicides, plus paraquat or Roundup® herbicide for preemergence (to crops and weeds) and post-emergence (to weeds) but prior to crop emergence, control of broadleaf and grass weeds where soybeans will be planted into a field with previous crop residues (e.g., corn, small grain stubble).

**WEEDS CONTROLLED:** When used as directed, the paraquat portion of the tank mix controls most emerged annual weeds and suppresses many perennial weeds. Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Roundup label. Command 4 EC or Command 4 EC plus Sencor®, Lexone®, or Lorox® provides preemergence control of the weeds listed in the Command 4 EC Applied Alone and Command 4 EC Applied in Combination sections of this label.

**APPLICATION AND MIXING INSTRUCTIONS**

**Paraquat Combinations:** Tank mix Command 4 EC alone or with Sencor®, Lexone®, Lorox®, or Lora®. Observe mixing instructions in the General Mixing Instructions portion of this label, as well as individual product label use directions. Add 1 to 2 pints paraquat per acre on product containing 2 pounds active ingredient per gallon per acre. Use lower rate for annual weeds less than 4 inches tall and the higher rate for weeds 4 to 6 inches tall. Add the surfactant and a suitable surfactant (refer to paraquat label).

**Roundup Combinations:** Tank mix Command 4 EC alone or with Sencor®, Lexone®, Lorox®, or Lora®. Observe mixing instructions in the General Mixing Instructions portion of this label, as well as individual product label use directions. Add 1 to 4 quarts of Roundup per acre for control of emerged annual weeds or 2 to 4 quarts per acre for control of emerged perennial and annual weeds.

Apply at 2 to 6 gallons of water per acre. Use the higher rate for a dense stand of vegetation.

**PUMPKINS**

**GENERAL INFORMATION**

Command 4 EC herbicide may be used as a preemergent soil application for the control of annual grass and broadleaf weeds in pumpkins.

Command 4 EC may be soil incorporated in the following states or as approved 24 C states for use under with that State's rules with the following states: Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, Wisconsin.

Note: After utilizing Command 4 EC (a soil incorporated treatment), incorporate Command 4 to a depth of 1 inch of soil and place the seed below the chemical barrier when planting.

Command 4 EC herbicide may be tank mixed with other pumpkins herbicides for control of weeds. Do not apply Command 4 EC herbicide to pumpkins that are in the first or second trifoliate leaf stage. Do not apply Command 4 EC herbicide to pumpkins that are in the first or second trifoliate leaf stage. Do not apply Command 4 EC herbicide to pumpkins that are in the first or second trifoliate leaf stage. Do not apply Command 4 EC herbicide to pumpkins that are in the first or second trifoliate leaf stage.

**BEST AVAILABLE COPY**



### COMMAND plus ATRAZINE

Command 4 EC at 1 to 1.5 pints (0.5 to 0.75 liter) per acre plus 1 to 2 pints of atrazine 3L or equivalent rates of atrazine 400 E or 400 G. Weeds controlled in addition to those noted under Command 4 EC alone will depend on atrazine rate used (see atrazine label). Use the higher rate of Command 4 EC when heavy weed infestation is anticipated or for longer residual control.

There is significant weed control in several trials conducted according to local Experiment Station and state or county statistics. Consult nearest Experiment Station for recommendations regarding rates for use in your area.

Refer to the atrazine label for spraying, mixing and other instructions. Follow all precautions, statements and restrictions on the atrazine label.

#### Mixing Instructions

Fill spray tank to 3/4 full with water and add recommended amount of atrazine with agitator operating. Mix thoroughly and fill tank to full working capacity. Add Command 4 EC to tank to make uniform mixture. Complete filling the spray tank with water. Mix thoroughly during mixing and application.

#### Application Instructions

Follow the atrazine Application Instructions for recommended rates.

### Dealers Should Sell in Original Packages Only.

**Terms of Sale or Use.** On purchase of this product buyer and user agree to the following conditions:

**Warranty.** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and of handling and of storage of the material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations.** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product.** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with directions or established sale practice.

**Damages.** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Loco 4/11/68

COMMAND and FMC—Trademarks of FMC Corporation  
CANOPY, LEXON and PREVIEW—Trademarks of E. I. du Pont de Nemours and Co., Inc.  
OPTION—Trademark of Hoechst Celanese Agricultural Company  
SONALAN, THE FLAN—Trademarks of E. I. du Pont de Nemours and Co., Inc.  
LASSO—Trademark of Monsanto Company  
SENCOR—Trademark of Parent Company of Farbwerke Bayer GmbH, Leverkusen  
PROWL, SCOPER, SQUADRON, TRISOPER—Trademarks of American Cyanamid Company  
DIALIN—Trademark of Ciba Geigy Corporation  
BLAZER—Trademark of BASF AG  
TACLER—Trademark of Rhone-Poulenc  
© 1968 FMC Corporation

4-13-68 revised

**BEST AVAILABLE COPY**