

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

APR 1 5 1994

Warren L. Smith
FMC
ACG SPECIALTY PRODUCTS
1735 MARKET STREET
PHILADELPHIA, PA 19103

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

16

•

Subject:

Label Amendment Submission of 02/15/94 in Response to PR Notice 93-7

EPA Reg. No. 279-3053

COMMAND 4EC

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- PEFORE selling or distributing any product bearing the final printed labeling AND
- WITHIN one year from date of this acceptance.

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

Jim Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Attachment

FMC CORP. 279-3053 02/15/94 COMMAND 4EC Original Submission

Delete the crossed-out statements on your proposed label. They are redundant statements or phrases.

Correct the typographical errors circled on your proposed label.

Command, 4 EC

Herbicide

For Agricultural or Commercial Use Only NOT FOR SALE OR USE IN CALIFORNIA

EPA Reg. No. 279-3053

EPA Est. 279-

Active Ingredient:	By Wt.
Clomazone: 2-(2-Chlorophenyl)methyl-4,	_
4-dimethyl-3-isoxazolidinone	46.7%
Inert Ingredients:*	53.3%
	100.0%

^{*}Contains xylene range aromatic solvents

Contains 4 pounds of active ingredient per gallon U.S. Patent No. 4,405,357

KEEP OUT OF REACH OF CHILDREN WARNING

-AVISO

Si usted no entiende la etiqueta, busque a algulen para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If in eyes: Flush immediately with plenty of water. Get medical attention as soon as possible.

If swallowed: Drink promptly large quantities of milk or water, /.void alcohol. Do not induce vomiting. Call a physician.

If inhaled: Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention as soon as possible.

If on skin: Wash skin with plenty of soap and water.

For Emergency Assistance Call (800) 331-3148

See other panels for additional precautionary information.

PRECAUTIONARY STATEMENTS **Hazards to Humans (and Domestic Animals)**

Causes eye injury (corneal opacity) that is temporary. Do not get in eyes or on clothing. Harmful it swallowed, inhaled, or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin. Wash-thoroughly with soap and water after handling:



FMC Corporation Agricultural Chemical Group Philadelphia PA 19103

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pents; Chemical-resistant gloves, such as Barrier Laminate or Vitori, Shoes plus socks; and Protective eyewer

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's contentrate. Do not reuse them. Follow manufacturer's instructions for cleaning maintaining PPE. If no such instructions for wishables, use detergent and hot water. Keep and wash PPE separately from other laundry.

What harder use closed systems renclosed cabs, or aircraft in a manner that meets the requirements like in the Woods Proficion Science (VPS) to a requirements pesticles (40 GF2) (0 (40 (4-8)), the requirements may be reduced of modified as specimal living WPS.

User Safety Recommendations:

Users should:

- · Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- · Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- · Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC here. bicide can cause foliar whitening or yellowing of some plants; Prior to making applications, read and strictly follow all precautions and application instructions on this label.

TABLE OF CONTENTS

Read all information on pages 1-4 before referring to specific crop

	Page
Ingredients	1
First Aid	1
Precautionary Statements	1
Directions for Use	2
Storage and Disposal	
Sprayer Cleanup	
Spray Drift Precautions	
General Mixing Instructions	
General Application Precautions	3
General Application Instructions	4
Command Geographical Areas	5
ROW CROPS	
Cotton	
Soybeans	
VEGETABLE CROPS	14
Peppers	
Pumpkins	14
Succulent Peas	15
FALLOW LAND	

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Workes Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on ferms, forests, nurseries, and greenhouses, and handlers of agricultural posticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions, pertaining to the statements on this label about personal protective equipment (RPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker party into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Erotection Standard under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated; such as plants, soil, or water, is: Coverals; Chemical-resistant gloves; such as Barrier Laminate or Viton; Shoes plus socks; and Protective eyewear.

STORAGE AND DISPOSAL

Pesticide Storage
Do not freeze. Do not store below 40°F. If solid crystals are observed, warm material to above 60°F by placing container in warm location. Shake or roll container periodically to redissolve

Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers, After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC (800) 331-

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Régional Office for guidance.

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If humed, stay out of smoke.

Returnable/Refillable Sealed Containers

Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase,

- Some species of trees, shrubs, flowers, agronomic crops, and fruits and vegetables are sensitive to Command® 4 EC herbicide.
- Foliar contact with spray drift or vapors may cause whitening of sensitive plants. Symptoms are generally temporary in nature but may persist on some plants.
- Plants such as the following may show symptoms of toliar whitening or yellowing if contacted by Command 4 EC herbicide.

Trees (Evergreen) Trees (Deciduous) Fir Species Apple (inc. fruit & Spruce species
Shrubs & Vines
Azeles omamental types)
Ash (Green; White,
Mountain;
Basswood)
Botelder (Marie Catalpa Marie Catalpa Marie Catalpa Marie Chemy (Inc. Iruk & Burningbush A (Winged Euonymus) Grape Grape Honeysuckie Content (Inc. Iroll &

Cottonwood

Em The

Ginkgos

Hackberry

Mulbary

Peach

Pear Inc. In It Roses Signature
Yew Manager
Autonomic Crops
Allalia gronomic Crops Alfalia Oats Vegetable and Pear (inc. fruit & ornamental ... Flower plants types)
Pecan
Pecan
Russian olive
Tree-of Heaven
Tulip tree <u>Others</u> Ferns, Herbs 📜 Strawberry Tulip tree Walnut trees Raspberry 🎊 Blackberry Willow species Do not apply Command 4 EC herbicide to non-field areas

SPRAYER CLEANUP

Do not drain or flush equipment on or near desirable trees or other plants, or in areas where their roots may extend or in locations where the chemical may be washed or move into contact with their roots. Do not contaminate any body of water including irrigation water that may be used on other crops. Carefully follow sprayer clean-up instructions noted below to prevent spray tank residues from damaging other crops.

including fence rows, waterways, ditches, and roadsides......

Sprayer equipment should be thoroughly rinsed to remove residues of herbicide that might injure other subsequently sprayed crops. The steps below should be followed for the thorough rinsing of spray equipment following applications of Command 4 EC herbicide or tank mixes of Command 4 EC with other labeled products.

1) Drain any remaining spray solution from tank, pump, hoses and boom and discard in an approved manner (See Note that follows).

2) Clean tank and fittings by:

. Thoroughly hosing down the inside walls of the spray tank with a quantity of water equal to 1/s of the total tank capacity and operating the pump to circulate this solution through the sprayer system for 15 minutes.

Washing down the outside surfaces of equipment.

- Removing nozzle tip and screen from end nozzle in each boom section and allowing several gallons of rinsate solution to flush completely through boom (collect rinsate while flushing).
- 3) Thoroughly drain remaining rinsate solution from tank, pump and hoses. Combine with boom flushing and dispose of all rinsates from this first rinsing in an approved manner (see Note that follows).
 - When switching from water dilutions to applications utilizing crop oil or liquid fertilizer as a carrier, a small volume of crop oil or liquid fertilizer should be flushed through the tank, pump, hoses, and boom prior to the next use. Dispose of crop oil or liquid fertilizer rinsate in an approved manner (see Note for local, state and federal quidelines).
- 4) Remove the remaining nozzle tips, and screens and the line filter and wash in a pail of warm soapy water, thoroughly rinse and replace.
- 5) Hose down the inside walls of the spray tank a second time and circulate this solution using the same procedure as noted in #2 above.



preplant incorporated pesticide on any crop for which Command 4 EC is registered, rinsate from this second rinsing may be utilized by diluting with the water for the next pesticide load;

HOWEVER

If the next use of the sprayer will be a postemergence applied pesticide on any crop, drain rinsate solution from this second rinsing. Retain rinsate solution for use only with a preemergent soil incorporated pesticide to be applied on a crop and in a manner for which Command 4 EC is registered. Refill tank (after draining second rinsate solution) in accordance with post-emergence product label directions.

NOTE: Dispose of excess spray mixture and/or rinsate from first tank rinsing by application to cropland as described on this label. If excess spray mixture and/or rinsate from first rinsing cannot be disposed of according to label instructions, dispose of in compliance with local, state and federal guidelines. Contact your state pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance.

GENERAL MIXING INSTRUCTIONS

Care must be taken when mixing Command 4 EC herbicide. Avoid mixing in areas adjacent to desirable plants (See Section headed "IMPORTANT")

Command 4 EC Alone: Mix Command 4 EC with water or liquid fertilizer in the following manner: Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 4 EC, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

Tank Mixtures: Fill spray tank one-fourth to one-third full with water, with agitator operating add the recommended amount of ingredients using the following order: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's). Mix thoroughly and fill tank one-half full continuing agitation. Addition and the sprayer tank with water. Where use of a surfactant is recommended, add as the last ingredient to the spray tank. Maintain agitation during filling, mixing and application. When using drift reducing additives, follow specific product label instructions for the order of addition of all products to the spray tank. Fertilizer Spray Mixtures: Applications of all products to the spray tank. Fertilizer Spray Mixtures: Applications of all products to the spray tank. Fertilizer Spray Mixtures: Applications of with fertilizer solutions may be used unless use directions specifically state otherwise. Small quan-tities should be tested for compatibility by the following procedure before mixing tankful quantities.

- 1) Put 1 pint of fertilizer solution in a quart jar.
- 2) Add the appropriate amount of herbicide based on the table at the top of page 3. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).

Herbicide	Rate/Acre	Amount To Be Added Per Pint of Fluid Fertilizer
Wettable Powders	1/2 pound	3/4 level teaspoon 1 1/2 level teaspoons
or Dry Flowables (Dry Granules)	1 pound 2 pounds 3 pounds	3 level teaspoors, 41/2 level teaspoons
Emulsifiable	1 pint	1/2 level teaspoon
Concentrates	1 quart	1 level teaspoon
Or	2 quarts	2 level teaspoons
Flowables	3 quarts	3 level teaspoons

- Based on a spray volume of 25 gpa. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.
- 3) Close jar and shake well.
- 4) Watch mixture for several seconds, again in 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, get or become (umpy), the mixture is compatible and can be mixed in full volumes and sprayed. If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below:
- Command 4 EC Alone: Dilute with 2 parts of water and add slowly to spray tank of fertilizer with agitator running. Maintain agitation during application.
- 2) Command 4 EC plus Tank Mixtures: Dilute individual products with two (2) parts of water, then add to the spray tank of fertilizer, while maintaining agitation, using the following order—sturry of dry formulations (wettable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Continue agitation during application.


GENERAL APPLICATION PRECAUTIONS

Do not apply aerially or through irrigation equipment.

geography) or there is an approved SLN (24C) for counties within these states:

Connecticut Delaware Michigan Minnesota Minnesota New Hampshire Iowa New Jersey Maine New York Maryland Ohio

Pennsylvania Rhode Island Vermont Virginia West Virginia Wisconsin



For Soil Incorporated Applications: See Geographic Restrictions above.

Sensitive plants may show symptoms of foliar whitening or yellowing if contacted by Command 4 EC herbicide. It is recommended that prior to application, adjacent properties be checked and that spraying within 200 feet of desirable plants be avoided.

avoided.

Do not apply Command 4 EC herbicide to non-field areas including fence rows, waterways, ditches, and roadsides.

Do not apply Command 4 EC within 1,000 feet of the following areas: Towns and Subdivisions, Commercial Fruit/Nut of Vegetable* Production, Commercial Greenhouses or Nursedes.

Apply Command 4 EC herbicide only to surfaces that will be soil incorporated unless specific use directions and precautions permit otherwise

*except sweet com

For Preemergent Applications: See Geographic Restrictions above.

Except as noted above (see Geographic Restrictions), Command may be used as a preemergent soil applied treatment without incorporation, however, additional precautions must be taken as follows:

Do not apply Command 4 EC herbicide to non-field areas including tence rows waterways, ditches, and roadsides.

Do not apply Command 4 EC within 1,500 feet of the following areas: Towns and Subdivisions, Commercial Fruit/Nut or Vegetable Production, Commercial Greenhouses or Nurseries.

Sensitive plants may show symptoms of foliar whitening or yellowing if contacted by Command 4 EC herbicide. Prior to application, adjacent properties must be checked, and spraying within 1,500 feet of desirable plants must be avoided.

Refer to individual crop use directions for additional requirements.

*except sweet com. *



prevent whitening or desirable plants. Unit is minoripod by in which include wind speed, spray pressure, particle size, nozzle type, and boom height.

Do not apply when weather conditions favor drift.

Do not apply in winds above 10 miles per hour.

A minimum spray volume of 10 gallons per acre is recommended with appropriate nozzle types and sizes that produce coarser sprays

The use of agriculturally approved drift reducing additives is recommended for application volumes of 15 to 40 gallons per acre when spraying in the proximity of sensitive plants (see list on page 2).

The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre.

Use minimum nozzle pressure and boom height while maintaining

uniform spray pattern.

Do not exceed 30 psi spray pressure.

Selection and proper use of spray equipment is critical in minimizing spray drift. The following table suggests pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types.

When moving spray equipment to noncontiguous sites, do not allow spray solution to spray or drip from tanks, hoses, fittings or spray

nozzles and tips.

Suggested Nozzle Types, Minimum Size and Recommended Pressure Ranges for Minimizing Drift (See Special Precautions in the cotton use section for further restrictions)

Nozzle	Pressure Range (PSI)	Minimum Flow Rate Within Pressure Range (GPM)	Minimum Nozzle Size
Flat-fan	15-30	0.3	#4*
LP-flat-fan	10-25	0.3	#3
Even flat-fan**	15-30	0.3	#4
Flood	10-25	0.3	#2.5°
Whirl-chamber	5-20	0.3	#5
Raindrop	15-30	0.15	#2
Wide angle	_		
full cone	15-30	0.3	#5

Refers to tip number such as 8004 or LF2.5.

ROTATIONAL CROPPING PRECAUTIONS

Under some conditions, temporary whitening or yellowing of leaves may occur on approved rotational crops where undesirable soil residues of Command 4 EC exist.

Under abnormal conditions, carryover injury to rotational crops can occur. The following factors can contribute to increased risk of injury to rotational crops:

Exceeding label recommended rates.

Overapplication resulting from use of worn nozzles, excessive overlapping of spray swaths, failing to shut off spray booms when turning (end row areas), or slowing or stopping sprayer.

Soil with pH less than or equal to 5.9.

- Extreme dryness in the four months following application., Use of organophosphate soil insecticide followed by use of some 5) postemergence com herbicides.

Choice of rotational crop hybrid.

Additional recommendations to prevent rotational crop injury may be provided in the form of service bulletins for location where risk of injury is significantly increased due to extremely dry conditions.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications read and strictly follow all precautions and application instructions on this label.

GENERAL APPLICATION INSTRUCTIONS

Ground Applications

Broadcast or Banded Applications: Apply Command 4 EC alone or in tank mix combinations by ground equipment using a finished spray volume of 10 to 40 gallons of water per acre. NOTE: The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre. Use nozzles suitable for broadcast boom or banded application of herbicides. Coarse sprays are less likely to drift out of the target area than fine sprays. See "APPLICATION PRECAUTIONS" section for specific recommendations to reduce spray drift. For Command 4 EC tank mixtures with wettable powder or dry flowable formulations, nozzle screens and strainers should be no finer than 50-mesh.

Band width Broadcast Band rate in inches Rate Per = per acre Row width Acre in inches Band width x Broadcast Band volume in inches Volume **Row width** per acre Per Acre in inches

Follow broadcast minimum gallonage requirements adjusted to band width. Do not exceed 30 psi spray pressure.

Refer to Spray Drift Precautions for suggested pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types. For banded applications an even flat-fan nozzle is recommended

Soil Incorporated Application: Apply Command 4 EC to the soil surface and uniformly incorporate. Select equipment suited for proper incorporation. Shallow incorporation normally results in better weed control than a deep incorporation.

In states where soil incorporation is required (see Geographical Restrictions) application and immediate incorporation to a maximum depth of 3 inches is required unless the soil is dry. On dry soils, incorporation to a maximum depth of 3 inches must be completed within 8 hours of Command 4 EC herbicide application.

Soil must be in good tilth to allow for thorough mixing of the soil. Do not apply under dry, windy conditions which allow movement of soil from treated areas. Do not apply under wet soil conditions. Application to wet soils will increase the potential for off-site movement of Command 4 EC herbicide vapors and may result in poor soil incorporation and unsatisfactory weed control.

INCORPORATION EQUIPMENT

ONE PASS INCORPORATION

Suggested Implements

For one pass incorporation use C or S shank field cultivators equipped with a leveling device such as a spike or coil time harrow or rolling basket attachment. Field cultivator must have 3 or 4 rows of sweeps spaced at intervals of 7 inches or less and staggered so that no soil is left unturned. Equipment design, including any drag attachments, must be adequate to avoid soil ridging which may result in streaked or reduced waed control. The implement should be operated at 6 to 8 mph and set to cut 3 to 4 inches deep to incorporate herbicide 1 to 2 inches.

Combination equipment (three or more tillage devices combined and used as a single tool) should be operated at a depth of 3 to 4 inches and a minimum speed of 6 mph.

PTO-drive seed bed preparation equipment should be set to cut 2 inches deep and operated at 4 mph or less.

TWO PASS INCORPORATION

Suggested Implements

When using two pass incorporation, the second pass should be at an angle to the first pass.

Disc set to cut 3 to 4 inches deep at 4 to 6 mph. Blades should be less than 20 inches in diameter and spaced 9 inches or less apart.

Rolling cultivator set to cut 2 to 3 inches deep at 6 to 8 mph (use on coarse or medium textured soils only).

Field cultivator with 3 to 4 rows of sweeps spaced at 7-inch intervals or less and set to cut 3 inches deep at 5 mph or greater.

Cultivation After Planting

Soil treated with Command 4 EC may be shallow-cultivated, rotary-hoed or hand-hoed without reducing the weed control activity of Command 4 EC. Do not cultivate deeper than the Command 4 EC freated layer of soil since this may bring untreated soil to the surface and poor weed control may result.

COMMERCIAL IMPREGNATION AND APPLICATION OF COMMAND 4 EC ON DRY BULK FERTILIZERS

Command herbicide may be impregnated on dry bulk fertilizers. When applied as directed Command/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of Command applied in

The Command/fertilizer mixtures must be soil incorporated if required according to the Geographic Restrictions of the Application Precautions section of this label.

BEST AVAILABLE COPY

^{**} Recommended for banded application.

per acre and up to a maximum of 450 pounds per acre with the recommended amount of Command 4 EC herbicide per acre. Use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform, fine spray pattern over the tumbling fertilizer for thorough coverage. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Command provides a satisfactory, dry mixture. If the absorptive capacity is inadequate use of a highly absorptive powder is required to provide a dry, flowable mixture. Microcel E (Johns-Manvill Products Corporation) is a recommended absorbent powder. Generally less than 2% by weight of Microcel E is required. DO NOT impregnate Command 4 EC onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with Command.

The amount of Command actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with Command 4 EC herbicide should be applied immediately, not stored. Care should be taken to ensure spreading procedures in the field do not place fertilizer treated with Command in areas which cannot be incorporated, if incorporation is required within the Geographic Restrictions of this label.

All state regulations, labeling, etc., of the mixtures are the responsibility of the seller.

For those rates not listed in the following table, calculate the amount of Command to be impregnated on a ton of dry bulk fertilizer using the following formula:

2000	PINTS of Command	PINTS of
pounds dry	× per acre =	Command per ton
fertilizer	(recommended rate	of fertilizer
per acre	for soil texture)	

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZERS WITH COMMAND 4 EC HERBICIDE


PINTS OF COMMAND 4 EC PER TON OF FERTILIZER

Fertilizer	Command 4 EC Rate Per Acre		
Rate Lbs/Acre	1½ pts. (.75 lbs A.I.)	2 pt. (1 lb A.l.)	2½ pts. (1.25 lbs A.I.)
200	15	20	25
250	12	16	20
300	10	13 1/3	16 3/3
350	8 1/2	11 1/2	14
400	7 1/2	10	12 1/2
450	6 7 ⁄3	9	11

If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fert lizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

SPECIAL PRECAUTIONS FOR DRY BULK FERTILIZER IMPREGNATION AND APPLICATION WITH COMMAND 4 EC

- All equipment used to apply Command must be thoroughly cleaned immediately following use to ensure no contamination results which could cause injury to non-labeled crops or desirable vegetation. Refer to "Sprayer Cleanup" section for additional details and disposal of rinsates.
- Equipment used to physically transport Command treated fertilizer, including boots, augers, conveyers, bins, etc. must be covered to prevent loss of fine particles an bjected to clear up procedures previously described.
- Mixing of Command herbicide and dry fertilizer must be conducted in a blender which will ensure that airborne particles and potential for volatilization are avoided.
- Bulk containers must be tightly covered while the product is being transported and applied to reduce chances of Command volatilization or product loss.
- It is recommended that Dry Bulk Fertilizer impregnation operations. using Command not be conducted within 1/4 mile of residential areas or areas where Command symptomology or desirable vegetation would not be tolerated.



Southern area includes Missouri bootheel which incorporates the following counties: Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott, and Stoddard.

COTTON

DIRECTIONS FOR USE

Command 4EC Herbicide may be applied, as a preemergent banded or a soil incorporated broadcast treatment in conventional tillage or conservation tillage systems for the control of annual grasses and broadleaf weeds in cotton.

SPECIAL PRECAUTIONS FOR COTTON

Off-site movement of spray drift or vapors of Command® 4 EC. herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

Additional Precautions For Cotton Applications:

Completion of proper qualification training required for sale, purchase, and use.

Do not apply aerially or through irrigation equipment.

Observe all buffer restrictions.

Do not apply Command 4 EC within 1,500 feet for preemergent applications or 1,000 feet for soil incorporated applications of the 4 following areas: Towns and Subdivisions, Commercial Fruit/Nut or :: Vegetable* Production, Commercial Greenhouses or Nurseries,

except sweet com.

Prior to application, adjacent properties must be checked, and preemergent spraying within 1,500 feet or soil incorporated spraying within 200 feet of desirable plants must be avoided.

Before application, determine air movement and direction.

Do not apply in winds above 10 miles per hour,

Do not apply without an FMC approved drift reducing additive. Follow specific product label instructions for the order of addition. of all products to the spray tank.

Do not exceed 30 psi spray pressure.

Do not apply at boom heights above 24 inches.

Do not apply at boom neights above 24 inches.

Do not apply with nozzles other than flood, whirtchamber, or Raindrop[®] or similar tips that produce coarse sprays. Refer to nozzle table under SPRAY DRIFT PRECAUTIONS.

REQUIREMENTS FOR PLANTING TIME **APPLICATIONS**

Do not apply Command 4EC Herbicide to cotton unless either disulfoton or phorate organophosphate insecticide is applied in-furrow with the seed at planting time at a minimum of 0.75 pound per acre of active ingredient. Do not reduce the application rate of the organophosphate insecticide when Command 4EC herbicide is applied as a banded treatment. Failure to apply either disulfoton or phorate insecticides with Command in accordance with in-furrow label use directions can result in crop phytotoxicity (bleaching) and/or stand reduction. Combinations of at planting systemic granular carbamate and organophosphate insecticides in conjunction with Command may result in injury to cotton. Crop injury may occur with higher Command rates on sandy soils. Diuron is not recommended at planting when Command is used as plant injury may

Refer to the insecticide product labels for appropriate in-furrow application directions and maximum use rates. Monitor application equipment to insure accurate and uniform placement of the insecticide.

Do not apply Command 4 EC Herbicide as a broadcast treatment in cotton without soil incorporation.

SOIL INCORPORATED APPLICATIONS

For soil incorporated treatments, apply Command as an incorporated treatment prior to or immediately before planting. Refer to the Soil Incorporated Application section of the GENERAL APPLICATION INSTRUCTIONS for further directions.

REPLANTING INSTRUCTIONS

If the initial planting of cotton fails to produce a uniform stand, cotton may be replanted in fields treated with Command 4EC alone. Do not retreat fields with a second application of Command 4EC. When tank mixing with a labeled product, refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP GUIDELINES on the Command 4EC label. Where a tank mix is used, refer to the product's labels for any additional replant instructions.

If replanting is required follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS noted above.

Command 4 EC Herbicide Applied Alone

RATES AND WEEDS CONTROLLED'

1.0 PT.2	1.5PT.	2.0 PT.	2.5 PT.
(0.50 LB. A.I.)	(0.75 LB, A.I.)	(1.0 LB. A.I.)	(1.25 LB. A.L)
GRASSES	<u>GRASSES</u>	GRASSES	GRASSES
Suppression of some annual grasses	Barmyardgrass Broadleaf Signalgrass CrabgrassLargeSmooth FoxtailGiantGreen Goosegrass PanicumCommonFallTexas Seedling Johnsongrass	All grasses controlled at 1.5 pints plus: Cupgrass —South- western —Woolty Field Sandbur Bermudagrass* Red Rice* Itchgrass*	All grasses controlled at 2.0 pints plus; Foxtail —Purple —Robust —Yellow Junglerice* Shattercane* Wild Proso Millet*

1.5PT.

(0.50 LB. A.I.)	(0.75 LB. A.I.)
BROADLEAVES	BROADLEAVE
Velvetleaf Spurred Anoda	Velvetleaf Spurred Anoda plus: Common Ragweed* Jimsonweed* Lambsquarter Pennsylvania Smartweed* Prickly Sida Purslane Redweed
	Venice Mallow

(1.0 LB. A.I.) (1.25 LB, A.I.) S BROADLEAVES BROADLEAVES All broadleaves All broadleaves controlled at controlled at 1.5 pints *plus:* 2.0 pints plus: Cocklebur **Balloonvine** Black Common Nightshade* Curly Dock Joint Vetch* Ragweed Dayflower Florida Beggarweed Kochia Florida Pusley Morningglory Jimsonweed -Pitted Kochia¹ Pennsylvania Redvine* Smartweed Tropic Croton Pigweed Wild Poinsettia⁴ -Redroot*

2.0 PT.

*Partially Controlled

1.0 PT.2

¹ Higher label rates (a minimum of 2 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, a minimum application rate of 2½ pints per acre for cocklebur or common ragweed control is recommended.

²For heavy weed pressure or heavy soils use 1.25 pints per acre to control Velveteat or Spurred Anoda.

For control of itchgrass, do not incorporate Command 4 EC. For high weed pressure and heavy soils use 2.5 pints.

*Use In AR/LA/MS/TX/OK only—for high pressure and heavy soils use 2.5 pints.

COMMAND 4 EC HERBICIDE MAY BE APPLIED WITH OTHER HERBICIDES REGISTERED FOR THESE USES ON COTTON.

Command 4EC may be applied in combination with other herbicides registered for use in cotton to broaden the weed control spectrum. Refer to the labels of the other herbicides for additional information. Carefully observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Note: When using Command 4EC with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

1.0 PT. (0.50 LB. A.I.)	1.5 to 2.6 PT. (0.75 to 1.0 LB. A.I.)	2.5 PT. (1.25 LB. A.I.)
ANYTIME	ANYTIME	ANYTIME
Cotton° Soybeans Peppers Pumpkins Peas	Cotton* Soybeans Pumpkins Peppers	Cotton* Soybeans
9 MONTHS	9 MONTHS	9 MONTHS
Com —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Potatoes Pumpkins Rice Snap Beans Sorghum Sugar Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)	Same as 1.0 pint, plus: Peas	Same as 1.5 to 2.0 pints, <i>plus</i> ; Pumpkins
12 MONTHS	12 MONTHS	12 MONTHS
All crops	All crops	All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4EC treatment.

"Refer to the section "Requirements for Planting Time Applications" and "Replanting Instructions."

RESTRICTIONS

Do not apply Command within 65 days of harvest.

Do not apply more than 1.25 pounds active ingredient per acre per season,

Do not allow livestock to graze on treated cotton forage or trash, or feed treated cotton forage or trash to livestock.

SOYBEANS

GENERAL INFORMATION

Command 4 EC herbicide may be utilized as a preemergent soil applied or soil incorporated treatment for the control of annual grass and broadleaf weeds in soybeans.

Command 4 EC must be applied as a soil incorporated treatment in the following states, unless early preplant (EPP) applications are allowed (see Conservation Tillage Soybeans for specified geography) or there is an approved SLN (24C) for counties within these states (check with your local dealer): Connecticut, Delaware, Iliinois, Indiana, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, and Wisconsin.

Command 4 EC herbicide may be tank mixed with other herbicidos to broaden the weed control spectrum compared to the products applied alone. Command 4 EC may be tank mixed with Canopy®, Dual®, Lasso®, Lexone®, Lorox®, Lorox Plus®, Preview®, Prowl®, Scepter®, Sencor®, Sonalan®, Squadron®, Tri-Scept®, or other registered herbicides. Observe all precautions instructions, and rotational cropping guidelines of each product's label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with the herbicides listed above, unless use directions specifically state otherwise.

REPLANTING INSTRUCTIONS

If initial seeding of soybeans fails to produce a stand, soybeans may be replanted in fields treated with Command 4 EC alone (or with recommended tank mixtures). Do not retreat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the soybean replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Guidelines on this label.

Northern Area-See Map **COMMAND® 4 EC HERBICIDE** APPLIED ALONE RATES AND WEEDS CONTROLLED'

1 0 PT. ²	1.5 PT.	2.0 PT.
(0.50 LB. A.I.)	(0.75 LB. A.I.)	(1.0 LB. A.I.)
GRASSES	GRASSES	GRASSES
Suppression of	Barnyardgrass	All grasses controlled at
some annual	Crabgrass	1.5 pints <i>plus:</i>
grasses	Large	Bernudagrass*
•	-Smooth	Broadleaf Signalgrass
	Field Sandbur	Cupgrass
	Foxtail	—Southwestern
	Giant	—Woolly*
	Green	Seedling Johnsongrass
	-Robust	Shattercane*
	Yellow	Wild Proso Millet*
	Goosegrass	
	Panicum —Common	
	—Fall	
	—Texas	
	Seedling	
	Johnsongrass*	
BROADLEAVES	BROADLEAVES	BROADLEAVES
Velvetleaf	Velvetleaf	All broadleaves
Spurred Anoda	Spurred Anoda	controlled at 1.5 pints
Spaned Alkoda	plus:	plus:
	Common Ragweed*	Black Seeded Plantain
	Galinsoga	Cocklebur*
	Jimsonweed*	Common Ragweed
	Lambsquarters	Jimsonweed
	Prickly Sida	Kochia
	Purslane	Smartweed
	Smartweed*	—Ladysthumb
	-Ladysthumb*	—Pennsylvania
	Pennsylvania*	Tropic Croton
*Partially Controlled	Venice Mallow	

Partially Controlled

NOTE: When using Command 4 EC more than 30 days prior to planting. the highest label rate should be used.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur Refer to sections headed "Rotational Cropping Precautions" and "General Information."

NOTE: When using Command 4 EC with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

1.0 PT. (0.50 LB. A.i.)	1.5 PT. (0.75 LB. A.I.)	2.0 PT. (1.0 LB, A.I.)
ANYTIME	ANYTIME	ANYTIME
Scybeans Cotton*	Soybeans Cotton*	Soybeans Cotton*
Peppors Pumpkins	Peppers Pumpkins	Peppers Pumpkins
Peas	•	•
9 MONTHS Corn	9 MONTHS	9 MONTHS
Field	Same as 1.0 pint, plus:	Same as 1.5 pints, except
—Pop —Seed	Peas	Seed Com and Sweet Com require a 12
—Sweet Cotton		month rotation.
COROL		

Cucurbits Dry Beans Peanuts Peopers **Potatoes** Rice Snap Beans Sorghum Sugar Beets **Sweet Potatoes** Tobacco Tomatoes (Transplanted)

12 MONTHS Cabbage Tomatoes (all) Wheat	12 MONTHS Same as 1 0 pint.	12 MONTHS Same as 1.5 pints, plus: Seed Corn Sweet Corn
16 MONTHS	16 MONTHS	16 MONTHS
All crops	All crops	All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

2 n PT

25 DT

Prostrate Spurge

Southern Area—See Map

1 A DT?

COMMAND® 4 EC HERBICIDE APPLIED ALONE RATES AND WEEDS CONTROLLED'

1 5 PT

(0.50 LB. A.I.)	(0.75 LB. A.I.)	(1.0 LB. A.I.)	2.5 Pl. (1.25 LB. A.I.)
<u>GRASSES</u>	GRASSES	<u>GRASSES</u>	<u>GRASSES</u>
Suppression of some annual grasses	Bamyardgrass Broadleaf Signalgrass Crabgrass —Large —Smooth Foxtail —Giant —Green Goosegrass Panicum —Commc·, —Fall —Texas Seedling Johnsongrass Field Sandbur*	All grasses controlled at 1.5 pints plus: Cupgrass —South—westem —Woolly Field Sandbur Bermudagrass* Red Rice Itchgrass³	All grasses controlled at 2.0 pints plus: Foxtail —Purple —Robust —Yellow Junglerice* Shattercane* Wild Proso Miltet*
BROADLEAVES Velvetleaf Spurred Anoda	BROADLEAVES Velvetleaf Spurred Anoda plus: Common Ragweed* Jimsonweed* Lambsquarter Pennsylvania Smartweed* Prickly Sida Purslane Redweed Venice Mallow	BROADLEAVES All broadleaves controlled at 1.5 pints plus: Cocklebur Common Ragweed Dayllower Florida Begganveed Florida Pusley Jimsonweed Kochia* Redvine* Tropic Croton Wild Poinsettia*	BROADL EAVES All broadleaves controlled at 2.0 pints plus: Balloonvine* Black Nightshade* Curty Dock Joint Vetch* Kochia Morningglory —Pitted Pennsylvania Smartweed Pigweed —Redroot* —Smooth*

* Partially Controlled

² For heavy weed pressure or heavy soils use 1.25 pints per acre to control Velvetical or Spurred Anoda.

^a For control of litchgrass, do not incorporate Command 4 EC. For high weed pressure and heavy soils use 2.5 plnts.

4 Use In AR/LA/MS/TX/OK only—for high pressure and heavy soils use 2.5 pints.

NOTE: When using Command 4 EC more than 30 days prior to planting, the highest label rate should be used.

¹Higher label rates (a minimum of 2 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavler soil types.

²For heavy weed pressure or heavy soils use 1.25 pints per acre to control Velvetleaf or Spurred Anoda.

^{*}Refer to the "Requirements for Planting Time Applications" and "Replanting Instructions" in the Cotton Directions for Use section.

¹ Higher label rates (a minimum of 2 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, a minimum application rate of 21/2 pints per acre for cocklebur or common ragweed control is recommended.

ROTATIONAL CHUP GUIDELINES

Route to Crops as listed below, otherwise crop injury may Occur.

Refer to sections headed "Rotational Cropping Precautions" and "General Information.

NOTE: When using Command 4 EC with other registered herbicides always refer to rotational restrictions and precautions on the other

product's label. 1.5 to 2.0 PT. .0 LB. A.I.)

1.0 P I.	1.5 10
(0.50 LB. A.I.)	(0.75 to 1.
NYTIME	ANYTIME

(1.25 LB. A.I.)

Soybeans Cotton* **Peppers Pumpkins**

Soybeans Soybeans Cotton* Cotton*

Peopers **Pumokins**

9 MONTHS Same as 1.0 pint, plus: Peas

9 MONTHS Same as 1.5 to 2.0 pints, *plus:* Pumpkins

-Pop -Seed -Sweet Cotton Cucurbits Dry Beans Peanuts **Potatoes Pumpkins** Rice Snap Beans Sorahum Sugar Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)

Peas

Corn

9 MONTHS

--Fiek

12 MONTHS All crops

12 MONTHS All crops

12 MONTHS All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

*Refer to the "Requirements for Planting Time Applications" and "Replanting Instructions" in the Cotton Directions for Use section.

COMMAND® 4 EC HERBICIDE APPLIED IN COMBINATIONS FOR SOYBEANS

Command 4 EC herbicide may be tank mixed with other soybean herbicides to broaden the weed control spectrum compared to the products applied alone. In the sections that follow FMC recommends a combination of products and application rates to broaden weed control spectrum. FMC makes no claims as to weed control for any combination of rates or products other than as recommended in this label. Additional recommendations for product use with other herbicides not included in the following sections may become available subsequent to the date of publicat on for this label. Please check with your dealer for more recent FMC technical bulletins and label supplements. When applying mixtures of Command 4 EC with other herbicides, observe all precautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixes.

Command 4 EC Herbicide plus Canopy® Herbicide— Northern & Southern Areas

Tank mix combination for broadened weed control spectrum.

Broadcast Rates Per Acre*

Nonn	South	Canopy	
1½ to 2 pints	2 to 21/2 pints	4 to 6 oz.	
1½ to 2 pints	2 to 2½ pints	5 to 8 oz.	
1½ to 2 pints	2 to 2½ pints	6 to 8 oz.	
	North 1½ to 2 pints 1½ to 2 pints 1½ to 2	1½ to 2 2 to 2½ pints 1½ to 2 2 to 2½ pints 1½ to 2 2 to 2½ pints 1½ to 2 2 to 2½	

Select lower to higher rates within the range to correspond to the coarser to liner soil types within the textural group.

Rate within the range should correspond to anticipated weed pressure.

" Do not apply to soil, less than 1/2% Organic Matter.

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

TTELUU WWITH WELLS.

Broadleaf Weeds All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following: Cocklebur** Copperleaf, hophombeam* Hemp sesbania Morning-glory* (Annual, Ivy-leaf, Entireleaf, Pitted, Tall) Pigweed (Redroot, Smooth)

Sicklepod^{*}

Partially Controlled

"Large seeded weeds, germinating deep in the soil, such as morningglory, sicidepod and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a postemergence herbicide for season long control.

When applying mixtures of Command 4 EC plus Canopy, observe all cautions and limitations appearing on the labels of each product. Special attention must be paid to sections regarding planting depth, phosphate pesticide interaction, previous pesticides, and sensitive soybean varieties on the Canopy label. Observe application precautions rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Dual® 8 E Herbicide— Northern & Southern Areas

Tank mix combination for broadened weed control spectrum.

Broadcast Rates Per Acre

	Com		
Soil Texture	North	South	Dual 8 E
Coarse (light) Soils: (loamy sand, sandy loam)	1½ to 2 pints	2 to 21/2 pints	1½ to 2 pints
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	2 to 2½ pints	2 to 21/2 pints
Fine (heavy) Soils: (silty clay, clay loam, clay)	1½ to 2 pints	2 to 21/2 pints	2 to 21/2 pints

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Nutsedge Red Rice Witchgrass Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Black Nightshade Carpetweed Pigweed

When applying mixtures of Command 4 EC plus Dual, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Lasso® or Lasso® Micro-Tech—Northern & Southern Areas

A tank mix combination of Command 4 EC herbicide and Lasso or Lasso Micro-Tech herbicide will broaden the weed control spectrum compared to either product used alone.

Broadcast Rates Per Acre

	Broadcast Hates Per Acre			
Soil Texture	Com North	Lasso*/Lasso Micro-Tech		
Coarse (light) Soils: (loamy sand, sandy loam)	1½ to 2 pints	2 to 2½ pints	2 to 2½ quarts	
Medium Soils: (k.am, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	2 to 2½ pints	2½ to 3 quarts	
Fine (heavy) Soils: (sitty clay, clay loam, sitty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	2½ to 3 quarts	

Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., sand, loamy sand, sandy loam) within a textural group.

Use higher rates in areas with heavy pressure.

Grass Weeds All species listed under the section Command Applied Alone Rates and Weeds Controlled

Broadleaf Weeds All species listed under the section Command Applied Alone Hates and Weeds Controlled, plus the following. Black Nightshade Carpetweed Pigweed

When applying mixtures of Command 4 EC herbicide plus Lasso or Lasso Micro-Tech, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Lorox®
A preemergence tank mix combination of Command 4 EC herbicide and Lorox herbicide will broaden the weed control spectrum compared to either product alone.

The tank mix combination may not be incorporated. Where Command 4 EC must be incorporated (see Geographical Restrictions), apply Lorox as an overlay treatment.

Broadcast	Rates	Per	Acre'

	Broaccast Hates Per Acre			
Soil Texture**	Command North South		Lorox DF+	
Coarse (light) Soils: (loamy sand**, sandy loam)	1½ to 2 pints	2 to 21/s pints	²/s to 1 lb.	
Medium Soile: (loam, silt, silt loam, sandy day, sandy clay loam)	1½ to 2 pints	2 to 21/e pints	1 to 1½ lbs.	
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	1 1/4 to 2 lbs.	
	<u> </u>	<u> </u>		

- Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, sitt loam, sandy clay, sandy clay loam) within a textural
- Plant seed at least 1% inches deep or Lorox injury to soybeans may result. Refer to Lorox label for other planting and application precautions, one pint of Lorox L equals one pound of Lorox DF, use equivalent rates.
- Do not use on sand, loamy sand, gravelly soils, or exposed subsoils or soils with less than 1% organic matter. Use only on loamy sand with over 1% organic

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

<u>Broadleaf Weeds</u>

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Carpetweed Chickweed Pigweed (Redroot, Smooth) Wild Mustard Wild Radish

When applying mixtures of Command plus Lorox observe all cautions and limitations appearing on the labels of each product. Observe application precautions, rotational crop/grazing and feeding restrictions and replanting instructions sections of this label when using tank mixtures.

Command 4 EC Herbicide plus Lorox Pluse

A preemergence tank mix combination of Command 4 EC herbicide and Lorox Plus herbicide will broaden the weed control spectrum compared to either product alone.

The tank mix combination may not be incorporated. Where Command 4 EC must be incorporated (see Geographical Restrictions), apply Lorox Plus as an overlay treatment.

	Command		Lorox*	
Soil Texture**	North	South	Plus	
Coarse (light) Soils: (loamy sand**, sandy loam)	1½ to 2 pints	2 to 2½ pints	9 to 12 oz.	
Medium Soits: (loam, sitt, sitt loam, sandy clay, sandy clay loam)	1½ to 2 pints	2 to 2½ pints	10 to 14 oz.	
Fine (heavy) Soils: (sifty clay, clay loam, sifty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	11 to 14 oz.	

- * Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- Plant seed at least 134 inches deep or Lorox injury to soybeans may result. Refer to Lorox Plus label for other planting and application precautions.
- ** Do not use un sand, loamy sand, gravelly soils, or exposed subsoils or soils with less than 1% organic matter. Use only on loamy sand with over 1% organic

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled

Broadleat Weeds All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following: Black Nightshade* Carpetweed Cocklebur* Giant Ragweed* Morning-glory* (Annual, Ivy-leaf, Entireleaf, Pitted, Tall)

Mustards Pigweed (Redroot, Smooth)

Partially Controlled

Large seeded weeds, germinating deep in the soil, such as morningglory, sicklepod and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a postemergence herbicide for season long control.

When applying mixtures of Command plus Lorox Plus observe all cautions and limitations appearing on the labels of each product. Observe application precautions, rotational crop/grazing and feeding restrictions and replanting instructions sections of this label when using tank mixtures.

Command 4 EC Herbicide plus Metribuzin (Sencor® or Lexone⁶)—Northern Area Önly

A tank mix combination of Command 4 EC heroicide and metribuzin will broaden the weed control spectrum compared to either product used

	Broadcast Rates Per Acre*		
Soil Texture**	C-∡mmand	Sencor DF* Lexone DF	
Coarse (light) Soils:(loamy sand**, sandy loam)	1½ to 2 pints	¼ to ⅓ lb.	
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	11/2 to 2 pints	1/4 to 1/2 lb.	
Fine (heavy) Soils: (silty clay, clay loam, silty clay luam, clay)	11/2 to 2 pints	⅓ to ½ lb.	

- Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.
- Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
Carpetweed

Pigweed* (Redroot, Smooth)
Wild Mustard

Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC herbicide plus Metribuzin (Sencor® or Lexone®)—Southern Area Only

A tank mix combination of Command 4 EC herbicide and metribuzin will broaden the weed control spectrum compared to either product used alone.

	Broadcast Rates Per Acre*	
Soil Texture**	Command	Sencor DF* Lexone DF
Coarse (light) Soils:(loamy sand**, sandy loam)	2 to 21/2 pints	1/4 to 1/3 lb.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	2 to 21/2 pints	1⁄4 to 1∕2 lb.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	2 to 21/2 pints	1/3 to 3/3 lb.

- Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- * For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.
- ** Do not use Command 4 EC in combination with metribuzin on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur. Use % to 1 pound metribuzin on silty clay and heavy clay soils of the Mississippi Delta.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Carpetweed Hemp sesbania (Coffeeweed) Pigweed (Redroot, Smooth)

Sicklepod**
Wild Mustard

Reduced Competition

* Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 4 EC plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Preview— Northern Area Only

A tank mix combination of Command 4 EC herbicide and Preview herbicide will broaden the weed control spectrum compared to either product used alone.

Broadcast Ra	atos P	or Aci	-0

Soil Texture	Command	Preview
Coarse (light) Soils:(loarny sand, sandy loarn)	1½ to 2 pints	5 to 6 oz.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1⅓ to 2 pints	6 to 7 oz.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	11/2 to 2 pints	6 to 8 oz.

Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

Do not use Command 4 EC in combination with Preview on soils with less than 0.5% organic matter, or on soils with pH greater than 6.8 or on soils having a calcareous surface area as Preview injury to rotational crops may occur.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alo. e Rates and Weeds Controlled, plus the following:

Burcucumber* Carpetweed Cocklebur*

Copperleaf, hophombeam Giant Ragweed* Morning-glory* (Annual, Tall, Ivyleaf, Pitted, Entireleaf)

Mustards Pigweed Spurge, spotted Sunflower*

NOTE: Control of morningglory, sunflower, cockdebur and giant ragweed may be erratic ranging from poor to excellent depending upon soil temperature, time of weed seed germination, depth of weed seed in the soil, and the amount and timing of soil moisture. Control may be improved with timely cultivation.

When applying mixtures of Command 4 EC plus Preview, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions or each product's label when using tank mixtures.

Command 4 EC Herbicide plus Prowl® 4 EC Herbicide— Northern & Southern Areas

A tank-mix combination of Command 4 EC and Prowl will broaden the weed control spectrum of either product used alone.

Broadcast Rates Per Acre*

Soil Texture	Com North	mand] South	Prowl 4 EC
Coarse (light) Soils: (loamy sand, sandy loam)	1½ to 2 pints	2 to 21/2 pints	1 to 1½ pints
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	2 to 2½ pints	2 pints
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	2 to 2½ pints

Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled. Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Carpetweed

Pigweed (Redroot, Smooth)

When applying mixtures of Command 4 EC plus Prowl, observe all cautions and fimitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Scepter® Herbicide— Northern Area Only

See Scepter label for appropriate geographic and rainfall restrictions.

A tank mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

NOTE: The Scepter label does not allow rotation to Com, Wheat, Barley, or Oats prior to 16 months after last Scepter application when Scepter is applied preemergent or preplant incorporated within the Northern Use Area as defined on the Scepter label. See Scepter label for appropriate geographic restrictions.

	Broadcast Hales Per Acre		
oil Texture	Command	Scepter	
I Soil Types	11/2 to 2 pints	1/3 to 1/2 pint	

Use higher use rates, within the range stated above, when heavier weed pressure is anticipated.

IEEDS CONTROLLED:

rass Weeds
If species listed under the ection Command Applied Jone Rates and Weeds controlled.

Broadleat Weeds
All species listed under the section Command Applied
Alone Rates and Weeds Controlled, plus the following:

and and Dates Day Asset

Black Nightshade Cocklebur Common Sunflower Giant Ragweed* Pigweed spp.

Partially controlled

n the event of a crop loss due to weather conditions, soybeans can be eplanted. DO NOT work the soil deeper than two inches, or man - additional Command 4 EC or Command-Scepter® tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

When applying mixtures of Command 4 EC plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Scepter Herbicide— Southern Area Only

See Scepter label for appropriate geographic and rainfall restrictions.

A tank mix combination of Command 4 EC and Scepter will broaden the weed control spectrum of either product used alone.

,	Broadcast Rates Per Acre*		
Soil Texture	Command	Scepter	
All Soil Types	2 to 21/2 pints	½ to ⅔ pint	

Use higher use rates, within the range stated above, when heavier weed pressure is anticipated.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
Black Nightshade
Cocklebur
Common Sunflower
Pigweed spp.
Wild Poinsettia

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT work the soil deeper than two inches, or make additional Command 4 EC or Command-Scepter tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

When applying mixtures of Command 4 EC plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Sonalan® EC Herbicide— Northern Area Only

A tank mix combination of Command 4 EC and Sonalan will broaden the weed control spectrum of either product used alone.

	Broadcast Rates Per Acre*		
Soil Texture	Command	Sonalan EC	
Coarse (light) Soils:(loamy sand, sandy loam)	11/2 to 2 pints	11/4 to 2 pints	
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	11/2 to 2 pints	1% to 2½ pints	
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	21/4 to 3 pints	

Select lower to higher rates within the range to correspond to the coarser to liner soil types within the textural group.

Rates within the range should correspond to anticipated weed pressure.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Carpetweed Pigweed (Redroot, Smooth)

When applying mixtures of Command 4 EC plus Sonalan, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting

Command 4 EC Herbicide plus Squadron— Southern Area Only

See Squadron label for appropriate geographic restrictions.

instructions of each product's label when using tank mixtures.

A tank mix combination of Command 4 EC and Squadron, will broaden the weed control spectrum of either product used alone.

	Broadcast Rates Per Acre		
Soil Texture	Command	Squadron*	
All Soil Types	1½ to 2 pints	11/2 to 21/4 pint	

Do not exceed 1½ pints in Northern Area (as defined on the Squadron label).
 Use higher rate of Squadron in areas of heavy cocklebur infestation.

WEEDS CONTROLLED:

Grass Weeds
All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds
All species listed under the

section Command Applied Alone Rates and Weeds Controlled, plus the following:

Black Nightshade Cocklebur Common Sunflower Giant Ragweed* Pigweed spp.

Partially controlled

When applying mixtures of Command 4 EC plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Squadron— Southern Area Only

See Squadron label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Squacron will broaden the weed control spectrum of either product used alone.

	Broadcast Rates Per Acre		
Soil Texture	Command	Squadron*	
All Soil Types	2 to 21/2 pints	21/4 to 3 pints	

^{*} Use higher rate of Squadron in areas of heavy pigweed and cocklebur infestation.

WEEDS CONTROLLED:

Grass Weeds
All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Black Nightshade Cocklebur Common Sunflower Pigweed spp. Wild Poinsettia

When applying mixtures of Command 4 EC plus Squadron, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Treflan[®] EC (Trifluralin)—Northern & Southern Areas

A tank mix combination of Command 4 EC herbicide and Treflan EC herbicide will broaden the weed control spectrum compared to either product used alone.

Broadcast Rates Per Acre*

Soil Texture	Command North South		Treflan EC
Coarse (light) Soils: (loamy sand, sandy loam)	1½ to 2 pints	2 to 2½ pints	1 pint
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ to 2 pints	2 to 2½ pints	1 to 1½ pints
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pints	2 to 2½ pints	1 to 1½ pints

Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Carpetweed Chickweed Field Bindweed Pigweed (Redroot, Smooth)

When applying mixtures of Command 4 EC plus Treflan, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Tri-Scept Herbicide— Northern Area Only

See Tri-Scept label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Tri-Scept will broaden the weed control spectrum of either product used alone.

	Broadcast Hates Per Acre	
Soil Texture	Command	Tri-Scept*
All Soil Types	1½ to 2 pints	11/s to 13/4 pints

Do not exceed 1½ pints in Northern Area (as defined on the Tri-Scept label). Use higher rates of Tri-Scept in areas of heavy cocklebur infestation.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled. **Broadleaf Weeds**

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Black Nightshade Cocklebur Common Sunflower Giant Ragweed* Pigweed spp.

Partially controlled

When applying mixtures of Command 4 EC plus Tri-Scept observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 4 EC Herbicide plus Tri-Scept—Southern Area Only

See Tri-Scept label for appropriate geographic restrictions.

A tank mix combination of Command 4 EC and Tri-Scept will broaden the weed control spectrum of either product used alone.

	Broadcast Rates Per Acre	
Soil Texture	Command	Tri-Scept*
All Soil Types	2 to 21/2 pints	1¾ to 2⅓ pints

Use higher rate of Tri-Scept in areas of heavy pigweed and cocklebur infestation.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds
All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
Black Nightshade
Cocklebur
Common Sunflower

Pigweed spp. Wild Poinsettia

When applying mixtures of Command 4 EC plus Tri-Scept observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY PREEMERGENCE TREATMENTS

Preemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of additional weed species listed on the preemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information that appear on the product labels. DO NOT apply compounds containing metribuzin after a soil incorporated Command 4 EC plus Lexone® or Sencor® tank mix treatment, unless split applications are used where the pounds of active ingredient do not exceed the total allowed.

DO NOT apply Canopy, Preview, Pursuit Plus, Scepter, Squadron, or Tri-Scept herbicides after Command 4 EC plus any of those products in a tank mix treatment, as these other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

COMMAND 4 EC HERBICIDE APPLICATION FOLLOWED BY POSTEMERGENCE TREATMENTS

Postemergence herbicides approved for use on soybeans may be applied following treatments of Command 4 EC alone or in tank mix combinations for control of emerged weeds as listed on the postemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information appearing on the product labels.

DO NOT apply Classic, Pursuit or Scepter as a postemergence treatment after Command 4 EC plus Canopy, Preview, Pursuit Plus, Scepter, Squadron, or Tri-Scept in a tank mix treatment, unless the other product labels specifically recommend such a treatment, as those other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

COMMAND 4 EC HERBICIDE FOLLOWED BY ACIFLUORFEN (BLAZER® 2L OR TACKLE®) HERBICIDE

Following an application of Command 4 EC or tank mixture with Command 4 EC, a postemergence application of acifluorien will control the following emerged broadleaf weeds:

Broadleaf Weeds
Black Nightshade
Cocklebur
Hemp sesbania
Pigweed (Redroot)
Pitted Morning-glory
Purple Moonflower
Wild Mustard

Apply acifluorien at a rate of 1 pint per acre with .25% crop oil concentrate to actively growing weeds at no more than the 4-leaf growth stage (do not count cotyledonary leaves but only the fully developed true leaves). This timing generally correlates to soybean growth stages of the first to third trifoliate leaves.

NOTE: Application of Command 4 EC herbicide generally retards the development rate of weeds which may extend the period where acifluor-ten can be effectively applied.

Read and follow all precautions, restrictions, and warnings on all product labels.

COMMAND 4 EC HERBICIDE FOLLOWED BY SCEPTER® HERBICIDE

Following an application of Command 4 EC or tank mixtures with Command 4 EC, a postemergence application of Scepter herbicide will control the following emerged broadleaf weeds:

Broadleaf Weeds
Cocklebur, common
Pigweed
Wild Poinsettia

Apply Scepter after crop emergence but before weeds exceed a height of 12 inches (6 inches for Wild Poinsettia). Apply at a broadcast rate of 1/2 pints per acre. Apply when weeds are actively growing. DO NOT apply Scepter postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes. The total amount of Scepter must not exceed 1/3 pint per acre per season in northern areas or 73 pint per acre per season in southern areas as defined on the Scepter label. Squadron and Tri-Scept both include the same active ingredient as Scepter.

For postemergence applications, the addition of a nonionic surfactant or crop oil concentrate is required. The nonionic surfactant should contain at least 80% active ingredient and should be applied at a rate of 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate stated on the COC label.

Read and follow all precautionary statements and restrictions, directions for use and all other information appearing on the Scepter label.

COMMAND® 4 EC HERBICIDE APPLICATION-CONSERVATION TILLAGE (INCLUDING MINIMUM AND NO-TILLAGE) SOYBEANS

SURFACE APPLICATIONS

Command 4 EC herbicide may be applied to the soil surface where permitted (see General Application Precautions, Geographical Restrictions) alone, or in combination with other soil residual preemergence herbicides registered for use in conservation tillage soybeans. If weeds are present at the time of treatment, include a burndown herbicide to control or suppress existing weeds. Observe all application precautions, rate restrictions, rotational guidelines and replanting instructions of each product's label when using tank mixes.

EARLY PREPLANT (EPP) APPLICATIONS

Where soil incorporated applications are required (see General Application Precautions, Geographical Restrictions), Command 4 EC herbicide may be surface applied under conservation tillage practices, but only as an early preplant (EPP) treatment. EPP applications of Command 4 EC must be made prior to field green-up (i.e., the time at which annual weeds are emerging at the treatment site, and trees or shrubs bordering the treatment is the and trees or shrubs bordering. the treatment site have broken dormancy and are showing green leaf tissue) and no later than the date listed in the table below

EPP Treatment Cutor Date Table			
	Geography	Date	
	South of I-80 (IA, IL, IN, OH, PA)	4/1	
	North of I-80 (IA, IL, IN, MI, NY, OH, PA)	4/10	
	MN WI	4/20	

When applied EPP Command 4 EC herbicide may be used at rates of 1 5 to 2.0 pints per acre (refer to RATES AND WEEDS CONTROLLED under Northern Area COMMAND 4EC HERBICIDE APPLIED ALONE) alone, or in a tank mixture with other herbicides registered for use in conservation tillage soybeans. If weeds are present at the time of treatment, tank mix with a bumdown herbicide to control or suppress existing weeds.

Observe all use directions, precautions and restrictions on the label of any herbicide(s) included in the tank mixture. When applying Command 4 EC as an EPP treatment strictly observe all restrictions and precautions found in the Preemergent Applications and Spray Drift Precautions por-tions under the General Application Precautions section of this label.

Command 4 EC Herbicide plus Canopy® or Preview® Herbicides EPP Conservation Tillage

Tank mix combination for broadening weed control spectrum.

Broadcast Rates Per Acre Soil Texture* Canopy* Command Preview** Coarse, Medium 6-8 oz. 1.5-2 pints 5-6 oz. and Fine

Select lower to higher rates within the ranges noted based on lighter to heavier soil types.

Refer to Canopy label for geographical restrictions. Do not apply to soils with less than 0.5% organic matter. Observe all cautions and limitations on the labels of each product. Special attention must be paid to sections regarding planting depth, phosphat, pesticide interpolation, previous pesticide and pest action, previous pesticides and sensitive soybean varieties on the Canopy label

Refer to Preview label for geographical restrictions. Do not use Command 4 EC in combination with Preview on soils with less than 0.5% organic matter, or on soils with pH greater than 6.8 or on soils having a calcareous surface area as Preview injury to rotational crops may occur.

WEEDS CONTROLLED: Command plus Canopy

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadle af Weeds All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following: Cocklebur** Copperleaf, hophombeam* Hemp sesbania Momingglory* (Annual, Ivyleaf, Entireleaf, Pitted, Tall) Pigweed (Redroot, Smooth)

Partially Controlled

** Large seeded weeds, germinating deep in the soil such as morninggiory, sicklered and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a posternergence herbicide for season long control.

WEEDS CONTROLLED: Command plus Preview

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

Burcucumber* Carpetweed Cocklebur*

Copperleaf, hophombeam

Giant Ragweed

Morningglory* (Annual, Tall, lyyleaf, Pitted, Entireleaf)

Mustards **Pigweed** Spurge, spotted Sunflower'

Note: Control of morningglory, sunflower, cocklebur and giant rag-weed may be erratic ranging from poor to excellent depending upon soil temperature, time of weed seed germination depth of weed seed in the soil, and the amount and timing of soil moisture. Control may be improved with timely cultivation.

When applying tank mixtures of Command 4 EC plus Canopy or Preview observe application precautions, rotational guidelines and replanting instructions of each product's label.

Command 4 EC Herbicide EPP Followed by Postemergence Applications of Pursuit® Herbicide

Following an EPP application of Command 4 EC herbicide a postemergence application of Pursuit herbicide can be made to broaden the weed control spectrum.

	Broadcast Rates Per Acre		
Soil Texture*	Command*	Pursuit*,**	
Coarse, Medium and Fine	1.5-2 pints	2-3 oz.	

Select lower to higher rates within the ranges noted based on lighter

to heavier soil types.

Application of Command 4 EC herbicide will reduce competition from the above weeds. This allows for control of light to moderate infestations with a postemergence application of Pursuit herbicide at the noted rate. Application should be made to weeds tess than one inch tall .

** Postemergence application requires specific additives. Refer to current Pursuit label for specific directions.

Command 4 EC herbicide EPP followed by Pursuit Postemergence

Continuing : Eo hairicide Ei i Tone.	ion of i gindult concluding
Weeds Controlled*	Maximum Height
Cocklebur	1"
Pigweed	1"
Nightshade	1*
- Black	
- Eastern black	
~ Hairy	
Mustard species	1*
Smartweed, Pennsylvania	1″

*In addition to the weeds controlled by Command 4 EC alone.

Erratic control may occur under heavy weed infestations. Control may be improved by cultivating approximately 14 days after the postemergence

DO NOT apply Pursuit herbicide with ground equipment when wind velocity is greater than 10 mph, or when spray may be carried to sensitive crops. Sensitive crops include leafy vegetables, sugar beets and cotton.

Refer to the General Application Instructions portion of this label and to the Pursuit herbicide label for specific application instructions. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when applying each product.

DRILL 'N DRAG APPLICATIONS

For conservation tillage use of Command 4 EC herbicide where soil incorporated applications are required, and the timing for EPP treatments has passed Command 4 EC may be applied using no-tilt drills capable of incorporating Command 4 EC (i.e., "Drill 'N Drag" type equipment). The following conditions apply:

- 1) Drills which result in incorporation, including Great Plains® Drills, Tye® Drills and Crustbuster® Drills which are equipped with fluted or rippled coulters and spring tines or harrows may be used. John Deere® Drills may be used but must employ spring tines or the equivalent to enhance incorporation Drills should be operated at speeds greater than 6 mph to ensure enough soil is moved to incorporate Command 4 EC. Other minimum-till/no-till drill equipment may be used for "Drill and Drag" application of Command 4 EC herbicide as long as their operation results in soil mixing.
- 2) When utilizing the "Drill and Drag" technique DO NOT APPLY Command 4 EC herbicide within 1,000 feet of subdivisions and towns, commercial vegetable production (except sweet corn) commercial fruit production commercial nurseries and commercial greenhouses. Caution must be taken to minimize spray drift as off-site movement of Command herbicide can cause foliar whitening or yellowing of some plants (refer to plant listing, and Spray Drift Precautions under the General Application Precautions section of this label).
- 3) Where soybean planting and Command 4 EC application using the "Drill and Drag" technique is not a one-pass operation, drilling of soybean seeds must occur immediately after Command 4 EC application on wet soils and within 8 hours on dry soils.
- 4) All other cautions and precautionary statements (sprayer cleanup, use rates, rotational cropping precautions and restrictions, etc.) on the Command 4 EC label must be followed.

PEPPERS

GENERAL INFORMATION

Command 4 EC herbicide may be utilized as a soil applied treatment prior to weed emergence, for the control of annual grass and broadleaf weeds in bell, hot, pimento and sweet peppers. Do not use on banana peppers.

Make a single herbicide application in a minimum of 10 gallons of water per acre at the rate shown in the application rate table. Apply to the soil surface prior to seeding or transplanting of peppers and incorporate to a depth of 1-inch or less. Place seed, or roots of the transplants, below the chemical barrier when planting.

Command 4 EC must be soil incorporated in the following states unless an approved 24(c) permitting otherwise is on file within that state (check with your local dealer): Connecticut, Delaware, Illinois, Indiana, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, and Wisconsin.

Command 4 EC herbicide may be tank mixed with other herbicides registered for use in peppers to broaden the weed control spectrum compared to the products applied alone. When applying mixtures of Command 4 EC plus other bell and non-bell pepper herbicides, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational muidelines and replanting instructions of each product's label when using tank mixes. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with other pepper herbicides unless use directions specifically state otherwise.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

REPLANTING INSTRUCTIONS

If initial seeding or transplants fail to produce a uniform stand, peppers may be replanted in a field treated with Command 4 EC alone. Do not retreat field with a second application of Command 4 EC. When tank mixing with a labeled product, refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Guidelines on the Command label, Where a tank mix is used refer to the product(s) labels for any additional replant instructions.

NOTE: Seed or transplants should be planted below the herbicide barrier to minimize early bleaching of seedlings. Not all varieties have been tested for tolerance to a Command treatment. Pepper type and variety, soil type and amount of organic matter rate of application depth of incorporation, planting depth, amount of moisture throughout the growing season and other factors can impact the degree of sensitivity that may be observed.

COMMAND 4 EC HERBICIDE APPLIED ALONE RATES AND WEEDS CONTROLLED

Soil Texture	Broadcast Rates Per Acre* 0.5 to 0.75 pint	
Coarse (light) Soils: (loamy sand, sandy loam)		
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	0.75 to 1.0 pint	
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1.0 to 2.0 pints	

^{*}Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

WEEDS CONTROLLED:

WEEDS CONTROLLED.		
RATE/ACRE 0.5-0.75 pint (0.25-0.38 LB. A.I.)	RATE/ACRE 0.75-1.0 pint (0.38-0.5 LB, A.I.)	RATE/ACRE 1.0-2.0 pints (0.5-1.0 LB. A.I.)
GRASSES Barnyardgrass* Crabgrass* —Large —Smooth Foxtail* —Giant —Green —Robust Goosegrass* Panicum* —Common —Fall	GRASSES Bamyardgrass* Crabgrass* —Large —Smooth Field Sandbur* Foxtail —Giant —Green —Robust Goosegrass Panicum —Common —Fall Seeding Johnsongrass*	GRASSES Barnyardgrass Crabgrass —Large —Smooth Field Sandbur Foxtail —Giant —Green —Robust —Yellow Goosegrass Panicum —Common —Fall Seedling Johnsongrass*
BROADLEAVES Galinsoga spp.* Lambsquarters* Spurred Anoda* Velvetleaf*	BROADLEAVES Galinsoga spp." Lambsquarters" Prickly Sida" Spurred Anoda Velvetleaf Venice Mallow"	BROADLEAVES Velvetleaf Spurred Anoda Common Ragweed* Galinsoga Jimsonweed* Lambsquarters Prickly Sida Purslane Smartweed* Venice Mallow

^{*} Partialty Controlled

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed under Soybean Rotational Crop Guidelines, otherwise crop injury may occur.

Cover crops may be planted at anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment Do not allow livestock to graze on treated peppers or feed treated pepper trash to livestock.

PUMPKINS

GENERAL INFORMATION

Command 4 EC herbicide may be utilized as a preemergent soil applied treatment for the control of annual grass and broadleaf weeds in pumpkins.

Command 4 EC must be soil incorporated in the following states unless an approved 24(C) is on file for counties within that state (check with your local dealer): Connecticut, Delaware, Illinois, Indiana, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, and Wisconsin.

NOTE: When utilizing Command 4 EC as a soil incorporated treatment, incorporate Command to a depth of 1 inch or less and place the seed below the chemical barrier when planting.

Command 4 EC herbicide may be tank mixed with other pumpkin herbicides to broaden the weed control spectrum compared to the products applied alone. When applying mixtures of Command 4 EC herbicide plus other pumpkin herbicides, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with the herbicides listed above unless use directions specifically state otherwise.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

REPLANTING INSTRUCTIONS

If initial seeding of pumpkins fails to produce a stand, pumpkins may be replanted in fields treated with Command 4 EC alone. Do not retreat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the pumpkin replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Guidelines on this label.

COMMAND 4 EC HERBICIDE APPLIED ALONE RATES AND WEEDS CONTROLLED

Do not exceed a rate of 2 pints per acre. Weeds controlled (by rate) are as listed under Soybean for Northern and Southern areas.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed under Soybean Rotational Crop Guidelines, otherwise crop injury may occur.

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated pumpkin vines or feed treated vines or vine trash to livestock.

SUCCULENT PEAS

GENERAL INFORMATION

Command 4 EC herbicide may be utilized as preemergent soil applied treatment for the control of annual grass and broadleaf weeds in peas.

Command 4 EC must be soil incorporated in the following states unless an approved 24(C) is on file for counties within that state (check with your local dealer): Connecticut, Delaware, Illinois, Indiana, Iowa, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, and Wisconsin.

NOTE: When utilizing Command 4 EC as a soil incorporated treatment, incorporate Command to a depth of 1 inch or less and place the seed below the chemical barrier when planting.

Command 4 EC herbicide may be tank mixed with other herbicides registered for use in peas to broaden the weed control spectrum compared to the products applied alone. When applying mixtures of Command 4 EC plus other succulent pea herbicides, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions rotational guidelines and replanting instructions of each product's label when using tank mixes. Water or liquid fertilizer may be used as a carrier for Command 4 EC when applied alone, or when tank mixed with other succulent pea herbicides unless use directions specifically state otherwise.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command® 4 EC her-bicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

REPLANTING INSTRUCTIONS

If initial seeding of peas fails to produce a stand, peas may be replanted in fields treated with Command 4 EC alone (or with recommanded tank mixtures). Do not retreat field with a second application of Command 4 EC. When tank mixing with a labeled product refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Guidelines on this label.

COMMAND 4 EC HERBICIDE APPLIED ALONE

FOR PEAS ONLY: Apply only at 1 pint (0.5 Lb. A.I.) Per Acre

RATE AND WEEDS CONTROLLED

1 PT. (0.5 LB. A.I.) (for all soil types)

GRASSES
Barnyardgrass*
Crabgrass (Large, Smooth)
Field Sandbur*
Foxtail (Green, Giant,
Robust)
Goosegrass
Panicum (Fall, Common)
Seedling Johnsongrass*

BROADLEAVES
Galinsoga spp*
Lambsquarters*
Prickly Sida*
Spurred Anoda
Velvetleaf
Venice Mallow*

*Partially Controlled

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed under Soybean Rotational Crop Guidelines, otherwise crop injury may occur.

Cover crops may be planted at anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 4 EC treatment. Do not allow livestock to graze on treated pea vines or feed treated vines or vine trach to livestock.

FALLOW LAND

GENERAL INFORMATION

Command[®] 4 EC is a herbicide which may be utilized as a preemergence surface applied treatment to control many annual grass and broadleaf weeds on fallow land. Command 4 EC may be tank mixed with other herbicides registered for fallow cropland use (e.g., atrazine) to broaden the weed control spectrum compared to the products applied alone.

SPECIAL PRECAUTION >

Off-site movement of spray drift or vapors of Command 4 EC herbicide can cause toliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

Additional Precaution For Fallow Land Applications Only: Do not apply within 500 feet of Emerged Winter Wheat or within 1,000 feet of the areas listed below:

Towns and Subdivisions
Commercial Vegetable Production*
Commercial Fruit Production
Commercial Nursenes
Commercial Greenhouses
*except sweet com

Supplemental Spray Drift Precautions For Fallow Land Applications: Non-target spray drift of Command 4 EC herbicide should be avoided to prevent whitening of desirable vegetation. Specific recommendations and requirements for minimizing spray drift are included in the Application Precautions section of this label. For Fallow Land applications only:

 The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 5 to 15 gallons per acre and is recommended for application volumes of 15 to 40 gallons per acre when spraying in the proximity of desirable plants (see list under Application Precautions).

ROTATIONAL CROP/GRAZING AND FEEDING RESTRICTIONS: Do not plant wheat sooner than nine (9) months after a late summer or fall application. Do not allow livestock to graze on treated fields.

APPLICATION RATES:

Use Command 4 EC herbicide alone at a rate of 1½ to 2 pints (24 to 32 ounces) per acre in a surface applied broadcast application for preemergent control of weeds. Where Glean is registered and was applied at ½ to ½ ounce per acre for weed control in the growing wheat crop, a rate of 1½ pints (24 ounces) per acre of Command 4 EC is recommended. However, the additional weed control benefit from Glean may not last throughout the following spring (see Glean label). Use the higher rate of Command 4 EC when heavier weed infestations are anticipated or for longer residual control. practiced, make application after wheat harvest. For optimum control, make applications after August 15 but before October 31. Command 4 EC can also be used in areas where spring wheat is planted following an 18 month fallow period.

Treatments should be made only to fields where good weed control has been maintained in the wheat crop through wheat harvest. Treatment recommendations are only designed to control weeds that have recently emerged and those that germinate throughout the fall, winter and spring period. Do not apply in situations where dense weed growth exists.

WEEDS CONTROLLED:

Grass Weeds
Cheat Grass
Downy Bromegrass
Jointed Goatgrass
Volunteer Bye
Volunteer Rye

Broadleaf Weeds
Common Lambsquarters
Common Purslane
Kochia
Prickly Lettuce
Tansy Mustard
Tumble Mustard
Wild Buckwheat

COMMAND 4 EC APPLIED IN COMBINATIONS

Tank mix combinations of Command 4 EC herbicide with atrazine will broaden the weed control spectrum and length of control compared to the herbicides used alone.

Command 4 EC Plus Atrazine

Command 4 EC at 1 to 1½ pints (16 to 24 ounces) per acre plus 1 to 2 pints of Atrazine 4L or equivalent rates of Atrazine 90 DF and 80 W. Weeds controlled in addition to those noted under Command 4 EC alone will depend on atrazine rate used (see atrazine label). Use the higher rate of Command 4 EC when heavier weed infestations are anticipated or for longer residual control.

There is significant risk of atrazine carryover if not used according to local Experiment Station and label recommendations. Consult your local Experiment Station for recommended arrazine rates for use in fallow in your area.

Refer to the atrazine label for planting, cropping and other restrictions. Follow all precautions, statements, and restrictions on this and the atrazine label.

Mixing Instructions:

Fill spray tank ½ to ½ full with water and add recommended amount of atrazine with agitator operating. Mix thoroughly and fill tank ½ full continuing agitation. Add Command 4 EC herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Maintain agitation during mixing and application.

Application Instructions:

Follow the General Application Instructions as for Command 4 EC alone.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/o; handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

COMMAND and FMC—Trademarks of FMC Corporation
CANOPY, GLEAN, LEXONE, LOROX, LOROX PLUS and PREVIEW—
Trademarks of E.I. duPont de Nemours and Co., Inc.
OPTION—Trademark of Hoechst-Roussel Agri-Vet Company
SONALAN, TREFLAN—Trademarks of DowElanco
LASSO and ROUNDUP—Trademarks of Monsanto Company
SENCOR—Trademark of Bayer AG
PROWL, SCEPTER, SQUADRON, TRI-SCEPT—Trademarks of Amorican Cyanamid Company
DUAL—Trademark of Ciba-Geigy Corporation
BLAZER—Trademark of BASF Corporation
TACKLE—Trademark of Rhone-Poulenc, Inc.
C1987 FMC Corporation
All rights reserved

All rights reserved (618-1/14/94-A)