

Code 619

Net Contents

Command® 6 EC

Herbicide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-3054

EPA Est. 279-


Active Ingredient:	By Wt.
2-(2-Chlorophenyl)methyl-4,4-dimethyl-3-isoxazolidinone.....	64.3%
Inert Ingredients:	35.7%
	<u>100.0%</u>

Contains 6 pounds of active ingredient per gallon
U.S. Patent No. 4,405,357

**KEEP OUT OF REACH OF CHILDREN
DANGER**

FIRST AID

- If in eyes:** Flush immediately with plenty of water. Get medical attention.
 - If swallowed:** Drink promptly large quantities of milk or water. Avoid alcohol. Do not induce vomiting. Call a physician.
 - If inhaled:** Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
 - If on skin:** Remove contaminated clothing. Wash skin with plenty of soap and water.
- See other panels for additional precautionary information.
For Emergency Assistance Call: (716) 735-3765



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

2/86

PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

Danger
Causes persistent but reversible eye injury. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes or on clothing. Avoid contact with skin. When measuring or mixing, wear goggles or face shield, rubber gloves and full arm and leg length clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS
Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Pesticide Storage
Do not freeze. Do not store below 40°F. If solid crystals are observed, warm material to about 60°F by placing container in warm location. Shake or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC Collect: (716) 735-3765.

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal
Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal
Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.
Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

GENERAL INFORMATION
Command 6 EC selective herbicide may be utilized as a preemergence surface applied or preplant incorporated treatment for the control of many annual grass and broadleaf weeds in soybeans.

Command 6 EC herbicide may be tank mixed with other soybean herbicides to broaden the weed control spectrum compared to the products applied alone. Command 6 EC may be tank mixed with Sencor® or Lexone® herbicides and applied preemergence or preplant incorporated. It may be applied preemergence in tank mixture with Lorox® or Amiben® herbicides. Command 6 EC alone or tank mixed with Sencor, Lexone, Lorox or Amiben, plus either paraquat or Roundup® herbicide may be used for weed control in minimum-tillage or no-tillage soybeans.

Water or liquid fertilizer may be used as a carrier for Command 6 EC when applied alone, or when tank mixed with Sencor, Lexone, Lorox or Amiben unless use directions specifically state otherwise.

IMPORTANT

Injury to desirable plants and/or loss of crops may result from failure to observe the following:

Do not plant wheat, oats, barley, rye or alfalfa in the fall of the year of application or in the spring of the following year as crop injury may occur (See General Application Instructions—Rotational Crop section).

Some desirable plants, including ornamentals (e.g., roses), trees (e.g., flowering and edible cherries), agronomic crops (e.g., small grains, alfalfa, sunflowers) and vegetables (e.g., lettuce, cole crops, radish) are sensitive to Command herbicide. Foliar contact with spray drift or vapors may cause visual symptoms of chlorosis or bleaching to sensitive plants growing in the proximity of treated fields. These symptoms are temporary. Extra caution is advised when spraying near desirable plants found in residential or other commercial agricultural plantings. (See General Application Instructions section.)

Do not drain or flush equipment on or near desirable trees or other plants, or in areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. Do not contaminate any body of water including irrigation water that may be used on other crops. Carefully follow sprayer clean-up instructions noted below to prevent spray tank residues from damaging other crops.

Spray equipment should be thoroughly cleaned to remove all traces of herbicide that might injure other subsequently sprayed crops. Drain any remaining spray solution from tank and discard in non-crop areas away from water supplies. Wash tank, pump, hoses and boom with a suitable solution of heavy-duty detergent. Recycle this mixture through the equipment for 5 minutes and spray out in non-crop area away from water supplies. Repeat this procedure twice. Rinse tank, pump, hoses and boom with several changes of water. Remove nozzle tips and screens and clean separately if tank mixtures with wettable powders were utilized. Wash away any spray mixture from the outside of spray tank or other portions of the spray rig.

GENERAL MIXING INSTRUCTIONS

Command 6 EC Alone: Mix Command 6 EC herbicide with water or liquid fertilizer in the following manner: Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 6 EC, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing to maintain a uniform spray mixture.

Tank Mixtures: Fill spray tank one-fourth to one-third full with water; with agitator operating add the recommended amount of ingredients using the following order: dry formulations (e.g., wettable powder, dry granules) first, liquid suspensions (e.g., flowables) next. Mix thoroughly and fill tank one-half full continuing agitation. Add Command 6 EC herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Maintain agitation during filling, mixing and application. If paraquat or Roundup[®] is used in the tank mix, add the required amount of either product prior to, but near the end of the filling process. Where use of a surfactant is recommended, add as the last ingredient to the spray tank.

Fertilizer Spray Mixtures: Application of Command 6 EC alone, or in combination with either Sencor[®], Lexone[®], Lorox[®], Amiben[®], paraquat or Roundup[®] herbicide, in conjunction with fertilizer solutions may be used unless otherwise directed. Small quantities should be tested for compatibility by the following procedure before mixing tankful quantities:

- 1) Put 1 pint of fertilizer solution in a quart jar.
- 2) Add the appropriate amount of herbicide based on the tables below. If more than one product is to be used, add each separately using the following sequence: wettable powders or dry flowables first, flowable liquids second and liquids (e.g., EC's) last.

Herbicide	Rate/Acre	Amount To be Added Per Pint of Fluid Fertilizer*
Wettable Powders or Dry Flowables (Dry Granules)	1/2 pound	3/4 level teaspoon
	1 pound	1 1/2 level teaspoons
	2 pounds	3 level teaspoons
	3 pounds	4 1/2 level teaspoons
Emulsifiable Concentrates or Flowables	1 pint	1/2 level teaspoon
	1 quart	1 level teaspoon
	2 quarts	2 level teaspoons
	3 quarts	3 level teaspoons

*Based on spray volume of 25 gpa. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

- 3) Close jar and shake well.
- 4) Watch mixture for several seconds, again in 5 minutes and again after 30 minutes.
- 5) If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed.

If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below:

- 1) **Command 6 EC Alone:** Dilute with 2 parts of water and add slowly to spray tank of fertilizer with agitator running. Maintain agitation during application.
- 2) **Command 6 EC plus Tank Mixtures:** Dilute individual products with two (2) parts of water, then add to the spray tank of fertilizer, while maintaining agitation, using the following order—slurry of dry formulations (wetable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Paraquat or Roundup[®] should be added as the last ingredient. Continue agitation during application.

GENERAL APPLICATION INSTRUCTIONS

Ground Applications

Broadcast Application: Apply Command 6 EC alone or in tank mix combinations by ground equipment using a finished spray volume of 10 to 40 gallons of water per acre. Use nozzles suitable for broadcast boom application of herbicides. For Command 6 EC tank mixtures with wettable powder or dry flowable formulations, nozzle screens and strainers should be no finer than 50-mesh. Over-application through excessive overlapping of spray swaths or failing to shut off spray booms when turning (end row areas), slowing or stopping can result in undesirable soil residues.

Band Application: Use nozzles suitable for banded application of herbicides. Calculate the rates and volumes required by using the formulas below:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast rate per acre}}{\text{per acre}} = \frac{\text{Band rate per acre}}{\text{per acre}}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \frac{\text{Broadcast volume per acre}}{\text{per acre}} = \frac{\text{Band volume per acre}}{\text{per acre}}$$

APPLICATION PRECAUTIONS

- (1) Care should be taken to minimize spray drift when applying Command herbicide.
 - Do not apply when weather conditions favor drift.
 - Use nozzles which produce coarser sprays.
 - Use lower pressures to reduce production of fine spray droplets.
 - The use of agriculturally approved drift reducing additives, particularly when spraying in the proximity of desirable plants is recommended.
- (2) In the proximity of desirable plants use soil incorporation to reduce vapor movement from treated fields.
- (3) Improper application of Command herbicide may result in injury to rotational crops. Particular attention should be paid to the following:
 - Do not exceed label recommended rates.
 - Avoid over-application resulting through use of worn nozzles, excessive overlapping of spray swaths, failing to shut off spray booms when turning (end row areas) or slowing or stopping sprayer.

ROTATIONAL CROP/GRAZING AND FEEDING RESTRICTIONS

Do not rotate to wheat, oats, barley, rye or alfalfa in the fall of the year of application or in the spring of the following year as crop injury may occur. Other rotational crops may be planted nine (9) months after application. Cover crops may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest these cover crops for food or feed. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

REPLANTING INSTRUCTIONS

If initial seeding of soybeans fails to produce a stand, soybeans may be replanted in fields treated with Command 6 EC alone (or with recommended tank mixtures). Do not retreat field with a second application of Command 6 EC alone (or with recommended tank mixtures). Do not replant treated fields with any crop (except soybeans) at intervals which are inconsistent with the above Rotational Crop statement.

COMMAND 6 EC HERBICIDE APPLIED ALONE
WEEDS CONTROLLED:

<u>Grass Weeds</u>	<u>Broadleaf Weeds</u>
Barnyardgrass	Cocklebur*
Broadleaf signalgrass (Brachiaria)	Florida pusley
Crabgrass	Florida beggarweed
(Large crabgrass)	Jimsonweed
(Smooth crabgrass)	Lambsquarters
Field sandbur	Pitted morningglory
Foxtail	Prickly sida
(Green foxtail)	Common purslane
(Giant foxtail)	Ragweed
(Yellow foxtail)	(Common ragweed)
(Robust purple foxtail)	Smartweed
Goosegrass	(Pennsylvania smartweed)
Johnsongrass (Seedling)	Spurge (Prostrate)
Panicum	Tropic croton
(Fall panicum)	Velvetleaf
(Texas panicum)	Venice mallow
Southwestern cupgrass	
Woolly cupgrass	

WEEDS PARTIALLY CONTROLLED:

<u>Grass Weeds</u>	<u>Broadleaf Weeds</u>
Bermudagrass	Black nightshade
Red rice	Pigweed
Shattercane	(Redroot pigweed)
Wild proso millet	(Smooth pigweed)
	Redvine

*Control in southern states (including AL, AR, FL, GA, LA, KY, MS, TN, VA and W. VA) and northern states (including DE, MD, MI, NJ, NY, OH and PA). Partial control (suppression) in midwest states (including CO, IA, IL, IN, KS, MN, MO, NE, ND and SD) and in NC and SC. Where suppression is claimed, control may be improved with timely cultivation.

SOYBEANS

Preemergence Broadcast Application: Use Table 1 to select the recommended rate of Command 6 EC herbicide based on soil texture and percent organic matter. Apply as a broadcast spray during planting, or as a separate operation after planting but before weeds or soybeans emerge.

(NOTE: Command 6 EC herbicide when used alone may be applied through "cracking stage" of soybeans. When applied at cracking stage temporary yellowing of cotyledonary leaves may occur.)

Table 1: Command 6 EC Alone

Preemergence Broadcast Rates Per Acre (Pints)

Soil Texture*	Less than 3% Organic Matter	3% or Greater Organic Matter
Coarse (light) Soils: (sand, loamy sand, sandy loam)	1 to 1 1/2 pts. or 16 to 21.3 fl. oz.	1 1/2 pts. or 21.3 fl. oz.
Medium Soils: (loam, silt loam, silt, sandy clay, sandy clay loam)	1 1/2 pts. or 21.3 fl. oz.	1 1/2 to 1 3/4 pts. or 21.3 to 26.7 fl. oz.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1 1/2 to 1 3/4 pts. or 21.3 to 26.7 fl. oz.	1 1/2 to 1 3/4 pts. or 21.3 to 26.7 fl. oz.

*Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., sand, loamy sand, sandy loam) within a textural group, and increasing organic matter levels.

Band Application: Apply Command 6 EC herbicide at a broadcast equivalent rate and volume per acre. Use the rates from the Preemergence Broadcast Application directions above (Table 1) and adjust rate and total spray volumes by using the formulas in the General Application Instructions portion of this label.

Preplant Incorporated Application: Use Table 2 to select the recommended rate of Command 6 EC herbicide. Apply Command 6 EC to the soil surface and uniformly incorporate. Select equipment suited accordingly for shallow or deeper incorporation. Equipment should be set to incorporate Command 6 EC herbicide (or tank mixes) to a depth of 3 inches or less in the final seedbed.

Table 2: Command 6 EC Alone
Preplant Incorporated Rates Per Acre (Pints)

Soil Texture*	Less than 3% Organic Matter	3% or Greater Organic Matter
Course (light) Soils: (sand, loamy sand, sandy loam)	1 to 1 1/2 pts. or 16 to 21.3 fl. oz.	1 1/2 to 1 3/4 pts. or 21 to 26.7 fl. oz.
Medium Soils: (loam, silt loam, silt, sandy clay, sandy clay loam)	1 1/2 to 1 3/4 pts. or 21.3 to 26.7 fl. oz.	
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1 1/2 to 1 3/4 pts. or 21.3 to 26.7 fl. oz.	

*Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., sand, loamy sand, sandy loam) within a textural group, and increasing organic matter levels.

COMMAND 6 EC APPLIED IN COMBINATIONS

COMMAND 6 EC herbicide Plus Sencor* or Lexone*
Tank mixed combinations of Command 6 EC herbicide and Sencor or Lexone herbicides will control more broadleaf weeds than either herbicide used alone. Command 6 EC herbicide broadleaf activity complements Sencor or Lexone, thereby permitting a reduction in the Sencor or Lexone use rate for broad spectrum control of many weed species, and a widening of the margin of safety to soybeans.

COMMAND 6 EC HERBICIDE PLUS SENCOR/LEXONE "REDUCED" RATE TANK MIX

WEEDS CONTROLLED:

<u>Grass Weeds</u>	<u>Broadleaf Weeds</u>
All species listed under "Command 6 EC Herbicide Applied Alone" section of the label.	All species listed under "Command 6 EC Herbicide Applied Alone" section of the label plus the following:
	Carpetweed
	Pigweed
	(Redroot pigweed)
	(Smooth pigweed)
	Spurred anoda
	Wild mustard

WEEDS PARTIALLY CONTROLLED:

<u>Grass Weeds</u>	<u>Broadleaf Weeds</u>
All species listed under Command 6 EC Herbicide Applied Alone" section of the label.	Black nightshade
	Redvine

Table 3: Command 6 EC Plus Sencor/Lexone "Reduced" Rates

Soil Texture**	Broadcast Rates Per Acre			
	Less Than 3% Organic Matter		3% or Greater Organic Matter	
	Command 6 EC	Sencor DF* or Lexone DF	Command 6 EC	Sencor DF* or Lexone DF
Coarse (light) Soils: (loamy sand**, sandy loam)	1 pt. or 15 fl. oz.	1/4 lb.	1 1/2 pts. or 21.3 fl. oz.	1/4 lb.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1 1/2 pts. or 21.3 fl. oz.	1/4 lb.	1 1/2 pts. or 21.3 fl. oz.	1/4 lb.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1 1/2 pts. or 21.3 fl. oz.	1/4 lb.	1 1/2 pts. or 21.3 fl. oz.	1/4 lb.

*For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.

**Do not use Command 6 EC in combination with Sencor or Lexone on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter as Sencor/Lexone injury to soybeans may occur. Use 1/2 to 1.0 pound Sencor or Lexone DF per acre in this tank mixture when applying on silty clay and heavy clay soils of the Mississippi Delta.

When applying mixtures of Command plus Sencor or Lexone observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop/Grazing and Feeding Restrictions and Replanting Instructions sections of this label when using tank mixtures.

COMMAND 6 EC HERBICIDE PLUS "FULL" RATE SENCOR[®]/LEXONE[®] TANK MIX

WEEDS CONTROLLED:

Grass Weeds

All species listed under "Command 6 EC Alone" and Command 6 EC Plus "Reduced" Rate section of label.

Broadleaf Weeds

All species listed under "Command 6 EC Alone" and Command 6 EC Plus "Reduced" Rate section of label, plus the following:
Hemp sesbania (coffee weed)

WEEDS PARTIALLY CONTROLLED:

Grass Weeds

All species listed under "Command 6 EC Alone" and Command 6 EC Plus "Reduced" Rate section of label.

Broadleaf Weeds

Black nightshade
Redvine
Sicklepod

Table 4: Command 6 EC Plus "Full" Rates of Sencor or Lexone

Broadcast Rates Per Acre

Soil Texture**	Less Than 3% Organic Matter		3% or Greater Organic Matter	
	Command 6 EC	Sencor DF* or Lexone DF	Command 6 EC	Sencor DF* or Lexone DF
Coarse (light) Soils: (loamy sand**, sandy loam)	1 pt. or 16 fl. oz.	1/2 lb.	1 1/2 pts. or 21.3 fl. oz.	1/2 lb.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1 1/2 pts. or 21.3 fl. oz.	1/2 lb.	1 1/2 pts. or 21.3 fl. oz.	3/4 lb.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1 1/2 pts. or 21.3 fl. oz.	3/4 lb.	1 1/2 pts. or 21.3 fl. oz.	3/4 lb.

*For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.

**Do not use Command 6 EC in combination with Sencor or Lexone on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter or on soils having a calcareous surface area or a pH of 7.5 or higher as Sencor/Lexone injury to soybeans may occur. Use 1 to 1 1/2 pounds Sencor or Lexone DF in this tank mixture when applying on silty clay and heavy clay soils of the Mississippi Delta.

When applying mixtures of Command plus Sencor or Lexone observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop/Grazing and Feeding Restrictions and Replanting Instructions sections of this label when using tank mixtures.

MIXING INSTRUCTIONS

Fill spray tank 1/4 to 1/2 full with water and add recommended amount of Sencor/Lexone with agitator operating. Mix thoroughly and fill tank 1/2 full continuing agitation. Add Command 6 EC herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Maintain agitation during mixing and application.

APPLICATION INSTRUCTIONS

Preemergence Application: Apply as a broadcast or banded spray during planting, or as a separate operation after planting but before weeds or soybeans emerge. Utilize nozzles suited for broadcast boom application or banded application accordingly. Refer to the General Application Instructions portion of this label for appropriate rate and volume of application adjustment formulas for banded treatment.

Preplant Incorporated Application: Apply tank mix to the soil surface and uniformly incorporate. Select equipment suited accordingly for shallow or deeper incorporation. Equipment should be set to incorporate the tank mix to a depth of 3 inches or less in the final seedbed.

COMMAND 6 EC HERBICIDE PLUS LOROX[®]

A preemergence tank mix combination of Command 6 EC herbicide and Lorox herbicide will broaden the weed control spectrum compared to either product alone.

WEEDS CONTROLLED:

Grass Weeds

All species listed under "Command 6 EC Herbicide Applied Alone" section of the label.

Broadleaf Weeds

All species listed under the "Command 6 EC Herbicide Applied Alone" section of the label plus the following:

- Carpetweed
- Chickweed
- Common day flower
- Galinisoga
- Pigweed (Redroot pigweed) (Smooth pigweed)
- Wild mustard
- Wild radish

WEEDS PARTIALLY CONTROLLED:

Grass Weeds

All species listed under "Command 6 EC Herbicide Applied Alone" section of the label.

Broadleaf Weeds

Black nightshade
Redvine

Table 5: Command 6 EC Plus Lorox

Broadcast Rates Per Acre

Soil Texture**	Less Than 3% Organic Matter		3% or Greater Organic Matter	
	Command 6 EC	Lorox DF or 50 WP*	Command 6 EC	Lorox DF or 50 WP*
Coarse (light) Soils: (loamy sand**, sandy loam)	1 pt. or 16 fl. oz.	2/3 to 1 lb.	1 1/2 pts. or 21.3 fl. oz.	1 to 1 1/2 lbs.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	1 1/2 pts. or 21.3 fl. oz.	1 to 1 1/2 lbs.	1 1/2 pts. or 21.3 fl. oz.	1 1/2 lbs.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1 1/2 pts. or 21.3 fl. oz.	1 1/4 to 2 lbs.	1 1/2 pts. or 21.3 fl. oz.	2 to 3 lbs.

*When using Lorox L or Lorox DF, use equivalent rates; one pint of Lorox L equals one pound of Lorox DF or Lorox 50 WP. Plant seed at least 1 1/4 inches deep or Lorox injury to soybeans may result. Refer to Lorox label for other planting and application precautions.

**Do not use on sand, gravelly soils, or exposed subsoils or soils with less than 1% organic matter. Use only on loamy sand with over 1% organic matter.

When applying mixtures of Command plus Lorox observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop/Grazing and Feeding Restrictions and Replanting Instructions sections of this label when using tank mixtures.

MIXING INSTRUCTIONS

Refer to mixing instructions as noted for Sencor/Lexone above.

APPLICATION INSTRUCTIONS

Preemergence Application: Refer to preemergence application instructions as noted for Sencor/Lexone above.

COMMAND 6 EC HERBICIDE PLUS AMIBEN[®]

A preemergence tank mix combination of Command 6 EC and Amiben herbicides will broaden the weed control spectrum compared to either product alone.

WEEDS CONTROLLED:

Grass Weeds

All species listed under "Command 6 EC Herbicide Applied Alone" section of the label.

Broadleaf Weeds

All species listed under "Command 6 EC Herbicide Applied Alone" section of the label plus the following:

- Black nightshade
- Carpetweed
- Common chickweed
- Pigweed (Redroot pigweed) (Smooth pigweed)
- Wild mustard

PARTIAL CONTROL:

Grass Weeds

All species listed under "Command 6 EC Herbicide Applied Alone" section of the label.

Broadleaf Weeds

Redvine

Table 6: Command 6 EC Plus Amiben*
Broadcast Rates Per Acre

Soil Texture**	Less Than 3% Organic Matter		3% or Greater Organic Matter	
	Command 6 EC	Amiben Liquid (Amiben DS)	Command 6 EC	Amiben Liquid (Amiben DS)
Coarse (light) Soils (loamy sand**, sandy loam)	1 pt or 16 fl oz	4 qts (2½ lbs)	1½ pts or 21 3 fl oz	4 qts (2½ lbs)
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1½ pts or 21 3 fl oz	4 to 5 qts (2½ to 3½ lbs)	1½ pts or 21 3 fl oz	4 to 5 qts (2½ to 3½ lbs)
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ pts or 21 3 fl oz	4 to 5 qts (2½ to 3½ lbs)	1½ pts or 21 3 fl oz	5 to 6 qts (3½ to 4 lbs)

When applying mixtures of Command plus Amiben observe all cautions and limitations appearing on the labels of each product. Observe Application Precautions, Rotational Crop/Grazing and Feeding Restrictions and Replanting Instructions sections of this label when using tank mixtures.

MIXING INSTRUCTIONS

Refer to mixing instructions as noted under Sencor*, Lexone* section of label.

APPLICATION INSTRUCTIONS

Preemergence Refer to preemergence application instructions as noted Application: under Sencor/Lexone section of label.

COMMAND 6 EC HERBICIDE APPLICATION—MINIMUM TILLAGE OR NO-TILLAGE SOYBEANS

Command 6 EC herbicide may be used alone, or in combination with Sencor*, Lexone*, Lorox*, or Amiben* herbicides plus paraquat or Roundup* herbicide for preemergence (to crop and weeds) and post-emergence (to weeds, but prior to crop emergence) control of broadleaf and grass weeds where soybeans will be planted into a cover crop or in previous crop residues (e.g., corn, small grain stubble).

WEEDS CONTROLLED: When used as directed, the paraquat portion of the tank mix controls most emerged annual weeds and suppresses many perennial weeds. Roundup combinations will control emerged annual and perennial weeds when applied as directed on the Roundup label. Command 6 EC or Command 6 EC plus Sencor, Lexone or Lorox provides preemergence control of the weeds listed in the "Command 6 EC Applied Alone" and "Command 6 EC Applied in Combinations" portions of this label.

USE RATES: Refer to Tables 1, 3, 4, 5 and 6 for use rates for Command alone, Command plus Sencor, Lexone, Lorox or Amiben.

APPLICATION AND MIXING INSTRUCTIONS

Paraquat Combinations: Tank mix Command 6 EC alone or with Sencor/Lexone, Lorox or Amiben (observe mixing instructions in the General Mixing Instructions portion of this label, as well as individual product label use directions). Add 1 to 2 pints paraquat (based on products containing 2 pounds active ingredient per gallon) per acre. Use lower rate for annual weeds less than 4 inches tall and the higher rate for weeds 4 to 6 inches tall. As the last ingredient, add a suitable surfactant (refer to paraquat label).

Roundup Combinations: Tank mix Command 6 EC alone or with Sencor/Lexone, Lorox or Amiben (observe mixing instructions in the General Mixing Instructions portion of this label, as well as individual product label use directions). Add 1½ quarts of Roundup per acre for control of emerged annual weeds or 2 to 4 quarts per acre for control of emerged perennial and annual weeds.

Apply in 20 to 60 gallons of water per acre. Use the higher gallnage for dense stubble or vegetation.

COMMAND 6 EC HERBICIDE APPLICATION FOLLOWED BY POSTEMERGENCE TREATMENTS

Postemergence herbicides approved for use on soybeans may be applied following preemergence or preplant incorporated treatments of Command alone (or tank mixtures of Command plus Sencor, Lexone, Lorox or Amiben as noted on this label) for control or emerged weeds as listed on the postemergent product labels. Read and follow the cautionary statements, directions for use, rates of application, and all other information appearing on the product labels.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use. On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

AMIBEN is a registered trademark of Union Carbide.
LEXONE and LOROX are registered trademarks of E. I. duPont de Nemours and Company.
ROUNDUP is a registered trademark of Monsanto Company.
SENCOR is a registered trademark of Bayer AG.

Command and FMC are registered trademarks of FMC Corporation.
© 1984 FMC Corporation All rights reserved