

PM 15 279-3059

1/1  
5

SEP 28 1988

216866 / 310  
3 / 18

FMC Corporation  
Agricultural Chemicals Division  
Product Registration Department  
2000 Market Street  
Philadelphia, PA 19103

Attention: S.E. Burkart, Room 2262

Gentlemen:

Subject: Amendment - Addition of Almonds and Pistachios to Label  
Pounce 1.5G  
EPA Registration No. 279-3059  
Your Application Dated February 22, 1988

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
  - a. Delete "... for at least 15 minutes" from the "If In Eyes" statement.
  - b. In the Environmental Hazards statement, change "... cleaning of equipment or disposal of wastes" to "... disposing of equipment washwaters."
3. The addition of use on almonds and pistachios has been accepted and made a part of the record.
4. Submit five (5) copies of your final printed labeling before you release the product for shipment.

53159:I/C:Richards:LLR-08:KENCO:09/22/89:11/03/88:aw:VO:EK:aw

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

1 of 6

2

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca  
Product Manager (15)  
Insecticide-Rodenticide Branch  
Registration Division (TS-767C)

Enclosure

2 of 6

Code 692

**RESTRICTED USE  
PESTICIDE**

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

**Net Weight**

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

3

SEP 28 1988

# Pounce 1.5 G

## Insecticide

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

279-3059

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

EPA Reg. No. 279-3059

EPA Est. 279-

<b>ACTIVE INGREDIENT:</b>	<b>By Wt.</b>
*Permethrin .....	1.50%
<b>INERT INGREDIENTS:</b> .....	98.50%
	100.0%

\*(3-Phenoxyphenyl) methyl (±) cis-trans 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate cis/trans ratio: Max. 55% (±) cis and min. 45% (±) trans  
U.S. Patent No. 4,024,163

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

**Caution**  
Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Remove and wash clothing before reuse.

### STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in eyes, flush with plenty of water for at least 15 minutes. Get medical attention.

If on skin, wash with soap and water.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment, or disposal of wastes.

### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

For Emergency Assistance Call (716) 735-3765



FMC Corporation  
Agricultural Chemical Group  
Philadelphia PA 19103

11/4-86-A

11/86

### STORAGE AND DISPOSAL

#### Pesticide Storage

Do not store in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

#### Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### Field Corn, Popcorn, Field Corn Grown for Seed

**Preemergent Use:** To control Armyworms and Cutworms—Use Pounce 1.5 G at a rate of 6.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre in the time period from five days prior to planting up to emergence of the crop. Apply either as a broadcast using ground or aerial equipment or as a 7-inch band. For the band application use Pounce 1.5 G at a rate of 8 to 16 ounces per 1,000 linear feet of row.

**Foliar Use:** To control Armyworms, Cutworms, European Corn Borer, Southwestern Corn Borer, Western Bean Cutworm, and Stalk Borers—Use Pounce 1.5 G at a rate of 6.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre. When treating for Armyworms, Cutworms, European Corn Borer, Southwestern Corn Borer and Western Bean Cutworm apply when eggs begin to hatch. For treatment of Stalk Borers, Pounce 1.5 G must be applied when or shortly before the Stalk Borer larvae are moving into the corn from the surrounding weeds and grasses. Mowing or burn-down herbicide is suggested to initiate movement of Stalk Borer larvae. Apply by aircraft broadcasting over corn plants or with ground equipment by directing the granule into the whorl.

Do not apply more than 0.4 pound active per acre after brown silk stage. Do not exceed a total of 1.0 pound active per acre per season.

### Sweet Corn

**Preemergent Use:** To control Armyworms and Cutworms—Use Pounce 1.5 G at a rate of 6.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre in the time period from five days prior to planting up to emergence of the crop. Apply either as a broadcast using ground or aerial equipment or as a 7-inch band. For the band application use Pounce 1.5 G at a rate of 8 to 16 ounces per 1,000 linear feet of row for 40-inch row spacing.

**Foliar Use:** To control Armyworms, Cutworms, European Corn Borer, Western Bean Cutworm, Southwestern Corn Borer and Stalk Borer—Use Pounce 1.5 G at a rate of 6.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre. When treating for Armyworms, Western Bean Cutworm, Southwestern Corn Borer, and European Corn Borer apply when eggs begin to hatch. For treatment of Stalk Borer, Pounce 1.5 G must be applied when or shortly before Stalk Borer larvae are moving into sweet corn from the surrounding weeds and grasses. Mowing or burn-down herbicide is suggested to initiate movement of Stalk Borer larvae. Apply by aircraft by broadcasting over sweet corn plants or with ground equipment by directing the granule into the whorl.

Apply every five days or as needed. Do not apply more than 1.2 pounds active ingredient per acre per crop season. Do not apply within 1 day of harvest of ears or forage or livestock grazing.

3 of 6

4

**Corn Insecticide Applicator Settings**

Pounce 1.5 G is a dense, free flowing insecticide requiring accurate equipment calibration to apply the recommended amount per acre. Calibrate your planter to the speed at which you plan to travel during planting operation. The following are approximate settings for guidance in calibration:

Application Rate Pounds/Acre	6.7			10.0			13.3		
Ounces/1000 Linear Feet	8			12			16		
Planter Speed (mph)	4	6	8	4	6	8	4	6	8
Equipment:									
				Gauge Setting					
Gandy Model 901	14	17	20	17	21	25	20	25	28
John Deere									
MaxEmerge									
(Even Notch)	8	12	16	12	17	22	15	22	27
(Odd Notch)	7	10	12	10	13	16	12	16	19
1-H Case	1.3	1.5	1.7	1.5	1.7	2.1	1.7	2.2	2.8

**Corn Application Rates**

When the granular application equipment is set to deliver Pounce 1.5 G at a rate of 12 ounces per 1,000 linear feet (0.15 pound active per acre) the total amount applied per acre will vary depending on row spacings.

Use table below to determine the Pounce 1.5 G needs for each acre:

Row Spacings (inches)	40	38	36	32	30	28
Pounce 1.5 G (pounds per acre)	9.8	10.3	10.9	12.2	13.0	14.0

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

4 of 6

Pounce 1.5 G

8/27/87

Almonds: To control ants -- Use Pounce 1.5 G at a rate of 26 pounds (0.4 pound active) per acre. Apply as a broadcast treatment by ground equipment or aircraft when ants appear. Overhead moisture following application will enhance activity.

Do not apply more than 0.8 pound active per acre during hull split. Do not apply more than 2 pounds active ingredient per acre per season. Do not apply within seven days of harvest. Do not graze livestock in treated orchards.

Pounce 1.5 G

10/21/87

Pistachios: To control ants -- Use Pounce 1.5 G at a rate of 26 pounds (0.4 pound active) per acre. Apply as a broadcast treatment by ground equipment or aircraft when ants appear. Overhead moisture following application will enhance activity.

Do not apply more than 0.8 pound active per acre per crop season. Do not apply after 10% hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated orchards.

6076