

Code 692

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

Net Weight

Pounce[®] 1.5 G Insecticide

EPA Reg. No. 279-3059

EPA Est. 279-

ACTIVE INGREDIENT:	By Wt.
"Permethrin"	1.50%
INERT INGREDIENTS:	98.50%
	100.0%

*[3-Phenoxyphenyl)methyl (+) cis-trans 3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate
**cis:trans ratio Max 55% (+) cis and min 45% (+) trans
U.S. Patent No 4,024,163

with COMMENTS
in EPA Letter # 100

SEP - 4 1991

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No.

279-3059

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in eyes, flush with plenty of water. Get medical attention if irritation persists.

If on skin, wash with soap and water.

For Emergency Assistance Call (716) 735-3785

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Caution

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Remove and wash clothing before reuse.

Environmental Hazards

This product is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Keep out of lakes, ponds or streams. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Pesticide Storage

Do not store in or around the home.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or discharge.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Contain spills. Call FMC collect (716) 735-3785.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable Refillable Containers: Do not reuse containers. Return empty containers to (716) 735-3785.

GENERAL INSTRUCTIONS

Almonds: To control ants. Use Pounce 1.5 G at a rate of 20 pounds (0.4 pound active) per acre. Apply as a broadcast treatment by ground equipment or aircraft when ants appear. (Weather measure adversely affects ant activity.)

Do not apply more than 0.8 pound active per acre during full bloom. Do not apply more than 2 pounds active ingredient per acre per season. Do not apply within seven days of harvest. Do not graze livestock in treated orchards.

Field Corn, Popcorn, Field Corn Grown for Seed: Preemergent Use To control Armyworms and Cutworms. Use Pounce 1.5 G insecticide at a rate of 8.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre in the time period from five days prior to planting up to emergence of the crop. Apply either as a broadcast using ground or aerial equipment or as a 7 inch band. For the band application use Pounce 1.5 G at a rate of 8 to 18 ounces per 1,000 linear feet of row.



FMC Corporation
Agricultural Chemical Group
Philadelphia PA 19103

BEST AVAILABLE COPY

Label Use: To control Armyworms, Cutworms, European Corn Borer, Southwestern Corn Borer, Western Bean Cutworm, and Stalk Borers—Use Pounce 1.5 G at a rate of 6.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre in treating for Armyworms, Cutworms, European Corn Borer, Southwestern Corn Borer and Western Bean Cutworm apply when eggs begin to hatch. For treatment of Stalk Borers, Pounce 1.5 G must be applied when shortly before the Stalk Borer larvae are moving into the corn from the surrounding weeds and grasses. Mowing or burn-down herbicide is suggested to initiate movement of Stalk Borer larvae. Apply by aircraft broadcasting over corn or with ground equipment by directing the granule into the whorl.

Do not apply more than 0.4 pound active per acre after brown silk stage. Do not exceed a total of 1.0 pound active per acre per season.

Label Use: To control ants—Use Pounce 1.5 G at a rate of 26 pounds (0.4 pound active) per acre. Apply as a broadcast treatment by ground equipment or aircraft when ants appear. Overhead moisture following application will increase activity.

Do not apply more than 0.8 pound active per acre per crop season. Do not apply after 10% hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated orchards.

Label Use:

To control Armyworms and Cutworms—Use Pounce 1.5 G at a rate of 6.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre in the period from five days prior to planting up to emergence of the crop. Apply as a broadcast using ground or aerial equipment or as a 7-inch band application use Pounce 1.5 G at a rate of 8 to 16 ounces per 1,000 linear feet of row for 40-inch row spacing.

Label Use: To control Armyworms, Cutworms, European Corn Borer, Western Bean Cutworm, Southwestern Corn Borer and Stalk Borer—Use Pounce 1.5 G at a rate of 6.7 to 13.3 pounds (0.1 to 0.2 pound active) per acre in treating for Armyworms, Western Bean Cutworm, Southwestern Corn Borer, and European Corn Borer apply when eggs begin to hatch. For treatment of Stalk Borer, Pounce 1.5 G must be applied when or shortly before the Stalk Borer larvae are moving into sweet corn from the surrounding weeds and grasses. Mowing or burn-down herbicide is suggested to initiate movement of Stalk Borer larvae. Apply by aircraft by broadcasting over sweet corn or with ground equipment by directing the granule into the whorl.

Apply every five days or as needed. Do not apply more than 1.2 pounds of active ingredient per acre per crop season. Do not apply within 1 day of use of ears or forage or livestock grazing.

Insecticide Applicator Settings

Pounce 1.5 G is a dense, free flowing insecticide requiring accurate equipment calibration to apply the recommended amount per acre. Calibrate your equipment to the speed at which you plan to travel during planting operation. The following are approximate settings for guidance in calibration:

Application Rate Pounds/Acre	6.7	10.0	13.3
Ounces/1000 Linear Feet	8	12	16
Travel Speed (mph)	4 6 8	4 6 8	4 6 8
Equipment	Gauge Setting		
Case Model 901	14 17 20	17 21 25	20 25 28
Case Model 901 (Even Notch)	8 12 16	12 17 22	15 22 27
(Odd Notch)	7 10 12	10 13 16	12 16 19
Case	13 15 17	15 17 21	17 22 28

Application Rates

As the granular application equipment is set to deliver Pounce 1.5 G at a rate of 12 ounces per 1,000 linear feet (0.15 pound active per acre) the amount applied per acre will vary depending on row spacings.

Refer to the table below to determine the Pounce 1.5 G needs for each acre.

Row Spacing (Inches)	40	36	32	30	28	
Pounce 1.5 G (pounds per acre)	9.8	10.3	10.9	12.2	13.0	14.0

Labels Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or storage of this material when such use and/or handling and storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practices.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

BEST AVAILABLE COPY

FILE 279-3059

1084

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

15397463
179

Ms. Linda A. Dansbury
FMC Corporation
Agricultural Chemical Group
1735 Market Street
Philadelphia, Pennsylvania 19103

Dear Ms. Dansbury:

Subject: Amendment to Revise Storage & Disposal Portion
of Label to Include U-Turn (returnable) Containers
Pounce 1.5G Insecticide
EPA Registration Number 279-3059
Your Application Dated June 3, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

0 Revise the environmental hazard section as follows:

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.


CONCURRENCES

SYMBOL	H7505C						
SURNAME	Altman						
DATE	9-14-91						

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely,



George T. LaRocca
Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

enclosure: