

**UNITED STATES ENVIRONMENTAL PROTECTION AGENCY**

WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES  
AND TOXIC SUBSTANCES

January 11, 2007

Christopher Davis  
Registration Manager  
FMC Corp  
1735 Market Street  
Philadelphia, PA 19103

Subject: Amendment – Response to Agency’s (10/21/06, 8/20/06, 9/26/05, 10/23/03 and 10/21/06) letters  
Capture 2EC Insecticide  
EPA Reg. No. 279-3069  
Your submission dated November 3, 2006

Dear Mr. Davis:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable subject to the comments listed below. Two (2) copies of the finished labeling must be submitted prior to releasing the product for shipment. A stamped copy of the label is enclosed for your records.

1. The maximum seasonal rate is 0.2lb ai/A for tobacco. Change “0.3lb ai/A per season” to “0.2lb ai/A per season” under tobacco.
2. Under the warranty statement change “should” to “must” in second paragraph and add, “To the extent consistent with applicable law” to the beginning of the sentence, “The exclusive remedy of the user or buyer .... Etc”.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

2 7 23

If you have any questions regarding this action, please contact BeWanda Alexander of my team at (703) 305-7460.

Sincerely,  
  
George T. LaRocca  
Product Manager 13  
Insecticide Branch  
Registration Division (7505C)

Enclosure

3723

**RESTRICTED USE PESTICIDE**  
Toxic to fish and aquatic organisms. For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

# CAPTURE 2EC

## Insecticide/Miticide

**CAPTURE 2EC IS NOT FOR SALE OR USE IN CALIFORNIA.**

EPA Reg. No. 279-3069

EPA Est. No. 279-

<b>Active Ingredient:</b>	<b>By Wt.</b>
Bifenthrin: (2 methyl[1,1'-biphenyl]-3-yl) methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)- 2,2-dimethyl-cyclopropanecarboxylate* .....	25.1%
<b>Other Ingredients:**</b> .....	74.9%
	100.0%

\*\*Cis isomers 97% minimum, trans isomers 3% maximum.  
\*\*Contains xylene range aromatic solvents.  
This product contains 2 pounds active ingredient per gallon.  
U.S. Patent No. 4,238,505

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated  
JAN 11 2007**

**ALL-WEATHER FORMULA**

**KEEP OUT OF REACH OF CHILDREN  
WARNING  
AVISO**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
279-3069

This label must be in the possession of the user at the time of application. Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)  
See other panels for additional precautionary information.

### FIRST AID

**If Swallowed:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.  
**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.  
**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

### HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

### NOTE TO PHYSICIAN

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided. Contains petroleum distillate - vomiting may cause aspiration pneumonia.  
For Emergency Assistance Call (800) 331-3148.

**FMC Corporation  
Agricultural Products Group  
Philadelphia, PA 19103**

**Net Contents:**

**PRECAUTIONARY STATEMENTS****Hazards to Humans (and Domestic Animals)****Warning**

May be fatal if swallowed. Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

**Personal Protective Equipment:** Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

**Handlers who may be exposed to the dilute through application or other tasks must wear:** long-sleeved shirt and long pants, chemical resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, and shoes plus socks.

**Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear:** long-sleeved shirt and long pants, chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations****Users should:**

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**Environmental Hazards**

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

**Physical/Chemical Hazards**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Resistance.** Some insects are known to develop resistance to products used repeatedly for control.

Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and

exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, and Shoes plus socks.

**STORAGE AND DISPOSAL**

**Pesticide Storage**  
 Do not freeze. Do not store below 40° F. If crystals are observed, warm material to above 60° F by placing container in warm location. Shake or roll container periodically to redissolve solids. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: 1-(800)-331-3148. To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal**  
 Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

**Container Disposal**  
 Metal or Plastic Container: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers. U-Turn® Container: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

**Chemigation Use Directions**

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

For LEPA irrigation a minimum of 0.75 inch of water per acre is recommended. Where non-emulsified oils are used as the diluent, 1 to 2 pints per acre is recommended.

Results from utilizing chemigation have been variable and depend upon the set up and calibration of equipment. Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent areas.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment. Capture® 2 EC insecticide/miticide should be applied continuously for the duration of the water application. Capture 2 EC should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

### Rotational Crops

Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days following the final application of bifenthrin.

### Tank-Mixture

Capture 2 EC Insecticide/Miticide may be applied in tank mixtures with other products approved for use on registered crops. Observe all restrictions and precautions which appear on the labels of these products. Test for compatibility of products before mixing.

### Application Instructions

Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations; higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

### Spray Drift Precautions

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes; reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm pond. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made in cotton. Use of ultra low volume (ULV) application on corn and hops is prohibited.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Do not cultivate within 10 feet of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Do not make aerial or ground applications to corn if heavy rainfall is imminent. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

**COTTON**

PEST	DOSAGE	
	LB A/A	FL OZ/A
European Corn Borer Soybean (Banded) Thrips Tobacco Thrips	0.02-0.10	1.3-6.4
Boll Weevil Bollworm Cabbage Looper Cotton Aphid Cotton Fleahopper Cotton Leafperforator Cutworms Fall Armyworm Plant Bugs Saltmarsh Caterpillar Southern Garden Leafhopper Stink Bugs Tobacco Budworm Whitely Yellow Striped Armyworm	0.04-0.10	2.6-6.4
Beet Armyworm Carmine Spider Mite Lygus Spp. Pink Bollworm Twospotted Spider Mite	0.06-0.10	3.8-6.4

**RESTRICTIONS—Cotton**

Do not apply more than 0.5 pound active per acre per season.  
 Do not apply within 14 days of harvest.  
 Do not graze livestock in treated areas or cut treated crops for feed.  
 Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Ammo®, Asana® XL, Baythroid®, Capture®, Danitol®, Karate®, Mustang®, and Scout X-TRA®.

**REMARKS—Cotton**

Capture 2 EC may be applied in water or refined vegetable oil (soybean/cottonseed).  
**Application in Water:** Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray.  
**ULV Application:** Apply the recommended rate of Capture 2 EC in refined vegetable oil in a minimum of 1 quart of finished spray per acre with aircraft calibrated to give adequate coverage.  
**To Control Boll Weevil:** Apply Capture 2 EC at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels.  
**To Control Mites and Aphids:** Apply when pests first appear. Repeat as necessary to maintain control. Higher rates will be required once a damaging threshold is established.  
 This product is not for sale or use in the State of California.

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (AT PLANT USE)**

PEST	DOSAGE		REMARKS
Corn Rootworm Larvae Northern Southern Western	0.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row	Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the Capture 2EC needs per acre. Apply in a minimum of 3 gallons of finished spray per acre.  Mix Capture 2EC with water or fertilizer in the following

Army Cutworm Cutworm Species Grubs Seed Corn Beetle Seed Corn Maggot True Armyworm or Armyworm Species Wireworm	0.0023 to 0.0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row	<p>manner Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of Capture 2EC, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.</p> <p>Applications of Capture 2EC alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of Capture 2EC and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.</p> <p>Do not apply to soil where there is greater than 30% cover of crop residue remaining.</p> <p>Do not apply within 30 days of harvest Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment. Do not apply more than 0.1 pound active per acre per season as an at plant application</p>
--	---	--	--

Row Spacings (inches)	40	38	36	30
Capture 2EC (pounds ai per acre)	0.060	0.064	0.069	0.080
Capture 2EC (formulated ounces per acre)	3.9	4.1	4.4	5.12

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (PRE & PPI)**

Pest	Dosage	Remarks
Black Cutworm White Grub Wireworm Seedcorn Maggot Armyworm spp. Stalkborer	0.047 to 0.062 LB A/A Pre-Plant Incorporated (PPI)	3 to 4 FL OZ/A Pre-Plant Incorporated (PPI)
Black Cutworm Armyworm spp. Stalkborer	0.040 LB A/A Pre- Emergence (PRE)	2.56 FL OZ/A Pre- Emergence (PRE)
		<p>The 3- 4oz/A rate must be applied as PPI and can be tankmixed and applied with PPI herbicides. Incorporation of Capture should not be any deeper than the intended planting depth and no deeper than 3 inches. Incorporation depth should be close to the intended seed planting depth</p> <p>The 2.56 oz/A rate may be applied PRE and can be tankmixed and applied with PRE herbicides.</p>

**FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (FOLIAR USE)**

PEST	DOSAGE	
	LB A/A	FL OZ/A
Aphids Army Cutworm Beet Armyworm Cereal Leaf Beetle Chinch Bug Common Stalk Borer Corn Earworm Corn Rootworm Adults Cucumber Beetle Adult Cutworm Species European Corn Borer Fall Armyworm Flea Beetle Grasshoppers Greenbug Japanese Beetle Adult Sap Beetle Southern Armyworm Southern Corn Leaf Beetle Southwestern Corn Borer Stinkbugs Tarnished Plant Bug True Armyworm or Armyworm Species Webworms Western Bean Cutworm Yellowstriped Armyworm	0.033-0.10	2.1-6.4

Banks Grass Mite Carmine Mite Twospotted Spider Mite	0.08-0.10	5.12-6.4
--	-----------	----------

**RESTRICTIONS—Corn**

Do not apply more than 0.3 pound active per acre per season including pre & ppi, at plant, plus foliar applications.  
 Do not apply within 30 days of harvest.  
 Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.  
 Use of ultra low volume (ULV) application on corn is prohibited.  
 Do not make aerial or ground applications to corn if heavy rainfall is imminent.  
 Use of Capture 2EC on corn is prohibited in all coastal counties.

**REMARKS—Corn**

**General:** Apply in a minimum of 2-5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment. To improve control by aircraft, use 5 gallons of finished spray per acre particularly when initial populations are heavier than normal. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  
**To control ear-attacking pests:** Apply Capture 2 EC just before silking and repeat as necessary to maintain control.  
**Southwestern Corn Borer, European Corn Borer:** Make application for corn borer control with initial application at or shortly before egg hatch.  
**For control of other insect pests:** Apply when pests first appear and repeat as necessary.  
**For Control of Mites:** Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.  
 For Twospotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.  
 Higher rates will be necessary for heavier initial populations and corn under heat or drought stress. Field experience with dimethoate at 0.5 lb. active per acre in tank mixture has demonstrated good control under these conditions.  
**For mite control in Texas, New Mexico, Oklahoma, Arizona:** Apply in a minimum of 5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment.

**SWEET CORN (GRAIN AND SILAGE) SWEET CORN GROWN FOR SEED (AT PLANT USE)**

PEST	DOSAGE		REMARKS
Corn Rootworm Larvae Northern Southern Western	0.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row	Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the Capture 2EC needs per acre. Apply in a minimum of 3 gallons of finished spray per acre.
Army Cutworm Cutworm Species Grubs Seed Corn Beetle Seed Corn Maggot True Armyworm or Armyworm Species Wireworm	0.0023 to 0.0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row	Mix Capture 2EC with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of Capture 2EC, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.  Applications of Capture 2EC alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of Capture 2EC and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.  Do not apply to soil where there is greater than 30% cover of crop residue remaining Do not apply within 30 days of harvest. Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment. Do not apply more than 0.1 pound active per acre per season as an at plant application.

Row Spacings (inches)	40	38	36	30
Capture 2EC (pounds ai per acre)	0.060	0.064	0.069	0.080
Capture 2EC (formulated ounces per acre)	3.9	4.1	4.4	5.12

**SWEET CORN (GRAIN AND SILAGE) SWEET CORN  
GROWN FOR SEED (FOLIAR USE)**

PEST	DOSAGE	
	LB AI/A	FL OZ/A
Aphids Army Cutworm Beet Armyworm Cereal Leaf Beetle Chinch Bug Common Stalk Borer Corn Earworm Corn Rootworm Adults Cucumber Beetle Adult Cutworm Species European Corn Borer Fall Armyworm Flea Beetle Grasshoppers Greenbug Japanese Beetle Adult Sap Beetle Southern Armyworm Southern Corn Leaf Beetle Southwestern Corn Borer Stinkbugs Tarnished Plant Bug True Armyworm or Armyworm Species Webworms Western Bean Cutworm Yellowstriped Armyworm	0.033-0.10	2.1-6.4
Banks Grass Mite Carmine Mite Twospotted Spider Mite	0.08-0.10	5.12-6.4

**RESTRICTIONS—Sweet Corn**

Do not apply more than 0.2 pounds active ingredient (12.8 ounces formulated) per acre per season.  
Do not apply within one day of harvest.  
Do not graze livestock in treated areas or cut treated crops for feed within 1 day of the last application.  
Use of ultra low volume (ULV) application on corn is prohibited.  
Do not make aerial or ground applications to corn if heavy rainfall is imminent.  
Use of Capture 2EC is prohibited in all coastal counties.

**REMARKS—Sweet Corn**

**General:** Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

**To control ear-attacking pests:** Apply Capture® 2EC when silking begins and repeat as necessary to maintain control.

**Southwestern Corn Borer, European Corn Borer:** Make 2 applications for corn borer control with the initial application at or shortly before egg hatch.

**For control of other insect pests:** Apply when pests first appear and repeat as necessary.

**For Control of Mites:**

Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.

For Twospotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.

Higher rates will be necessary for heavier initial populations and corn under heat or drought stress.

## SUCULENT PEAS AND BEANS

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Pea ( <i>Pisum</i> spp.): Dwarf pea, Edible-pod pea, English pea, Garden pea, Green pea, Snow pea, Sugar snap pea Pigeon pea Bean ( <i>Phaseolus</i> spp.): Broadbean (succulent) Lima bean (green) Runner bean, Snap bean, Wax bean Bean ( <i>Vigna</i> spp.): Asparagus bean, Blackeyed pea Chinese longbean, Cowpea, Moth bean, Southern pea Yardlong bean Jackbean Soybean (immature seed) Sword bean	Flea Beetle Grasshoppers Aster Leafhopper Leafhoppers	0.025 - 0.10	1.6 - 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season.  Do not apply within 3 days of harvest.
	Aphids Beet Armyworm Fall Armyworm Southern Armyworm Yellowstriped Armyworm Bean Leaf Beetle Cucumber Beetles Japanese beetle Adult Sap Beetle Plant Bug Stink Bugs Tarnished Plant Bug Alfalfa Caterpillar Cloverworm European Corn Borer Cutworms Western Bean Cutworm Corn Earworm Loopers Corn Rootworm Adult Thrips Webworms Pea Weevil Pea Leaf Weevil Whitefly	0.033 - 0.10	2.1 - 6.4	
	Banks Grass Mite Twospotted Spider Mite Carmine Mite Lygus Spp	0.08 - 0.10	5.12 - 6.4	

## BRASSICAS

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Head and Stem Brassica Vegetables including: Broccoli Chinese Broccoli (gailon, white flowering broccoli) Brussels Sprouts Cauliflower Cavalo broccolo Kohlrabi Cabbage Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy)	Cutworms Corn Earworm Tobacco Budworm Saltmarsh Caterpillar Leafhoppers Flea Beetles Imported Cabbageworm Cucumber Beetles Aphids Whitefly Armyworms Loopers Stink Bugs Crickets Ground Beetles Thrips Wireworm (adults) Diamondback Moth	0.033 - 0.10	2.1 - 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not apply more than 0.5 lb active ingredient (1 quart) per acre per season.  Do not make more than 5 applications after bloom.  Do not make applications less than 7 days apart.
	Banks Grass Mite Twospotted Spider Mite Carmine Mite Pacific Spider Mite Lygus Spp.	0.08 - 0.10	5.12 - 6.4	

**CANOLA, CRAMBE, RAPESEED**

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Canola Crambe Rapeseed	Aphids Cutworms Diamondback Moth Loopers Other Lepidopterous Larvae Flea Beetle Flea Hopper Grasshopper Plant Bug Stink Bugs Seedpod Weevil Thrips Whitefly Armyworms	0.033 - 0.04	2.1 - 2.6	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not apply more than 0.08 lb active ingredient (5.12 ounces) per acre per season.  Do not make applications less than 14 days apart.  Do not apply within 35 days of harvest.

**CUCURBITS**

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Chayote (fruit) Chinese waxgourd (Chinese preserving melon) Citron melon Cucumber Gherkin Gourd, edible (includes hyotan, cucuzza) ( <i>Luffa</i> spp.) (includes <i>hechima</i> , Chinese okra) ( <i>Momordica</i> spp.) (includes balsam apple, balsam pear, bitter melon, Chinese cucumber) Muskmelon (hybrids and/or cultivars of <i>Cucumis melo</i> ) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon) Pumpkin ( <i>Cucurbita</i> spp.) Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini) Squash, winter (includes butternut squash, calabaza, hubbard squash ( <i>C. mixta</i> , <i>C. pepo</i> ) includes acorn squash, spaghetti squash) Watermelon (includes hybrids and/or varieties of <i>Citrullus</i> spp.)	Aphids Cutworms Cabbage Looper Leafhoppers Cucumber Beetles Squash Bugs Melonworm Pickleworm Plant Bug Stink Bugs Rindworm Squash Vine Borer Armyworms Corn Earworm Tobacco Budworm Grasshopper	0.04 - 0.10	2.6 - 6.4	Apply in a minimum of 5 gallons of finished spray per acre by air or in a minimum of 20 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not apply more than 0.3 lb active ingredient (19.2 ounces formulated) per acre per season.  Do not make more than two applications after bloom.
	Whitefly Banks Grass Mite Twospotted Spider Mite Carmine Mite Lygus Spp.	0.08 - 0.10	5.12 - 6.4	Do not make applications less than 7 days apart.  Do not apply within 3 days of harvest.

**EGGPLANT**

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Eggplant	Colorado Potato Beetle Corn Earworm European Corn Borer Flea Beetle Cucumber Beetle Cabbage Looper Tomato Pinworm Tomato Hornworm Vegetable Leafminer Whitefly Armyworms Plant Bug Stink Bug Thrips	0.033 - 0.10	2.1 - 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not make applications less than 7 days apart  Do not apply more than 0.2 lb

	Banks Grass Mite Twospotted Spider Mite Carmin Mite Pacific Spider Mite Lygus Spp	0.08 - 0.10	5.12 - 6.4	active ingredient (12.8 ounces formulated) per acre per season.  Do not apply within 7 days of harvest.
--	---	-------------	------------	---

**LETTUCE, HEAD**

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Aphids Armyworms Corn earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Loopers Salt Marsh Caterpillar Stink bug Spp. Tobacco Budworm Whitefly	0.033 - 0.10	2.1 - 6.4	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not make applications less than 7 days apart.  A maximum of 0.5 lb. active ingredient may be applied per acre per season.  Do not apply within 7 days of harvest.
Lygus Spp. Carmin Mite Two Spotted Spider Mite	0.08 - 0.10	5.12 - 6.4	

**CANE BERRIES**

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Caneberries Including: Blackberries Bingleberries Dewberries Lowberries Marionberries Ollalieberries Youngberries Loganberries Raspberries	Leafrollers Orange Tortrix Root Weevils	0.05 - 0.10	3.2 - 6.4	Apply by air or ground equipment using sufficient water to obtain full coverage of foliage. (minimum of 10 gallons per acre by air and 50 gallons per acre by ground).  One application may be made pre-bloom and a second application may be made post bloom.  For Crown Borer, apply 0.1 lb ai/a post harvest (fall) or pre-bloom (spring), as a drench application directed at the crown of plants in a minimum of 200 gallons water/ acre. Greater efficacy is observed at higher water gallonages (up to 400 gallons/a) or in an application prior to a significant rainfall event. Do not make both pre-bloom foliar and pre-bloom drench applications.  Do not apply within 3 days of harvest.  Do not exceed 0.2 lb ai per acre per season.
	Raspberry Crown Borer Spider Mites	0.10	6.4	

**ARTICHOKE**

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	

Cribrate Weevil Artichoke Plume Moth	0.10	6.4	<p>Apply when pest population reaches damaging threshold and repeat as necessary to maintain control, but not more often than 15 day intervals.</p> <p><b>Application by ground:</b> Apply a full cover spray in a minimum of 75 gallons of finished spray per acre.</p> <p><b>Application by air:</b> Apply specified dosage in a minimum of 10 gallons per acre</p> <p>Do not exceed 0.5 lb ai per acre per season. A 5-day preharvest interval must be observed.</p>
---	------	-----	---

**PEPPERS, BELL AND NON-BELL**

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Armyworms Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Leafminers Loopers Pepper Weevil Thrips Whitefly	0.033 - 0.10	2.1 - 6.4	<p>Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft.</p> <p>Do not make applications less than 7 days apart.</p> <p>Do not apply more than 0.20 pound active ingredient per acre per season.</p> <p>Do not apply within 7 days of harvest.</p>
Lygus Spp. Broad Mite Carmine Mite Two Spotted Spider Mite	0.08 - 0.10	5.12 - 6.4	

**HOPS**

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Aphids Armyworms Cutworms Leafrollers Loopers	0.06 - 0.10	3.8 - 6.4	<p>Do not exceed 0.1 lb ai per acre per application.</p> <p>Do not exceed 0.3 lb ai per acre per season.</p> <p>A spray interval of 21 days between applications must be maintained.</p>
Root Weevils	0.05 - 0.10	3.2 - 6.4	A 14 day pre-harvest interval must be observed.
Twospotted spider mite	0.10	6.4	<p><b>Application by ground:</b> For best results, full coverage is essential. Early season recommend 100-150 gallons of spray per acre. Late season recommend 200-250 gallons of spray per acre.</p> <p>For Root Weevil control, make a directed spray to the base of the plant. Spray up the vine 3 feet and the soil surface 1.5 to 2 feet on either side of the plant.</p> <p><b>Application by air for late season control of twospotted spider mites:</b> Apply no less than 6.4 oz. (0.1 lb ai) per application in a minimum of 10 gallons per acre.</p> <p>Use of ultra low volume (ULV) application on hops is prohibited.</p>

**PEARS**

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	

<p>Aphids            Codling Moth            Cutworms            Green Fruitworm            Leafhoppers  <i>Leafminers</i>            Leafrollers            Lygus spp            Plant Bugs            Plum Curculio            San Jose Scale (Crawlers)            Stink Bugs            Tarnished            Plant Bugs</p>	0.04 to 0.20	2.6 to 12.8	<p><b>Application by ground.</b> Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.</p> <p><b>Application by air:</b> Apply the specified dosage in a minimum of 10 gallons per acre by air.</p> <p>Do not apply more than 0.5 pound active per acre per season with no more than 0.45 pound active per acre applied after petal fall.</p> <p>Apply as necessary to maintain control using a minimum of 30 day spray interval.</p> <p>Apply up to 14 days prior to harvest.</p>
<p>Twospotted Spider Mite            Yellow Mite</p>	0.06 - 0.20	3.8 to 12.8	<p>Do not graze livestock in treated orchards or cut treated cover crops for feed.</p>
<p>European Red Mite</p>	0.08 - 0.20	5.12 to 12.8	

**CITRUS** (1 day phi)

Insects Controlled	Rate of Application	Method of Application
<p><i>Diaprepes Root Weevil (Diaprepes abbreviatus)</i>                      Southern Blue Green Citrus Root Weevil (<i>Pachnaeus litus</i>)                      Blue Green Citrus Root Weevil (<i>Pachnaeus opalus</i>)                      Brown Leaf Notcher (<i>Epicraerus mexicanus</i>)                      Little Leaf Notcher (<i>Artipus flordanus</i>)</p>	<p>16 - 32 fluid ounces (0.25 - 0.50 pound active) per acre</p>	<p>Apply Capture 2EC by ground equipment to bare soil beneath citrus trees. Capture 2EC must be uniformly applied from the trunk to the drip line of tree; apply in a minimum of 40 gallons of dilute spray per acre. Greater spray volume should insure greater uniformity of coverage. A pre- and post-application irrigation may aid in the uniformity of coverage as well.</p> <p>Capture 2EC protects citrus tree roots from Diaprepes and other citrus root weevil feeding by forming a barrier which provides contact activity on newly hatched larvae (neonates). As citrus root weevil eggs hatch in new foliage, neonates fall to the soil surface beneath the tree and come in contact with Capture 2EC as they attempt to burrow into the root zone. Disturbance of the soil beneath trees should be minimized.</p> <p>Timing of Capture 2EC applications is critical. Current information suggests that peak emergence of adult Diaprepes Weevil varies by citrus growing region and these emergence peaks can be dramatically affected by environmental factors, such as soil moisture. Typically, two peaks are observed for Diaprepes, first in spring then late summer or early fall. Southern Blue-Green and Blue-Green Citrus Weevils and Fuller Rose Beetle typically exhibit a single emergence peak in the spring. Brown and Little Leaf Notchers typically exhibit three emergence peaks, spring, summer and fall. Since emergence varies seasonally and by location, timing of Capture 2EC application can be accurately forecast by observing adults. Adults are most active early morning and late afternoon, numbers can be estimated by trapping throughout spring and summer (emergence periods). Egg laying will occur for 8 to 10 weeks following adult emergence from the soil, larval invasion of the soil will begin 2-3 weeks following adult emergence. It is critical to have the Capture 2EC soil barrier in place prior to drop of the neonates.</p> <p>Capture 2EC is one of several effective tools in an integrated pest management program for Citrus Root Weevils. Application of Capture 2EC should be used in conjunction with good cultural practices, biological control of larvae and foliar control of adults. Consult local university extension personnel for current information to protect citrus trees from Citrus Root Weevils and other pests.</p>
<p>Fire ants (<i>Solenopsis spp.</i>)                      Asian cockroach (<i>Blattella asahinæ</i>)</p>	<p>6.4 - 16 fluid ounces (0.1 - 0.25 pound active) per acre.</p>	<p><b>Additional Instructions:</b>                      Apply to individual citrus resets, when not in solid planted rows, using hand-gun or shielded sprayer.</p> <p>Peak emergence of Diaprepes root weevil generally occurs in the spring. Depending on weather conditions, a minor emergence of Diaprepes root weevil may also occur in the fall.</p> <p>If the citrus grove to be treated is in an area where weather conditions are conducive to primary emergence occurring in the spring, 32 fluid ounces formulated product should be used to obtain the longest residual management of Diaprepes root weevil. If the citrus grove to be treated is in an area where weather conditions will promote more than one peak of pest emergence, 16 fluid ounces formulated product can be applied early season and 16 fluid ounces formulated product can be applied later in the season.</p>
<p>Do not apply through irrigation systems.</p> <p>Do not allow any application of Capture 2EC to contact fruit or foliage.</p> <p>Do not apply more than a total of 32 fluid ounces of formulated product (0.5 lb. a.i.) per acre per year.</p> <p>Apply the specified dosage in a minimum of 40 gallons of finished spray per acre.</p> <p>Ground application only. Do not apply by air.</p>		

**TOMATOES (1 day phi)**

Insects Controlled	Rate of Application	Method of Application
Aphids Armyworms Including: Beet Armyworm, Fall Armyworm, Southern Yellowstriped Armyworm Bean Leaf Beetle Cabbageworm Carmine Mite Cloverworm Corn earworm Corn Rootworm Cucumber Beetles Cutworms Diamondback Moth European Corn Borer Flea Beetles Flea Hopper Grasshopper Japanese Beetle (Adult) Leafhoppers Loopers Lygus Spp. Melonworm Pea Weevil Pea Leaf Weevil Pickleworm Plant Bug Rindworm Salt Marsh Caterpillar Sap Beetle Seedpod Weevil Squash Bugs Stink bug Spp. Tobacco Budworm Tarnished Plant Bug Thrips Two Spotted Spider Mite Whitefly	2.1 - 5.2 fluid ounces (0.033 - 0.08 pound active) per acre.	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment.  Thorough coverage is essential to achieve control.
Do not make applications less than 10 days apart. A maximum of 4 applications may be applied per season.		

**Spinach (40 day phi)**

Insects Controlled	Rate of Application	Method of Application
Colorado Potato Beetle Tomato Pinworm Tomato Hornworm Armyworms Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Leafminers Loopers Pepper Weevil Thrips Whitefly	2.1 to 6.4 ounces (0.033 to 0.10 pound active) per acre	For control of whiteflies apply foliar treatments of Capture® insecticide by ground or air at rates of up to 0.4 pt. (0.1 lb active) per acre at minimum 7-day intervals up to a maximum of 4 applications. Do not apply within 40 days of harvest.  For control of fire ants apply Capture® insecticide to the soil (at planting) or as a foliar treatment by ground or air at rates of up to 0.4 pt. (0.1 lb active) per acre at minimum 7-day intervals up to a maximum of 4 applications. Do not apply within 40 days of harvest.  Apply the specified dosage in 5-50 gallons of finished spray per acre by air or 10-50 gallons finished spray per acre by ground
Broad Mite Banks Grass Mite Twospotted Spider Mite Carmine Mite Pacific Spider Mite Lygus Spp. Fire Ants	5.12 to 6.4 ounces (0.08 to 0.10 pound active) per acre	
Do not make applications less than 7 days apart. Do not apply more than 0.4 pounds active ingredient per acre per season.		

**GRAPES**

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Cutworms Eastern grape leafhopper Grape berry moth Japanese beetles adults Variegated leafhopper Western grape leafhopper	0.05 - 0.10	3.2 - 6.4	Apply in a minimum of 10 gallons of finished spray by air or in a minimum of 25 gallons of finished spray with ground equipment.  When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray.  Thorough coverage is essential to achieve control.
Black vine weevil Glasswinged sharpshooter Twospotted spider mite	0.10	6.4	When pest pressure is moderate to severe, use higher rate.  Do not apply more than 0.10 lb ai per acre per season.  Do not apply within 30 days of harvest.

**OKRA**

PEST	DOSAGE		REMARKS
	LBS A/A	FL OZ/A	
Amyworms Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Leafminers Loopers Thrips Whitefly Aphids Japanese Beetle (Adult), Stink bugs	0.033 to 0.10	2.1 to 6.4	Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft.  Do not make applications less than 7 days apart.  Do not apply more than 0.20 pound active ingredient per acre per season.
Lygus Spp. Broad Mite Carmine Mite Two Spotted Spider Mite	0.08 to 0.10	5.12 to 6.4	Do not apply within 7 days of harvest.

**CILANTRO, CORIANDER**

PEST	DOSAGE		REMARKS
	LBS A/A	FL OZ/A	
Spotted Cucumber Beetle Beet Armyworm Cabbage Looper Aphids Whitefly Flea beetle Thrips Leafminer Cutworm Grasshoppers Saltmarsh caterpillar	0.033 to 0.10	2.1 to 6.4	Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft.  Do not make applications less than 7 days apart.  Do not apply more than 0.50 pound active ingredient per acre per season.
Two Spotted Spider Mite	0.08 to 0.10	5.12 to 6.4	Do not apply within 3 days of harvest.

**DRIED BEANS AND PEAS**

CROP	PEST	DOSAGE		REMARKS
		LBS A/A	FL OZ/A	
Dried cultivars of: Bean (Lupinus) Bean (Phaseolus) Field bean Kidney bean Lima bean (dry) Navy bean Pinto bean Tepary bean Bean (Vigna) Adzuki bean Blackeyed pea Cajana Cowpea Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean Broad bean (dry) Chickpea Guar Lablab bean Lentil Pea (Pisicum) Field pea Pigeon pea	Flea Beetle Grasshoppers Aster Leafhopper Leafhoppers	0.025 to 0.10	1.6 to 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment.  When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) to peas, or 0.3 active ingredient (19.2 ounces formulated) to beans per acre per season.  Do not apply within 14 days of harvest.  Do not make applications less than 7 days apart
	Aphids Beet Armyworm Fall Armyworm Southern Armyworm Yellowstriped Armyworm Bean Leaf Beetle Cucumber Beetles Japanese beetle Adult Sap Beetle Plant Bug Stink Bugs Tarnished Plant Bug Alfalfa Caterpillar Cloverworm European Corn Borer Cutworms Western Bean Cutworm Corn Earworm Loopers Corn Rootworm Adult Thrips Webworms Pea Weevil Pea Leaf Weevil Whitefly	0.033 to 0.10	2.1 to 6.4	
	Imported cabbageworm Saltmarsh caterpillar Tobacco budworm Leafminer	0.08 to 0.10	5.12 to 6.4	
	Banks Grass Mite Twospotted Spider Mite Carmine Mite Lygus Spp			

20 7 23

## LEAFY BRASSICAS

CROP	PEST	DOSAGE		REMARKS
		LBS A/A	FL OZ/A	
Broccoli Raab Bok Choy Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens	Cutworms Corn Earworm Tobacco Budworm Saltmarsh Caterpillar Leafhoppers Flea Beetles Imported Cabbageworm Cucumber Beetles Aphids Whitefly Armyworms Loopers Stink Bugs Crickets Ground Beetles Thrips Wireworm (adults) Diamondback Moth Japanese Beetle (adult) Grasshoppers Aphids	0.033 to 0.10	2.1 to 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.  Do not apply more than 0.4 lb. active ingredient per acre per season  Do not make applications less than 7 days apart  Do not apply within 7 days of harvest
	Banks Grass Mite Twospotted Spider Mite Carmine Mite Pacific Spider Mite Lygus Spp.	0.08 to 0.10	5.12 to 6.4	

**TUBEROUS AND CORM VEGETABLES**

CROP	PEST	DOSAGE		REMARKS
		LBS A/A	FL OZ/A	
Potato	Corn wireworm	0.30	19.2	Capture may be applied as an in-furrow planting time treatment for the control of wireworms, rootworms and white grubs. Apply Capture at the rate of 0.3 pounds active per acre as an in-furrow spray or T-band spray at planting time.
Sweet potato	Tobacco wireworm	(at-plant)	(at-plant)	
Arracacha				
Arrowroot				
Chinese artichoke	Southern potato wireworm	0.05 to 0.15	3.2 to 9.6	Capture may be applied as a lay-by treatment for the control of wireworms, rootworms and white grubs. Apply Capture to the drill area and cover with soil utilizing cultivation equipment set to throw soil to the drill area. Apply Capture as a banded spray over the row at a rate of 0.05-0.15 pounds active per acre (3.2 to 9.6 ounces formulated) in 10 gallons per acre of spray.
Jerusalem artichoke	Japanese beetle grubs	(lay-by)	(lay-by)	
Edible canna	June beetle			Capture may be applied as a lay-by treatment for the control of wireworms, rootworms and white grubs. Apply Capture to the drill area and cover with soil utilizing cultivation equipment set to throw soil to the drill area. Apply Capture as a banded spray over the row at a rate of 0.05-0.15 pounds active per acre (3.2 to 9.6 ounces formulated) in 10 gallons per acre of spray.
Cassava (bitter and sweet)	Sweetpotato flea beetle	0.033 to 0.10	2.1 to 6.4	
Chayote (root)	Cucumber beetle	(foliar)	(foliar)	Capture may be applied as a foliar spray for the control of the adult life stages of flea beetles, click beetles (wireworms), cucumber beetles (rootworms), white fringed beetles and May/June beetles (white grubs). Apply Capture at the rate of 0.1 lbs active per acre (6.4 ounces formulated) in 10 gallons of spray by ground and 3 gallons of spray by air.
Chufa	Sweetpotato weevil			
Dasheen (taro)	Banded Cucumber beetle			Do not make more than 2 foliar applications per season no sooner than 21 days apart.
Ginger	Black flea beetle			
Leren	Whitefringed beetle			Do not apply more than 0.5 lb active ingredient per acre per season, including soil application.
Tanier	White grub			
Turmer	Sugarcane beetle			Do not apply within 21 days of harvest
Yam bean	Rootworms			
True yam				

**TOBACCO**

CROP	PEST	DOSAGE		REMARKS
		LBS AI/A	FL OZ/A	
Tobacco	Cutworm spp. Tobacco Flea Beetle (larvae) White Grubs Wireworms Mole Crickets Armyworm spp. Stalkborers	0.0625 - 0.10	4.0 - 6.4	Pre-transplant soil applications: Apply 0.0625- 0.1 lb ai/A in a minimum of 10 gal/A to control soil pests. Use of suitable equipment to incorporate into top 4" of the soil is required to control below ground pests.  At-transplant water treatment application: Apply 0.0625- 0.1 lb ai/A in a water treatment application volume of 10-200 gal/A.
	Aphid spp. Armyworm spp. Flea Beetle (Adults) Chinch Bugs Stink Bugs Japanese Beetles Grasshoppers Cutworm spp. Tarnished Plant Bugs Green Bugs Thrips Whiteflies	0.04- 0.10	2.56 - 6.4	Foliar applications: Apply 0.04- 0.10 lb ai/A per foliar application up to, and including, layby in a minimum of 10 gal/A. Do not make more than 2 foliar applications per season.
	Spider mites Lygus spp.	0.1	6.4	
<p>Do not apply more than 0.3 lb ai/A per season.</p> <p>Do not apply later than layby.</p> <p>May be tank mixed with Command, Spartan and other herbicides approved for tobacco use.</p>				

**Dealers Should Sell in Original Packages Only.**

**Conditions of Sale and Limitation of Warranty and Liability:**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors. Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. **FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT.** Any warranties, express or implied, having been made are inapplicable if this product has been contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use. To the extent allowed by law, FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.** This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC, Ammo, Capture, Mustang, U-Turn—trademarks of FMC Corporation  
Ambush, Karate—trademarks of Zeneca, Inc.  
Asana—trademark of E.I. duPont de Nemours & Company  
Baythroid—trademark of Bayer Aktiengesellschaft  
Danitol—trademark of Sumitomo Chemical Company, Ltd.  
Scout Xtra—trademark of Hoechst Schering Agrevo S.A.

© 2003-2006 FMC Corporation All rights reserved.