



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Christopher Davis
Registration Manager
APG Specialty Products Business
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

OCT 10 2008

SUBJECT: Application for Pesticide Notification – Proposed Final Printed Label with Additional Changes via Notification for Label Revisions under PRN 2007-4 and PRN 98-10

Capture 2EC Insecticide/Miticide
EPA Registration No. 279-3069
September 2, 2008

Dear Mr. Davis:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 and 2007-4 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and 2007-4. PLEASE NOTE: final printed labels must be submitted to the appropriate Product Manager (PM), not as a Notification under PRN 98-10 and 2007-4. Therefore, the "proposed" label submitted for the above product were reviewed as a Notification under PRN 98-10 and 2007-4, not as final printed label which must still be submitted to the appropriate PM.

However, the Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and 2007-4 and finds that the actions requested fall within the scope of PRN 98-10. The labels submitted with the applications have been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2/39

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0080. Approval expires 2-28-95

| | | |
|--|--|-----------------------|
|  <p>United States Environmental Protection Agency Washington, DC 20460</p> | <input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other | OPP Identifier Number |
|--|--|-----------------------|

Application for Pesticide - Section I

| | | |
|---|-----------------------------------|--|
| 1. Company/Product Number 279-3069 | 2. EPA Product Manager LaRocca | 3. Proposed Classification <input type="checkbox"/> None <input checked="" type="checkbox"/> Restricted |
| 4. Company/Product (Name) Capture 2EC Insecticide/Miticide | PM# 13 | |
| 5. Name and Address of Applicant (Include ZIP Code) FMC Corporation 1735 Market Street Philadelphia, PA 19103 <input type="checkbox"/> Check if this is a new address | | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar and identical to the product and labeling to: <div style="text-align: center; font-size: 1.5em; font-weight: bold;">NOTIFICATION</div> EPA Reg. No. _____ <div style="text-align: center; font-size: 1.2em;">OCT 10 2008</div> Product Name _____ |

Section - II

| | |
|--|--|
| <input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below. | <input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated <u>January 17, 2008</u> <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - Explain below. |
|--|--|

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

- Amend Container Disposal statements per PR-Notice 2007-4
- Add company street address
- Modify warranty to be consistent with all FMC labels
- Add insects to the soybean use directions
- Add tomato aerial dilution
- Make changes required in Agency letter of 1/17/08

Section - III

1. Material This Product Will Be Packaged In:

| | | | |
|---|--|---|--|
| Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No | 2. Type of Container <input checked="" type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____ |
| * Certification must be submitted | | If "Yes" Unit Packaging wgt. No. per container | If "Yes" Package wgt No. per container |

| | | |
|--|--|--|
| 3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container | 4. Size(s) Retail Container 15 gal (metal), 1 gal, 1 qt | 5. Location of Label Directions <input type="checkbox"/> On leaflet attached to container |
| 6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Other Plastic sleeve <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled | | |

Section - IV


1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

| | | |
|---------------------------|-------------------------------|---|
| Name Christopher Davis | Title Registration Manager | Telephone No. (Include Area Code) 215-299-6334 |
|---------------------------|-------------------------------|---|

Certification


I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

8. Date Application Received
(Stamped)

| | | |
|---|----------------------------------|--|
| 2. Signature  | 3. Title Registration Manager | |
| 4. Typed Name Christopher Davis | 5. Date 9/24/08 | |

CERTIFICATION STATEMENT WITH RESPECT TO PR-NOTICE 98-10**Notification of label change per PR Notice 98-10:**

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.



Agent for FMC Corporation

9/26/08

Date

A collection of 25 small, stylized line drawings of various birds, including sparrows, finches, and songbirds, arranged in a grid-like fashion. The drawings are simple, using black outlines on a white background, and represent different species of birds.


Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

| PROPOSED LABEL | | |
|--------------------|-----------------------|--------------------------------------|
| EPA Registration # | Date Submitted to EPA | Electronic file name |
| 279-3069 | 9/2/08 | 000279-03069.20080902.FinalPrint.pdf |

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.


Signature

9/2/08
Date

Christopher Davis
Name (typed)

Registration Manager
Title

000001
000002
000003
000004
000005
000006
000007
000008
000009
000010
000011
000012
000013
000014
000015
000016
000017
000018
000019
000020
000021
000022
000023
000024
000025
000026
000027
000028
000029
000030
000031
000032
000033
000034
000035
000036
000037
000038
000039
000040
000041
000042
000043
000044
000045
000046
000047
000048
000049
000050
000051
000052
000053
000054
000055
000056
000057
000058
000059
000060
000061
000062
000063
000064
000065
000066
000067
000068
000069
000070
000071
000072
000073
000074
000075
000076
000077
000078
000079
000080
000081
000082
000083
000084
000085
000086
000087
000088
000089
000090
000091
000092
000093
000094
000095
000096
000097
000098
000099
000100

FMC Agricultural Products

6/39

FMC Corporation
1735 Market Street
Philadelphia, PA 19103

215.299.6000 Phone
215.299.6468 Fax

www.fmc.com

September 2, 2008

Mr. George T. LaRocca (PM-13)
United States Environmental Protection Agency
Document Processing Desk (FIN-LABEL)(E-SUB)
Office of Pesticide Programs (H7504P)
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

Dear Mr. LaRocca:

Subject: Capture[®] 2EC Insecticide/Miticide, EPA Reg. No. 279-3069
-Submission of Final Printed Label
-Notification of Additional Changes

FMC is submitting final printed labeling for the subject product. FMC has revised the labeling as requested in item #'s 6(a)-(c) of the Agency comment letter dated January 17, 2008.

Please note that we are also revising the labels as follows by notification:

- Amend the Container Disposal statements per PR-Notice 2007-4
- Add company street address
- Modify warranty to be consistent with all FMC labels
- Add an aerial use dilution to the tomato use directions
- Add insects to the soybean use directions table

This label also contains the new updated spray drift wording required for agricultural use pyrethroid insecticides in your letter of February 21, 2008.

To process this request, enclosed please find the following:

- Application for Pesticide (EPA Form 8570-1).
- Final Print Labeling (3 copies), including one hi-lited version for ease of review.
- Completed Certification with Respect to Label Integrity form
- A compact disk containing the label as amended
- A certification statement with respect to PR-Notice 2007-4

Thank you for your attention to this matter, please feel free to contact me at (215) 299-6334 or e-mail at chris.davis@fmc.com.



Sincerely,



Christopher Davis
Registration Manager
APG Specialty Products Business

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms. For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

CAPTURE 2EC

Insecticide/Miticide

NOTIFICATION

OCT 10 2008

CAPTURE 2EC IS NOT FOR SALE OR USE IN CALIFORNIA.

EPA Reg. No. 279-3069

EPA Est. No. 279-

Active Ingredient:

By Wt.

Bifenthrin: (2 methyl-1,1'-biphenyl-3-yl)

methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)-

2,2-dimethyl-cyclopropanecarboxylate* **25.1%**

Other Ingredients:** **74.9%**

100.0%

**Cis isomers 97% minimum, trans isomers 3% maximum.

**Contains xylene range aromatic solvents.

This product contains 2 pounds active ingredient per gallon.

U.S. Patent No. 4,238,505

ALL-WEATHER FORMULA

KEEP OUT OF REACH OF CHILDREN

WARNING

AVISO

This label must be in the possession of the user at the time of application. Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See other panels for additional precautionary information.

FIRST AID

If Swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

NOTE TO PHYSICIAN

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided. Contains petroleum distillate - vomiting may cause aspiration pneumonia.

For Emergency Assistance Call (800) 331-3148.

FMC Corporation

Net Contents:

**Agricultural Products Group
1735 Market Street
Philadelphia, PA 19103**

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

May be fatal if swallowed. Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment: Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Handlers who may be exposed to the dilute through application or other tasks must wear: long-sleeved shirt and long pants, chemical resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, and shoes plus socks.

Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, shoes plus socks, and protective eyewear. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not make applications when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

The use of bifenthrin is prohibited in areas that may result in exposure of endangered species to bifenthrin. Prior to use in a particular county contact the local extension service for procedures and precautions to use to protect endangered species.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Resistance. Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause,

Immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, and Shoes plus socks.

STORAGE AND DISPOSAL

Pesticide Storage

Do not freeze. Do not store below 40° F. If crystals are observed, warm material to above 60° F by placing container in warm location. Shake or roll container periodically to redissolve solids. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: 1-(800)-331-3148. To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

U-Turn® Container: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

Metal or Plastic Container: Non-refillable container (in sizes 5 gallons or less)- Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank and drain for 10 seconds after flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds, pour rinsate into application equipment or mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers.

Non-refillable container (in sizes greater than 5 gallons)- Do not reuse or refill this container. Triple rinse or pressure rinse. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Returnable/Refillable Containers (if other than U-Turn Container): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents into application equipment or mix tank. Fill the container about 10% full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

For LEPA irrigation a minimum of 0.75 inch of water per acre is recommended. Where non-emulsified oils are used as the diluent, 1 to 2 pints per acre is recommended.

Results from utilizing chemigation have been variable and depend upon the set up and calibration of equipment. Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent areas.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment. Capture® 2 EC insecticide/miticide should be applied continuously for the duration of the water application. Capture 2 EC should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

Rotational Crops

Crops for which bifenthrin tolerances exist may be rotated at any time. All other crops may be rotated 30 days following the final application of bifenthrin.

Tank-Mixture

Capture 2 EC Insecticide/Miticide may be applied in tank mixtures with other products approved for use on registered crops. Observe all restrictions and precautions which appear on the labels of these products. Test for compatibility of products before mixing.

Application Instructions

Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations; higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates. Cultivation within 10 feet of a water body is prohibited to allow for the growth of a vegetated filter strip.

In New York State this product may not be applied within 100 feet (using ground equipment) to 300 feet (using aerial equipment) of coastal marshes or streams that drain into coastal marshes.

BUFFER ZONES

Vegetative Buffer Zones

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing zeta-cypermethrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21pp.*

<http://www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf>.

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast) – Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application - Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application – Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

Spray Drift Requirements

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only Medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

COTTON

| CROP | PEST | DOSAGE | |
|--------|---|-----------|---------|
| | | LB AI/A | FL OZ/A |
| Cotton | European Corn Borer Soybean (Banded) Thrips Tobacco Thrips | 0.02-0.10 | 1.3-6.4 |
| | Boll Weevil Bollworm Cabbage Looper Cotton Aphid Cotton Fleahopper Cotton Leafperforator Cutworms Fall Armyworm Plant Bugs Saltmarsh Caterpillar Southern Garden Leafhopper Stink Bugs Tobacco Budworm Whitefly Yellow Striped Armyworm | 0.04-0.10 | 2.6-6.4 |
| | Beet Armyworm Carmine Spider Mite Lygus Species Pink Bollworm Twospotted Spider Mite | 0.06-0.10 | 3.8-6.4 |

RESTRICTIONS—Cotton

Do not apply more than 0.5 pound active per acre per season.

Do not apply within 14 days of harvest.

Do not graze livestock in treated areas or cut treated crops for feed.

Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Ammo®, Asana® XL, Baythroid®, Capture®, Danitol®, Karate®, Mustang®, and Scout X-TRA®.

REMARKS—Cotton

Capture 2 EC may be applied in water or refined vegetable oil (soybean/cottonseed).

Application in Water: Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray.

ULV Application: Apply the recommended rate of Capture 2 EC in refined vegetable oil in a minimum of 1 quart of finished spray per acre with aircraft calibrated to give adequate coverage.

To Control Boll Weevil: Apply Capture 2 EC at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels.

To Control Mites and Aphids: Apply when pests first appear. Repeat as necessary to maintain control. Higher rates will be required once a damaging threshold is established.

This product is not for sale or use in the State of California.

FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (AT PLANT USE)

| CROP | PEST | DOSAGE | | REMARKS |
|---|---|----------------------------|-----------------------------|---|
| | | LB AI/1000 linear row feet | FL OZ /1000 linear row feet | |
| Field Corn (Grain and Silage) Popcorn Field Corn Grown For Seed | Northern Corn Rootworm Larvae Southern Corn Rootworm Larvae Western Corn Rootworm Larvae | 0.0046 | 0.30 | Apply as a 5 to 7 Inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the Capture 2EC needs per acre. Apply in a minimum of 3 gallons of finished spray per acre. Mix Capture 2EC with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of Capture 2EC, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture. |
| | Army Cutworm Armyworm Species Cutworm Species Grubs Seed Corn Beetle Seed Corn Maggot True Armyworm Wireworm | 0.0023 - 0.0046 | 0.15-0.30 | Applications of Capture 2EC alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed before tank mixing with appropriate ratio of Capture 2EC and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained in the spray tank during mixing and application. Do not apply to soil where there is greater than 30% cover of crop residue remaining. Do not apply within 30 days of harvest Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment. Do not apply more than 0.1 pound active per acre per season as an at plant application. |

| | | | | |
|--|-------|-------|-------|-------|
| Row Spacings (inches) | 40 | 38 | 36 | 30 |
| Capture 2EC (pounds ai per acre) | 0.060 | 0.064 | 0.069 | 0.080 |
| Capture 2EC (formulated ounces per acre) | 3.9 | 4.1 | 4.4 | 5.12 |

FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (PRE & PPI)

| CROP | PEST | DOSAGE | | REMARKS |
|---|--|---|-------------------------------------|---|
| | | LB AI/A | FL OZ/A | |
| Field Corn (Grain and Silage) Popcorn Field Corn Grown For Seed | Armyworm Species Black Cutworm Seedcorn Maggot Stalkborer White Grub Wireworm | 0.047-0.062 Pre-Plant Incorporated (PPI) | 3-4 Pre-Plant Incorporated (PPI) | The 3- 4oz/A rate must be applied as PPI and can be tank mixed and applied with PPI herbicides. Incorporation of Capture should not be any deeper than the intended planting depth and no deeper than 3 inches total. Incorporation depth should be designed to place Capture close to the seed planting depth. |

| | | | | |
|--|--|-------------------------------------|------------------------------------|--|
| | Armyworm Species Black Cutworm Stalkborer | 0.040 Pre- Emergence (PRE) | 2.56 Pre- Emergence (PRE) | The 2.56 oz/A rate may be applied PRE and can be tank mixed and applied with PRE herbicides. |
|--|--|-------------------------------------|------------------------------------|--|

FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (FOLIAR USE)

| CROP | PEST | DOSAGE | |
|---|---------------------------|------------|----------|
| | | LB AI/A | FL OZ/A |
| Field Corn (Grain and Silage) Popcorn Field Corn Grown For Seed | Aphids | 0.033-0.10 | 2.1-6.4 |
| | Army Cutworm | | |
| | Armyworm Species | | |
| | Beet Armyworm | | |
| | Cereal Leaf Beetle | | |
| | Chinch Bug | | |
| | Common Stalk Borer | | |
| | Corn Earworm | | |
| | Corn Rootworm Adults | | |
| | Cucumber Beetle | | |
| | Adults | | |
| | Cutworm Species | | |
| | European Corn Borer | | |
| | Fall Armyworm | | |
| | Flea Beetle | | |
| | Grasshoppers | | |
| | Greenbug | | |
| | Japanese Beetle Adult | | |
| | Sap Beetle | | |
| | Southern Armyworm | | |
| | Southern Corn Leaf Beetle | | |
| | Southwestern Corn Borer | | |
| | Stinkbugs | | |
| | Tarnished Plant Bug | | |
| | True Armyworm | | |
| | Webworms | | |
| | Western Bean Cutworm | | |
| | Yellowstriped Armyworm | | |
| | Banks Grass Mite | 0.08-0.10 | 5.12-6.4 |
| | Carmine Mite | | |
| | Twospotted Spider Mite | | |

RESTRICTIONS—Corn

Do not apply more than 0.3 pound active per acre per season including pre & ppl, at plant, plus foliar applications.

Do not apply within 30 days of harvest.

Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application. Use of ultra low volume (ULV) application on corn is prohibited.

Do not make aerial or ground applications to corn if heavy rainfall is imminent.

Use of Capture 2EC on corn is prohibited in all coastal counties.

REMARKS—Corn

General: Apply in a minimum of 2-5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment. To improve control by aircraft, use 5 gallons of finished spray per acre particularly when initial populations are heavier than normal. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

To control ear-attacking pests: Apply Capture 2 EC just before silking and repeat as necessary to maintain control.

Southwestern Corn Borer, European Corn Borer: Make application for corn borer control with initial application at or shortly before egg hatch.

For control of other insect pests: Apply when pests first appear and repeat as necessary.

For Control of Mites: Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant. For Twospotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.

Higher rates will be necessary for heavier initial populations and corn under heat or drought stress. Field experience with dimethoate at 0.5 lb. active per acre in tank mixture has demonstrated good control under these conditions.

For mite control in Texas, New Mexico, Oklahoma, Arizona: Apply in a minimum of 5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment.

SWEET CORN (GRAIN AND SILAGE) SWEET CORN GROWN FOR SEED (AT PLANT USE)

| CROP | PEST | DOSAGE | | REMARKS |
|---|---|----------------------------|-----------------------------|--|
| | | LB AI/1000 linear row feet | FL OZ /1000 linear row feet | |
| Sweet Corn Sweet Corn grown for seed | Northern Corn Rootworm Larvae Southern Corn Rootworm Larvae Western Corn Rootworm Larvae | 0.0046 | 0.30 | Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the Capture 2EC needs per acre. Apply in a minimum of 3 gallons of finished spray per acre. |
| | Army Cutworm Armyworm Species Cutworm Species Grubs Seed Corn Beetle Seed Corn Maggot True Armyworm Wireworm | 0.0023 - 0.0046 | 0.15-0.30 | <p>Mix Capture 2EC with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of Capture 2EC, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.</p> <p>Applications of Capture 2EC alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of Capture 2EC and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.</p> <p>Do not apply to soil where there is greater than 30% cover of crop residue remaining. Do not apply within 30 days of harvest. Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment. Do not apply more than 0.1 pound active per acre per season as an at plant application.</p> |

| | | | | |
|--|-------|-------|-------|-------|
| Row Spacings (inches) | 40 | 38 | 36 | 30 |
| Capture 2EC (pounds ai per acre) | 0.060 | 0.064 | 0.069 | 0.080 |
| Capture 2EC (formulated ounces per acre) | 3.9 | 4.1 | 4.4 | 5.12 |

**SWEET CORN (GRAIN AND SILAGE) SWEET CORN
GROWN FOR SEED (FOLIAR USE)**

| CROP | PEST | DOSAGE | |
|--|---|------------|----------|
| | | LB AI/A | FL OZ/A |
| Sweet Corn Sweet Corn grown for seed | Aphids Army Cutworm Armyworm Species Beet Armyworm Cereal Leaf Beetle Chinch Bug Common Stalk Borer Corn Earworm Corn Rootworm Adults Cucumber Beetle Adult Cutworm Species European Corn Borer Fall Armyworm Flea Beetle Grasshoppers Greenbug Japanese Beetle Adult Sap Beetle Southern Armyworm Southern Corn Leaf Beetle Southwestern Corn Borer Stinkbugs Tarnished Plant Bug True Armyworm Webworms Western Bean Cutworm Yellowstriped Armyworm | 0.033-0.10 | 2.1-6.4 |
| | Banks Grass Mite Carmine Mite Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 |

| | |
|--|--|
| | <p>RESTRICTIONS—Sweet Corn Do not apply more than 0.2 pounds active ingredient (12.8 ounces formulated) per acre per season. Do not apply within one day of harvest. Do not graze livestock in treated areas or cut treated crops for feed within 1 day of the last application. Use of ultra low volume (ULV) application on corn is prohibited. Do not make aerial or ground applications to corn if heavy rainfall is imminent. Use of Capture 2EC is prohibited in all coastal counties.</p> <p>REMARKS—Sweet Corn General: Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. To control ear-attacking pests: Apply Capture® 2EC when silking begins and repeat as necessary to maintain control. Southwestern Corn Borer, European Corn Borer: Make 2 applications for corn borer control with the initial application at or shortly before egg hatch. For control of other insect pests: Apply when pests first appear and repeat as necessary. For Control of Mites: Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant. For Twospotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy. Higher rates will be necessary for heavier initial populations and corn under heat or drought stress.</p> |
|--|--|

SUCCULENT PEAS AND BEANS

| CROP | PEST | DOSAGE | | REMARKS |
|---|---|------------|----------|---|
| | | LB/AI/A | FL OZ/A | |
| Pea (Pisum Species): Dwarf pea, Edible-pod pea, English pea, Garden pea, Green pea, Snow pea, Sugar snap pea Pigeon pea Bean (Phaseolus Species): Broadbean (succulent) Lima bean (green) Runner bean, Snap bean, Wax bean Bean (Vigna Species): Asparagus bean, Blackeyed pea Chinese longbean, Cowpea, Moth bean, Southern pea Yardlong bean Jackbean Soybean (immature seed) Sword bean | Aster Flea Beetle Grasshoppers Leafhopper Leafhoppers | 0.025-0.10 | 1.6-6.4 | Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season. Do not apply within 3 days of harvest. |
| | Adult Sap Beetle Alfalfa Caterpillar Aphids Bean Leaf Beetle Beet Armyworm Cloverworm Corn Earworm Corn Rootworm Adult Cucumber Beetles Cutworms European Corn Borer Fall Armyworm Japanese beetle Loopers Pea Leaf Weevil Pea Weevil Plant Bug Southern Armyworm Stink Bugs Tarnished Plant Bug Thrips Webworms Western Bean Cutworm Whitefly Yellowstriped Armyworm | 0.033-0.10 | 2.1-6.4 | |
| | Banks Grass Mite Carmine Mite Lygus Species Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | |
| | | | | |

BRASSICAS

| CROP | PEST | DOSAGE | | REMARKS |
|--|--|------------|----------|---|
| | | LB/AI/A | FL OZ/A | |
| Head and Stem Brassica Vegetables Including: Broccoli Chinese Broccoli (gallon, white flowering broccoli) Brussels Sprouts Cauliflower Cavalo broccolo Kohlrabi Cabbage Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy) | Aphids Armyworm Species Budworm Corn Earworm Crickets Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Ground Beetles Imported Cabbageworm Leafhoppers Loopers Saltmarsh Caterpillar Stink Bugs Thrips Tobacco Whitefly Wireworm (adults) | 0.033-0.10 | 2.1-6.4 | Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.5 lb. active ingredient (1 quart) per acre per season. Do not make more than 5 applications after bloom. Do not make applications less than 7 days apart. Do not apply within 7 days of harvest. |
| | Banks Grass Mite Carmine Mite Lygus Species Pacific Spider Mite Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | |

CANOLA, CRAMBE, RAPESEED

| CROP | PEST | DOSAGE | | REMARKS |
|------------------------------|--|------------|---------|---|
| | | LB/AI/A | FL OZ/A | |
| Canola Crambe Rapeseed | Aphids Armyworm Species Cutworms Diamondback Moth Flea Beetle Flea Hopper Grasshopper Larvae Lepidopterous Loopers Moth Other Plant Bug Seedpod Stink Bugs Thrips Weevil Whitefly | 0.033-0.04 | 2.1-2.6 | <p>Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.</p> <p>Do not apply more than 0.08 lb active ingredient (5.12 ounces) per acre per season.</p> <p>Do not make applications less than 14 days apart.</p> <p>Do not apply within 35 days of harvest.</p> |

CUCURBITS

| CROP | PEST | DOSAGE | | REMARKS |
|--|---|------------|---------|--|
| | | LB/AI/A | FL OZ/A | |
| Chayote (fruit) Chinese waxgourd (Chinese preserving melon) Citron melon Cucumber Gherkin Gourd, edible (includes hyotan, cucurza) (Luffa Species) (includes hechima, Chinese okra) (Momordica Species) (includes balsam apple, balsam pear, bitter melon, Chinese cucumber) Muskmelon (hybrids and/or cultivars of Cucumis melo) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon) Pumpkin (Cucurbita Species) Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini) Squash, winter (includes butternut squash, calabaza, hubbard squash (C. mixta; C. pepo) includes acorn squash, spaghetti squash) Watermelon (includes hybrids and/or varieties of Citrullus Species). | Aphids Armyworms Beetles Budworm Cabbage Looper Corn Earworm Cucumber Cutworms Grasshopper Leafhoppers Melonworm Pickleworm Plant Bug Rindworm Squash Bugs Squash Vine Borer Stink Bugs Tobacco Banks Grass Mite Carmine Mite Lygus Species Twospotted Spider Mite Whitefly | 0.04- 0.10 | 2.6-6.4 | <p>Apply in a minimum of 5 gallons of finished spray per acre by air or in a minimum of 20 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.</p> <p>Do not apply more than 0.3 lb active ingredient (19.2 ounces formulated) per acre per season.</p> <p>Do not make more than two applications after bloom.</p> <p>Do not make applications less than 7 days apart.</p> <p>Do not apply within 3 days of harvest.</p> |

LETTUCE, HEAD

| CROP | PEST | DOSAGE | | REMARKS |
|---------------|--|------------|----------|--|
| | | LB/AI/A | FL OZ/A | |
| Lettuce, Head | Aphids Armyworms Corn earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Loopers Salt Marsh Caterpillar Stink bug Species Tobacco Budworm Whitefly | 0.033-0.10 | 2.1-6.4 | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not make applications less than 7 days apart. A maximum of 0.5 lb. active ingredient may be applied per acre per season. |
| | Carmine Mite Lygus Species Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | Do not apply within 7 days of harvest. |

CANEberries

| CROP | PEST | DOSAGE | | REMARKS |
|--|---|-----------|---------|---|
| | | LB/AI/A | FL OZ/A | |
| Caneberries Including: Blackberries Bingleberries | Leafrollers Orange Tortrix Root Weevils | 0.05-0.10 | 3.2-6.4 | Apply by air or ground equipment using sufficient water to obtain full coverage of foliage. |

| | | | | |
|---|---|-------------|------------|--|
| <p>Dewberries Lowberries Marionberries Ollalieberries Youngberries Loganberries Raspberries</p> | <p>Raspberry Crown Borer Spider Mites</p> | <p>0.10</p> | <p>6.4</p> | <p>(Minimum of 10 gallons per acre by air and 50 gallons per acre by ground).</p> <p>One application may be made pre-bloom and a second application may be made post bloom.</p> <p>For Crown Borer, apply 0.1 lb al/a post harvest (fall) or pre-bloom (spring), as a drench application directed at the crown of plants in a minimum of 200 gallons water/ acre. Greater efficacy is observed at higher water gallonages (up to 400 gallons/a) or in an application prior to a significant rainfall event. Do not make both pre-bloom foliar and pre-bloom drench applications.</p> <p>Do not apply within 3 days of harvest.</p> <p>Do not exceed 0.2 lb al per acre per season.</p> |
|---|---|-------------|------------|--|

ARTICHOKE

| CROP | PEST | DOSAGE | | REMARKS |
|-----------|---|---------|---------|--|
| | | LB/AI/A | FL OZ/A | |
| Artichoke | Cribrate Weevil Artichoke Plume Moth | 0.10 | 6.4 | <p>Apply when pest population reaches damaging threshold and repeat as necessary to maintain control, but not more often than 15 day intervals.</p> <p>Application by ground: Apply a full cover spray in a minimum of 75 gallons of finished spray per acre.</p> <p>Application by air: Apply specified dosage in a minimum of 10 gallons per acre.</p> <p>Do not exceed 0.5 lb ai per acre per season. A 5-day preharvest interval must be observed.</p> |

HOPS

| CROP | PEST | DOSAGE | | REMARKS |
|------|---|-----------|---------|--|
| | | LB/AI/A | FL OZ/A | |
| Hops | Aphids Armyworms Cutworms Leafrollers Loopers | 0.06-0.10 | 3.8-6.4 | <p>Do not exceed 0.1 lb ai per acre per application.</p> <p>Do not exceed 0.3 lb ai per acre per season.</p> <p>A spray interval of 21 days between applications must be maintained.</p> |
| | Root Weevils | 0.05-0.10 | 3.2-6.4 | <p>A 14-day pre-harvest interval must be observed.</p> <p>Application by ground: For best results, full coverage is essential. Early season recommend 100-150 gallons of spray per acre. Late season recommend 200-250 gallons of spray per acre.</p> <p>For Root Weevil control, make a directed spray to the base of the plant. Spray up the vine 3 feet and the soil surface 1.5 to 2 feet on either side of the plant.</p> <p>Application by air for late season control of Twospotted spider mites: Apply no less than 6.4 oz. (0.1 lb ai) per application in a minimum of 10 gallons per acre.</p> <p>Use of ultra low volume (ULV) application on hops is prohibited.</p> |
| | Twospotted spider mite | 0.10 | 6.4 | |

PEARS

| CROP | PEST | DOSAGE | | REMARKS |
|------|------|---------|---------|---------|
| | | LB/AI/A | FL OZ/A | |

| | | | | |
|-------|--|-----------|-----------|---|
| Pears | Aphids Codling Moth Cutworms Green Fruitworm Leafhoppers Leafminers Leafrollers Lygus Species Plant Bugs Plant Bugs Plum Curcullo San Jose Scale (Crawlers) Stink Bugs Tarnished | 0.04-0.20 | 2.6-12.8 | <p>Application by ground: Apply as a dilute (minimum of 200 gallons of finished spray per acre) or concentrate (minimum of 50 gallons of finished spray per acre) spray in sufficient water to provide thorough coverage.</p> <p>Application by air: Apply the specified dosage in a minimum of 10 gallons per acre by air.</p> <p>Do not apply more than 0.5 pound active per acre per season with no more than 0.45 pound active per acre applied after petal fall.</p> |
| | Yellow Spider Mite | 0.06-0.20 | 3.8-12.8 | <p>Apply as necessary to maintain control using a minimum of 30 day spray interval.</p> |
| | Twospotted Spider Mite European Red Mite | 0.08-0.20 | 5.12-12.8 | <p>Apply up to 14 days prior to harvest.</p> <p>Do not graze livestock in treated orchards or cut treated cover crops for feed.</p> |

CITRUS

| CROP | PEST | DOSAGE | | REMARKS |
|--------|---|-----------|---------|--|
| | | LB/AI/A | FL OZ/A | |
| Citrus | Blue Green Citrus Root Weevil Brown Leaf Notcher Diaprepes Root Weevil Citrus Root Weevil Little Leaf Notcher | 0.25-0.50 | 16-32 | <p>Apply Capture 2EC by ground equipment to bare soil beneath citrus trees. Capture 2EC must be uniformly applied from the trunk to the drip line of tree; apply in a minimum of 40 gallons of dilute spray per acre. Greater spray volume should insure greater uniformity of coverage. A pre- and post-application irrigation may aid in the uniformity of coverage as well.</p> <p>Capture 2EC protects citrus tree roots from Diaprepes and other citrus root weevil feeding by forming a barrier which provides contact activity on newly hatched larvae (neonates). As citrus root weevil eggs hatch in new foliage, neonates fall to the soil surface beneath the tree and come</p> |

| | | | | |
|--|------------------------------|-----------|--------|--|
| | Asian cockroach Fire ants | 0.10-0.25 | 6.4-16 | <p>In contact with Capture 2EC as they attempt to burrow into the root zone. Disturbance of the soil beneath trees should be minimized.</p> <p>Timing of Capture 2EC applications is critical. Current information suggests that peak emergence of adult Diaprepes Weevil varies by citrus growing region and these emergence peaks can be dramatically affected by environmental factors, such as soil moisture. Typically, two peaks are observed for Diaprepes, first in spring then late summer or early fall. Citrus Root Weevils and Fuller Rose Beetle typically exhibit a single emergence peak in the spring. Brown and Little Leaf Notchers typically exhibit three emergence peaks, spring, summer and fall. Since emergence varies seasonally and by location, timing of Capture 2EC application can be accurately forecast by observing adults. Adults are most active early morning and late afternoon; numbers can be estimated by trapping throughout spring and summer (emergence periods). Egg laying will occur for 8 to 10 weeks following adult emergence from the soil; larval invasion of the soil will begin 2-3 weeks following adult emergence. It is critical to have the Capture 2EC soil barrier in place prior to drop of the neonates.</p> <p>Capture 2EC is one of several effective tools in an integrated pest management program for Citrus Root Weevils. Application of Capture 2EC should be used in conjunction with good cultural practices, biological control of larvae and foliar control of adults. Consult local university extension personnel for current information to protect citrus trees from Citrus Root Weevils and other pests.</p> <p>Additional Instructions: Apply to individual citrus resets, when not in solid planted rows, using hand-gun or shielded sprayer.</p> <p>Do not apply within 1 day of harvest.</p> <p>Peak emergence of Diaprepes root weevil generally occurs in the spring. Depending on weather conditions, a minor emergence of Diaprepes root weevil may also occur in the fall.</p> <p>If the citrus grove to be treated is in an area where weather conditions are conducive to primary emergence occurring in the spring, 32 fluid ounces formulated product should be used to obtain the longest residual management of Diaprepes root weevil. If the citrus grove to be treated is in an area where weather conditions will promote more than one peak of pest emergence, 16 fluid ounces formulated product can be applied early season and 16 fluid ounces formulated product can be applied later in the season.</p> <p>Do not apply through irrigation systems.</p> <p>Do not allow any application of Capture 2EC to contact fruit or foliage.</p> <p>Do not apply more than a total of 32 fluid ounces of formulated product (0.5 lb. a.i.) per acre per year.</p> <p>Apply the specified dosage in a minimum of 40 gallons of finished spray per acre.</p> <p>Ground application only. Do not apply by air.</p> |
|--|------------------------------|-----------|--------|--|

Spinach

| CROP | PEST | DOSAGE | | REMARKS |
|---------|---|------------|----------|--|
| | | LB/AI/A | FL OZ/A | |
| Spinach | Colorado Potato Beetle Tomato Pinworm Tomato Hornworm Armyworms Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Leafminers Loopers Pepper Weevil Thrips Whitefly | 0.033-0.10 | 2.1-6.4 | For control of whiteflies apply foliar treatments of Capture® Insecticide by ground or air at rates of up to 0.4 pt. (0.1 lb active) per acre at minimum 7-day intervals up to a maximum of 4 applications. Do not apply within 40 days of harvest. For control of fire ants apply Capture® Insecticide to the soil (at planting) or as a foliar treatment by ground or air at rates of up to 0.4 pt. (0.1 lb active) per acre at minimum 7-day intervals up to a maximum of 4 applications. Do not apply within 40 days of harvest. |
| | Broad Mite Banks Grass Mite Twospotted Spider Mite Carmine Mite Pacific Spider Mite Lygus species Fire Ants | 0.08-0.10 | 5.12-6.4 | Apply the specified dosage in 5-50 gallons of finished spray per acre by air or 10-50 gallons finished spray per acre by ground. |

GRAPES

| CROP | PEST | DOSAGE | | REMARKS |
|--------|--|-----------|---------|--|
| | | LB/AI/A | FL OZ/A | |
| Grapes | Cutworms Eastern grape leafhopper Grape berry moth Japanese beetles adults Variegated leafhopper Western grape leafhopper | 0.05-0.10 | 3.2-6.4 | Apply in a minimum of 10 gallons of finished spray by air or in a minimum of 25 gallons of finished spray with ground equipment. When applying by air, 1 to 2 quarts of emulsified oil may be substituted for 1 to 2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. |
| | Black vine weevil Glassywinged sharpshooter Twospotted spider mite | 0.10 | 6.4 | When pest pressure is moderate to severe, use higher rate. Do not apply more than 0.10 lb ai per acre per season. Do not apply within 30 days of harvest. |

OKRA

| CROP | PEST | DOSAGE | | REMARKS |
|------|---|------------|---------|--|
| | | LBS AI/A | FL OZ/A | |
| Okra | Aphids Armyworms Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Japanese Beetle (Adult), Stink bugs Leafminers Loopers Thrips Whitefly | 0.033-0.10 | 2.1-6.4 | Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft. Do not make applications less than 7 days apart. |

| | | | | |
|--|---|-----------|----------|--|
| | Broad Mite Carmine Mite Lygus Species Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | Do not apply more than 0.20 pound active ingredient per acre per season. Do not apply within 7 days of harvest. |
|--|---|-----------|----------|--|

CILANTRO, CORIANDER

| CROP | PEST | DOSAGE | | REMARKS |
|-----------------------|--|------------|----------|--|
| | | LBS A/A | FL OZ/A | |
| Cilantro Coriander | Aphids Beet Armyworm Cabbage Looper Cutworm Flea beetle Grasshoppers Leafminer Saltmarsh caterpillar Spotted Cucumber Beetle Thrips Whitefly | 0.033-0.10 | 2.1-6.4 | Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft. Do not make applications less than 7 days apart. Do not apply more than 0.50 pound active ingredient per acre per season. |
| | Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | Do not apply within 3 days of harvest. |

DRIED BEANS AND PEAS

| CROP | PEST | DOSAGE | | REMARKS |
|---|--|------------|---------|--|
| | | LBS A/A | FL OZ/A | |
| Dried cultivars of: Bean (Lupinus) Bean (Phaseolus) Field bean Kidney bean Lima bean (dry) Navy bean Pinto bean Tepary bean Bean (Vigna) Adzuki bean Blackeyed pea Catjang Cowpea Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean Broad bean (dry) Chickpea Guar Lablab bean Lentil Pea (Pisum) Field pea Pigeon pea | Aster Flea Beetle Grasshoppers Leafhopper Leafhoppers | 0.025-0.10 | 1.6-6.4 | Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. |
| | Alfalfa Caterpillar Aphids Bean Leaf Beetle Beet Armyworm Cloverworm Corn Earworm Corn Rootworm Adult Cucumber Beetles Cutworms European Corn Borer Fall Armyworm Imported cabbageworm Japanese Beetle Adult Leafminer Loopers Pea Leaf Weevil Pea Weevil Plant Bug Saltmarsh caterpillar Sap Beetle Southern Armyworm Stink Bugs Tarnished Plant Bug Thrips Tobacco budworm Webworms Western Bean Cutworm Whitefly Yellowstriped Armyworm | 0.033-0.10 | 2.1-6.4 | Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) to peas, or 0.3 lb. active ingredient (19.2 ounces formulated) to beans per acre per season. Do not apply within 14 days of harvest. Do not make applications less than 7 days apart. |

33
39

| | | | | |
|--|--|-----------|----------|--|
| | Banks Grass Mite Camine Mite Lygus Species Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | |
|--|--|-----------|----------|--|

LEAFY BRASSICAS

| CROP | PEST | DOSAGE | | REMARKS |
|---|---|------------|----------|--|
| | | LBS A/A | FL OZ/A | |
| Broccoli Raab Bok Choy Collards Kale Mizuna Mustard Greens Mustard Spinach Rape Greens | Aphids Armyworm Species Beetles Budworm Cabbageworm Caterpillar Corn Earworm Crickets Cucumber Cutworms Diamondback Flea Beetles Grasshoppers Ground Beetles Imported Japanese Beetle (adult) Leafhoppers Loopers Moth Saltmarsh Stink Bugs Thrips Tobacco Whitefly Wireworm (adults) | 0.033-0.10 | 2.1-6.4 | Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.4 lb. active ingredient per acre per season. Do not make applications less than 7 days apart. Do not apply within 7 days of harvest |
| | Banks Grass Mite Camine Mite Lygus Species Pacific Spider Mite Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | |

TUBEROUS AND CORM VEGETABLES

| CROP | PEST | DOSAGE | | REMARKS |
|----------------------------|--------------------------|--------------|------------|---|
| | | LBS A/A | FL OZ/A | |
| Potato | Corn wireworm | 0.30 | 19.2 | Capture may be applied as an in-furrow planting time treatment for the control of wireworms, rootworms and white grubs. Apply Capture at the rate of 0.3 pounds active per acre as an in-furrow spray or T-band spray at planting time. |
| Sweet potato | Tobacco wireworm | (at-plant) | (at-plant) | |
| Arracacha | | | | |
| Arrowroot | Southern potato wireworm | 0.05 to 0.15 | 3.2 to 9.6 | Capture may be applied as a lay-by treatment for the control of wireworms, rootworms and white grubs. Apply Capture to the drill area and cover with soil utilizing cultivation equipment set to throw soil to the drill area. Apply Capture as a banded spray over the row at a rate of 0.05-0.15 pounds active per acre (3.2 to 9.6 ounces formulated) in 10 gallons per acre of spray. |
| Chinese artichoke | Japanese beetle grubs | (lay-by) | (lay-by) | |
| Jerusalem artichoke | June beetle | | | |
| Edible canna | Sweetpotato flea beetle | 0.033 to | 2.1 to 6.4 | Capture may be applied as a foliar spray for the control of the adult life stages of flea beetles, click beetles (wireworms), cucumber beetles (rootworms), white fringed beetles and May/June beetles (white grubs). Apply Capture at the rate of 0.1 lbs active per acre (6.4 ounces formulated) in 10 gallons of spray by ground and 3 gallons of spray by air. |
| Cassava (bitter and sweet) | Cucumber beetle | 0.10 | (foliar) | |
| Chayote (root) | Sweetpotato weevil | (foliar) | | |
| Chufa | Banded Cucumber beetle | | | Do not make more than 2 foliar applications per season no sooner than 21 days apart. |
| Dasheen (taro) | Black flea beetle | | | |
| Ginger | Whitefringed beetle | | | |
| Leren | White grub | | | Do not apply more than 0.5 lb. active ingredient per acre per season, including soil application.. |
| Tanier | Sugarcane beetle | | | |
| Turner | Rootworms | | | |
| Yam bean | | | | Do not apply within 21 days of harvest |
| True yam | | | | |
| | | | | |

SOYBEAN

| CROP | PEST | DOSAGE | | REMARKS |
|---------|---|------------|----------|---|
| | | LBS A/A | FL OZ/A | |
| Soybean | Alfalfa Caterpillar Aphids Aster Leafhopper Bean Leaf Beetle Beet Armyworm* Cloverworm Corn Earworm Corn Rootworm Adult Cucumber Beetles Cutworms European Corn Borer Fall Armyworm Flea Beetle Grasshoppers Imported cabbageworm Japanese beetle Adult Leafhoppers Leafminer Loopers Mexican Bean Beetle Adult Pea Leaf Weevil Pea Weevil Plant Bug Saltmarsh caterpillar Sap Beetle Southern Armyworm Stink Bugs Tarnished Plant Bug Thrips Tobacco budworm* Webworms Western Bean Cutworm Whitefly Yellowstriped Armyworm | 0.033-0.10 | 2.1-6.4 | Apply foliar treatments in at least 10 gallons per acre at the rate of 6.4 fl. oz/ (0.1 lb active) per acre at a minimum of 30 days intervals. Do not apply more than 0.3 lb ai per acre per season. Do not apply within 18 days of harvest * Pyrethroid resistance is common for Beet Armyworm and Tobacco Budworm. Please consult your local or state agricultural authority to determine if resistance pest populations are in your area. If so refer the resistance management statement in the DIRECTION FOR USE section of this label. |
| | Lygus Species Whitefly Two Spotted Spider Mite | 0.08-0.10 | 5.12-6.4 | |

FRUITING VEGETABLES

| CROP | PEST | DOSAGE | | REMARKS |
|------|------|---------|---------|---------|
| | | LBS A/A | FL OZ/A | |

| | | | | |
|--|--|------------|----------|---|
| Eggplant Pepper (Bell & Non-Bell) Groundcherry Pepino | Armyworms Including: Beet Armyworm, Fall Armyworm, Southern Yellowstriped Armyworm Cabbage Looper Colorado Potato Beetle Corn Earworm Cucumber Beetle Cutworms European Corn Borer Flea Beetle Leafminers Loopers Pepper weevil Plant Bug Stink Bug Thrips Tomato Hornworm Tomato Pinworm Vegetable Leafminer Whitefly | 0.033-0.10 | 2.1-6.4 | Apply in a minimum of 2 gallons of <i>finished spray per acre by air or in a</i> minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the <i>finished spray. Thorough coverage is</i> essential to achieve control. Do not make applications less than 7 days apart. Do not apply more than 0.2 lb active ingredient (12.8 ounces formulated) per acre per season. Do not apply within 7 days of harvest |
| | Banks Grass Mite Broad Mite Carmine Mite Lygus Species Pacific Spider Mite Two Spotted Spider Mite Twospotted Spider Mite | 0.08-0.10 | 5.12-6.4 | |

| | | | | |
|---------------------|---|------------|----------|--|
| Tomato Tomatillo | Aphids Armyworms Including: Beet Armyworm, Fall Armyworm, Southern Yellowstriped Armyworm Bean Leaf Beetle Cabbageworm Carmine Mite Cloverworm Corn earworm Corn Rootworm Cucumber Beetles Cutworms Diamondback Moth European Corn Borer Flea Beetles Flea Hopper Grasshopper Japanese Beetle (Adult) Leafhoppers Loopers Lygus Species Melonworm Pea Weevil Pea Leaf Weevil Pickleworm Plant Bug Rindworm Salt Marsh Caterpillar Sap Beetle Seedpod Weevil Squash Bugs Stink bug Species Tobacco Budworm Tarnished Plant Bug Thrips Whitefly | 0.033-0.08 | 2.1-5.2 | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment, or 5 gallons of finished spray per acre by air. Thorough coverage is essential to achieve control. Do not make applications less than 10 days apart. A maximum of 4 applications may be applied per season. Do not apply within 1 day of harvest. |
| | Two Spotted Spider Mite | 0.08-0.10 | 5.12-6.4 | |

PEANUT

| CROP | PEST | DOSAGE | | REMARKS |
|--------|---|-----------|----------|--|
| | | LBS AI/A | FL OZ/A | |
| Peanut | Beet armyworm Corn earworm Cutworm Species Fall armyworm Grasshoppers Green cloverworm Leafhoppers Lesser cornstalk borer Loopers Rednecked peanut worm Southern armyworm Southern corn rootworm Stink bugs Threecornered alfalfa hopper Velvetbean caterpillar Yellowstriped armyworm | 0.033-0.1 | 2.1-6.4 | Apply foliar treatments in at least 10 gallons per acre at the rate of 6.4 fl. oz/ (0.1 lb active) per acre at a minimum of 14 days intervals. Do not apply more than 0.5 lb ai per acre per season. Do not apply within 14 days of harvest. Do not feed green immature plants and peanut hay to livestock. |
| | Aphids Spider mites Thrips Whitefly | 0.08-0.1 | 5.12-6.4 | |

ROOT CROPS

| CROP | PEST | DOSAGE | | REMARKS |
|---|--|-----------|----------|--|
| | | LBS AI/A | FL OZ/A | |
| Burdock, edible Carrot Celeriac Chervil, turnip rooted Chicory Ginseng Horseradish Parsley, turnip rooted Parsnip Radish Radish, oriental Rutabaga Salsify Salsify, black Salsify, Spanish Skirret Turnip | Aphids Beet armyworm Celery leaf tier Corn earworm Cross-striped cabbageworm Cutworms Diamondback moth European corn borer Fall armyworm Fire Ants Flea Beetles Green cloverworm Hornworms Imported cabbageworm Loopers Southern armyworm Spider mites Tobacco budworm Velvetbean caterpillar Whitefly Yellowstriped armyworm | 0.08-0.10 | 5.12-6.4 | Apply foliar treatments in at least 25 gallons per acre. Apply no more than once every 7 days. Do not apply more than 0.5 lb ai per acre per season. Do not apply within 21 days of harvest |
| Garden Beet | Aphids Fire Ants Flea Beetles Lepidopterous larvae Spider mites Whitefly | 0.08-0.10 | 5.12-6.4 | Apply foliar treatments in at least 25 gallons per acre. Apply no more than once every 7 days. Do not apply more than 0.4 lb ai per acre per season. Do not apply within 1 day of harvest |

MAYHAW

| CROP | PEST | DOSAGE | | REMARKS |
|--------|---------------|----------|----------|--|
| | | LB/AI/A | FLOZ/A | |
| Mayhaw | Plum Curculio | 0.08-0.1 | 5.12-6.4 | Apply foliar treatments in at least 28 gallons per acre. Apply no more than once every 7 days. Do not apply more than 0.2 lb ai per acre per season. Do not apply within 30 days of harvest |

Dealers Should Sell in Original Packages Only.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

FMC, Ammo, Capture, Mustang, U-Turn—trademarks of FMC Corporation

Ambush, Karate—trademarks of Zeneca, Inc.

Asana—trademark of E.I. duPont de Nemours & Company

Baythroid—trademark of Bayer Aktiengesellschaft

Danitol—trademark of Sumitomo Chemical Company, Ltd.

Scout Xtra—trademark of Hoechst Schering Agrevo S.A.

© 2003-2008 FMC Corporation All rights reserved.