

US ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (TS-767)  
WASHINGTON, DC 20460

EPA REGISTRATION NO.

279-3083

DATE OF ISSUANCE

JUN 14 1990

TERM OF ISSUANCE

Until Reregistration

NAME OF PESTICIDE PRODUCT

2000-0000 Insecticide

**NOTICE OF PESTICIDE:**  REGISTRATION  
 REREGISTRATION  
(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

[Faint text]  
 [Faint text]

225576 / 165  
2 / 19  
238515 / 166  
2 / 18  
266002 / 166  
3 / 18

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

*[Faint, mostly illegible text, likely bleed-through from the reverse side of the page]*

ATTACHMENT IS APPLICABLE

**BEST AVAILABLE COPY**

SIGNATURE OF APPROVING OFFICIAL

*[Handwritten signature]*

DATE

June 14 1990

Richards

d. Add the heading "REENTRY STATEMENTS" above the paragraph beginning "Do not apply this product in such a manner ... etc."

e. The If In Eyes statement should read "Flush eyes with plenty of water. Get medical attention if irritation persists."

f. Update the environmental hazards statement by changing "cleaning of equipment or disposal of wastes" to "disposing of equipment washwaters."

g. Delete "Shake well before measuring" since this product is a water soluble bag.

3. In your letters dated August 25, and October 13, 1989 you wished to modify the peach section of the label and to also add cucurbits. These changes are acceptable provided the ~~labeling reflects the changes listed above.~~

4. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca  
Product Manager 15  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

CODE

NET WEIGHT 0.41 Lbs.

**POUNCE<sup>R</sup> WSB INSECTICIDE**

**RESTRICTED USE PESTICIDE**

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

EPA Reg. No. 279-  
EPA Est. 33967-NJ-1

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

June 14, 1990

|                           |               |
|---------------------------|---------------|
| <b>Active Ingredients</b> | <b>By Wt.</b> |
| *Permethrin**.....        | 24.7          |
| Inert Ingredients.....    | 75.3          |
|                           | <u>100.0%</u> |

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act as  
amended, for the pesticide  
registered under EPA Reg. No.

279-3083

\*(3-Phenoxyphenyl)methyl(±)cis-trans 3-(2,2-dichloroethenyl)-2,  
2-dimethylcyclopropanecarboxylate  
\*\*cis/trans ratio: Max. 55% (±)cis and min. 45% (±)trans  
U.S. Patent No. 4,024,163

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

**AVISO**

See other panels for additional precautionary statements.

Precaucion Al Usuario: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicado ampliamente.

**STATEMENT OF PRACTICAL TREATMENT:**

If in eyes: Flush eyes with water for 15 minutes. Get medical attention if irritation persists.

If swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

For Emergency Assistance Call 716-735-3765

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans (and Domestic Animals)**

**Warning**

Do not get in eyes, on skin or on clothing. Wash thoroughly after handling.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply this product or allow it to drift to crops or weeds on which bees are actively foraging. Additional information may be obtained from your Cooperative Extension Service. Apply this product only as specified on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until sprays have dried.

Protective clothing means at least a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**WARNING**

Area treated with Pounce WSB insecticide on (insert date). Do not enter without appropriate protective clothing until sprays have dried (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

#### CHEMIGATION USE DIRECTIONS

Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation systems. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump motor must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

For Pounce WSB mixing instructions, see general instructions. Pounce WSB should be diluted in sufficient volume of water to ensure accurate application over the area to be treated. Add the number amount of water soluble bag(s) to the supply tank. Maintain sufficient agitation during both mixing and application to ensure that the bags dissolve and that there is uniformity of the supply tank suspension. Hydraulic or mechanical agitation is recommended. Pounce WSB should be applied continuously for the duration of the water application. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended.

#### STORAGE AND DISPOSAL

**PESTICIDE STORAGE:** Keep out of reach of children and animals. Store in original containers only. Store in a cool dry place and avoid excess heat. Do not store at temperatures below 32°F (0°C). Rough handling may cause breakage, especially at low temperatures. Allow to warm above 50°(10°C) before use. Do not allow inner bags to become wet during storage. Do not handle inner bag with wet hands or wet gloves. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716)735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter, or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

#### PESTICIDE DISPOSAL:

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL:

When all water soluble bags are used, the outer package should be clean and may be disposed of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. If the outer container contacts formulated product in any way, it must be triple rinsed with clean water. Add rinsate to the spray tank and dispose of the outer package as described above.

GENERAL INSTRUCTIONS

The product in this outer package is a wettable powder formulation of the insecticide permethrin packaged in a translucent water soluble bag. Do not allow the inner bag to become wet before adding to spray or nurse tank. Do not handle the inner bag with wet hands or wet gloves. Rough handling may cause breakage. Do not store at temperatures below 32°F (0°C). Allow to warm above 50°F (10°C) before use. Cooler water temperatures increase the time needed for the inner bag to dissolve completely.

Apply Pounce WSB when insects appear or feeding is noticed. The higher rate should be used as pest populations increase. Repeat the application as necessary to maintain control. Pounce WSB may be applied by both ground and aerial equipment. Use sufficient water to obtain full coverage. With the exception of crops listed below, rotational crops should not be planted within 60 days of last application.

Pounce WSB insecticide is to be diluted with water for spray application. Do not use strainers finer than 50 mesh size. Determine the number of water soluble bag(s) to make up necessary spray suspension.

---

Use a minimum spray volume of 5 gallons of water per acre. ~~Fill the~~ spray or nurse tank 1/2 full with water. Open the outer wrapper of this product and immediately dump required contents into spray tank. Allow the water soluble bag(s) to dissolve completely. Start hydraulic or mechanical agitation. Air agitation is not recommended. Mix thoroughly to fully disperse and suspend the wettable powder. Fill the spray tank with the required amount of water. Maintain agitation during storage in nurse or supply tank and during application.

Mix as needed; do not store diluted material.

**Almonds:** To control Navel Orangeworm, Peach Twig Borer--Use Pounce WSB at a rate of 2 to 4 bags (0.2 to 0.4 pounds active) per acre. Do not make more than 2 applications during hull split. Do not apply more than 2.0 pounds active ingredient per acre per season. Use sufficient water to obtain full coverage of foliage. Do not apply within seven days of harvest. Do not graze livestock in treated orchards.

**Apples:** To control Plum Curculio, Redbanded Leafroller, Rosy Apple Aphid, Spotted Tentiform Leafminer, Tarnished Plant Bug, White Apple Leafhopper, Oblique Banded Leafroller and Green Fruitworm--Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pounds active) per acre with ground equipment only. Apply in 25 to 400 gallons of finished spray. Do not apply more than 0.6 pound active ingredient per acre per season. Do not apply after petal fall.

**Asparagus:** To control Cutworms and Asparagus Beetle - Use Pounce WSB at the rate of 1 bag (0.1 pound active) per acre. Apply by ground equipment as needed. Do not apply more than 0.4 pounds active ingredient per acre per season. Do not apply within 3 days of harvest.

For post-harvest application - Use Pounce WSB at a rate of 1 bag (0.1 pound active) per acre. Apply to the fern stage of the asparagus plant after spear harvest when larval and adult stage of the Asparagus Beetle, Tarnished Plant Bug, Lygus Bugs and Japanese Beetle\* are present.

\*Not for control of this insect in California.

**Broccoli, Chinese Broccoli (gai lan, white flowering broccoli), Brussels Sprouts, Cauliflower:** To control Beet Armyworm, Cabbage Looper, Imported Cabbageworm and Diamondback Moth--Use Pounce WSB at a rate of 1 bag (0.1 pound active) per acre. Apply in a minimum of 5 gallons of finished spray per acre by aircraft or 20 gallons per acre by ground equipment. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply within 1 day of harvest.

**Cabbage, Chinese Cabbage (tight-heading varieties only):** To control Beet Armyworm, Cutworms, Cabbage Looper, Diamondback Moth and Imported Cabbageworm--Use Pounce WSB at a rate of 1 bag (0.1 pound active) per acre. Apply in a minimum of 5 gallons of finished spray per acre by aircraft or 20 gallons per acre with ground equipment. Do not apply more than 1.0 pound of active ingredient per acre per season. Do not apply within 1 day of harvest.

**Cantaloupes:** To control Aphids, Leafminers, Squash Bug, and Whiteflies--Use Pounce WSB at a rate of 2 bags (0.2 pound active) per acre. To control Cucumber Beetle, Cutworms, Melon Worm, Pickleworm and Squash Vine Borer--Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply a minimum of 5 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active per acre per season. Do not apply within one day of harvest.

**Celery: Florence fennel (sweet anise, sweet fennel, finocchio):** For general use directions refer to the Leafy vegetable crop grouping.

**Cherries:** To control Lesser Peach Tree Borer, Rose Chafer, Green Fruitworm, Red-banded Leafroller, Plum Curculio and Tarnished Plant Bug--Use Pounce WSB at a rate of 1 bag (0.1 pounds active) per 400 gallons of water. Do not exceed 400 gallons per acre. Apply when insects appear. East of the Rockies, do not exceed 6 applications per season, with no more than 4 applications after petal fall. West of the Rockies, do not apply more than 4 applications per season, with no more than 3 applications after petal fall. Do not apply within 3 days of harvest. Do not graze livestock or feed crop forage from treated orchards.

**Chrysanthemums (Greenhouse\*)**: For control of *Liomyza* Leafminer flies - Use Pounce WSB at a rate of 5 bags (0.5 pound active) per 100 gallons. Avoid spraying the blooms. Pounce may be applied on a weekly schedule. Caution: Cultivars may vary in sensitivity and a small number of plants should be treated to determine plant safety prior to commercial use.  
\*Not for use in California

**Conifers (Container and Field Grown)**: For control of Nantucket Pine Tip Moth - Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Begin application when the adults appear and repeat at 5 to 7 day intervals or as needed throughout the season.

**Cotton: Preemergent Use**: To control Cutworms - Use Pounce WSB insecticide at a rate of 1 to 2 bags (0.1 to 0.2 pounds active) per acre. Apply prior to planting or preemergent to crop.

**Foliar Use**: To control Beet Armyworm (California and Arizona only), Bollworm, Cabbage Looper, Cotton Fleahopper, Cotton Leaf Perforator, Cutworms, Lygus Bugs, Pink Bollworm, Tarnished Plant Bug and Tobacco Budworm--Use Pounce WSB at a rate 1 to 2 bags (0.1 to 0.2 pound active) per acre. To control Boll Weevil, alternate Pounce WSB at 1 bag (0.1 pound active) per acre with Pounce 25 WP at 3.2 ounces (0.05 pound active) per acre every 3 to 4 days until control is achieved.

Use a *minimum* of 5 gallons of water per acre by air or ground application. Do not apply more than 3.0 pounds active ingredient per season. Do not apply within 14 days of harvest. Do not graze or feed cotton forage.

**Eggplants**: To control Colorado Potato Beetle - Use Pounce WSB at the rate of 2 bags (0.2 pound active) per acre. To control Vegetable Leafminer - Use Pounce WSB at the rate of 1 to 2 bags (0.1 to 0.2 pound) active per acre. Do not apply more than 3.2 pounds active ingredient per acre per season. Do not apply within 3 days of harvest.

**Field Corn, Popcorn, Field Corn Grown for Seed**: Preemergent Use: To control Armyworm and Cutworms - Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre in the time period from five days prior to planting up to emergence of the crop. Apply as a broadcast spray with ground equipment in a minimum of 20 gallons of finished spray per acre.

**Foliar Use**: For control of Cutworms, European Corn Borer, Flea Beetle, Southwestern Corn Borer, Stalk Borer, Corn Rootworm Beetle and Armyworm - Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre prior to ear formation. Ear formation occurs when pollination is completed as evidenced by initiation of brown silk in

field to be treated. Apply a minimum of 5 gallons of finished spray per acre by air or 10 gallons per acre with ground equipment. When treating for stalk borer, Pounce WSB must be applied when or shortly before the stalk borer larvae are moving into the corn from surrounding weeds and grasses. Mowing or burn-down herbicide is suggested to initiate movement.

Filberts: To control Filbertworm and Oblique Banded Leafroller-Use Pounce WSB at a rate of 2 to 4 bags (0.2 to 0.4 pound active) per acre. For full coverage application apply 1 bag (0.1 pound active) per 100 gallons (based on 400 gallons finished spray per acre), spray to runoff; or, for low volume application apply 2 to 4 bags (0.2 to 0.4 pound active) per acre (50 to 200 gallons finished spray per acre).

For aerial application apply 2 to 4 bags (0.2 to 0.4 pound active) in a minimum of 15 gallons of finished spray per acre. Apply when insects appear. Do not apply more than 1.6 pounds active ingredient per acre per season. Do not apply within 14 days of harvest. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock.

Head Lettuce: For general use directions refer to the Leafy Vegetable crop grouping.

Horseradish: To control Imported Crucifer Weevil (*Baris lepidii*). For foliar application -- Use Pounce WSB at a rate of 2 bags (0.2 pound active) per acre. Apply by ground equipment in a minimum spray volume of 20 gallons per acre. Make up to 3 foliar applications as needed to control weevil adults during oviposition.

Do not apply within 22 days of harvest.

Leafy Vegetables (except Brassica): Amaranth; arrugula; celery; celtuce; chervil, corn salad, chrysanthemum (edible-leaved and garland); cress, (garden and upland); dandelion; dock; endive; fennel; lettuce (leaf and head); orach; parsley; purslane (garden and winter); rhubarb; spinach; Swiss chard: To control Alfalfa Looper, Beet Armyworm, Cabbage Looper, Corn Earworm, Cutworms, European Corn Borer, Fall Armyworm, Green Cloverworm, Leafhoppers, Southern Armyworm, Tobacco Budworm and Vegetable Leafminer - Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply every 3 to 5 days or as needed by air or ground. Use sufficient water to obtain full coverage of foliage. Do not apply more than 2.0 pounds active ingredient per acre per season. Can be applied up to 1 day of harvest. Do not graze treated areas or feed crop refuse to livestock.

Ornamental Nursery Stock (Field Grown): To control Beet Armyworm, Cabbage Looper, Citrus Thrips, *Heliothis* spp., Leafhoppers, Leafminers and Whiteflies - Use 1 to 2 bags (0.1 to 0.2 pound active) per 100 gallons of water. Pounce WSB may be used to control specified pests on non-edible ornamentals and non-bearing plants of fruiting species.

**Caution:** Pounce WSB has demonstrated excellent plant safety; however, not all species and varieties have been tested. Before treating large numbers of plants of a particular variety, treat a few plants and observe prior to full-scale application. Marginal leaf burn may occur on Salvia, Dieffenbachia and Pteris Fern. Application to blooming plants may cause browning of petals.

**Peaches:** To control Lesser Peach Tree Borer, Peach Twig Borer, Rose Chafer, Plum Curculio, Tarnished Plant Bug, Green Fruitworm, Oriental Fruit Moth--Use Pounce WSB at a rate of 1 bag ( 0.1 pound active) per 100 gallons of water. Do not exceed 400 gallons per acre. Apply when insects appear. Make applications with ground equipment, using sufficient water to obtain uniform coverage. Do not apply within 7 days of harvest. Do not make more than 8 applications per season. Do not graze livestock in treated orchards.

**Pears (Dormant through Delayed Dormant):** To control Pear Psylla-- Apply Pounce WSB at a rate of 2 to 4 bags (0.2 to 0.4 pound active) per acre. Apply during the dormant through delayed dormant growth periods only. Apply in 5 to 20 gallons of finished spray per acre by aircraft and 25 to 400 gallons per acre by ground equipment. Do not apply more than 0.8 pound active ingredient per season. Do not graze orchard floor.

**Pears (Summer):** To control Pear Psylla, Green Fruitworm, Codling Moths--Use Pounce WSB at a rate of 2 to 4 bags (0.2 to 0.4 pound active) per acre. Use in sufficient water to achieve runoff and apply by ground equipment. Do not apply more than 1.2 pounds active ingredient per acre per season. Do not apply within 14 days of harvest.

**Peppers (Bell):** To control Vegetable Leafminer, Cabbage Looper, Cutworms, Flea Beetle, Corn Earworm, Pepper Weevil -- Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. To control European Corn Borer -- Use Pounce WSB at a rate of 2 bags (0.2 pound active) per acre. Do not apply more than 1.6 pounds of active ingredient per acre per season. Do not apply within 3 days of harvest.

**Pine Seed Orchards:** to control Coneworms, Seed Bugs -- Use Pounce WSB at the following rates:

For high volume sprayers: Use 3 bags in 100 gallons of water (0.025% dilution by weight). Apply 5 to 10 gallons of finished spray per tree.

For low volume sprayers: Use 16 bags in 100 gallons of water (0.125% dilution by weight). Apply 100 gallons per acre.

Apply first application within 1 week of female flower closure or peak pollen flight for webbing coneworm control. For other coneworms and seed bugs, apply first application within 30 days following female flower closure. Repeat applications at 4 week intervals, but do not apply more than 6 applications. Do not graze or harvest cover crop. Avoid contact with open water.

**Pistachios:** To control Peach Twig Borer and Navel Orangeworm -- Use Pounce WSB at a rate of 2 to 4 bags (0.2 to 0.4 pound active) per acre. Do not apply more than 0.8 pound active ingredient per acre per season. Do not apply after 10% hull split. Nuts may be harvested on the day of application. Do not graze livestock in treated orchards.

**Potatoes:** To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Cutworms, European Corn Borer, Potato Flea Beetle, Potato Leafhopper, Potato Psyllid, Potato Tuberworm, Aster Leafhopper and Tarnished Plant Bug. Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply in a minimum of 5 gallons of finished spray per acre by air or 20 gallons of finished spray per acre using ground equipment. Do not apply more than 2.4 pounds active ingredient per acre per season. Do not apply within 7 days of harvest. Do not graze or feed potato forage.

**Pumpkins:** To control Aphids, Leafminers, Squash Bug, and Whiteflies--Use Pounce WSB at a rate of 2 bags (0.2 pound active) per acre. To control Cucumber Beetle, Cutworms, Melon Worm, Pickleworm and Squash Vine Borer--Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply a minimum of 5 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 0.8 pound active per acre per season. Do not apply within one day of harvest.

**Roses (Field Grown):** For control of *Heliothis* spp., use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply in 5 gallons of finished spray per acre by ground or by air application.

**Roses (Greenhouse):** For control of Beet Armyworm, Cabbage Looper and Omnivorous Leafroller - Use Pounce WSB at a rate of 2 bags (0.2 pound active) per 100 gallons of water.

**Caution:** Varieties may vary in their sensitivity to Pounce 25 WSB, and a small number of plants should be treated under local conditions to determine plant safety prior to commercial use.

**Soybeans:** To control Cabbage Looper, Green Cloverworm, Mexican Bean Beetle, Soybean Looper, Velvetbean Caterpillar, Bean Leaf Beetle and Potato Leafhopper--Use Pounce WSB at 1 bag (0.1 pound active) per acre. To control Beet Armyworm and Corn Earworm--Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply a minimum of 5 gallons of finished spray per acre by air or 5 gallons with ground equipment. Do not apply more than 0.4 pound active ingredient per acre per season. Do not apply within 60 days of harvest. Do not graze or feed soybean forage or hay.

**Spinach:** For general use directions, refer to the Leafy Vegetable crop grouping.

**Sweet Corn:** To control Aster Leafhopper, Corn Earworm, European Corn Borer, Fall Armyworm, Flea Beetle, Corn Rootworm Beetle\* and cutworms -- Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply every 3 to 5 days or as needed. Do not apply more than 1.2 pounds active ingredient per acre per crop season. Do not apply within 1 day of harvest of ears or forage or livestock grazing.

\*Not for control of this insect in California.

**Sweet Corn (Florida Only) fresh market use only:** To control Aster Leafhopper, Cutworms Corn Earworm, European Corn Borer, Fall Armyworm, Southern Armyworm -- Use 1 to 2 bags (0.1 to 0.2 pound active) per acre. Apply every 3 to 5 days as needed.

Do not apply more than 3.25 pounds active ingredient per acre per crop season. Do not graze livestock in treated areas or harvest for forage.

**Tomatoes (Only for Application to Tomatoes Grown in Florida for Final Marketing as Fresh Tomatoes):** To control Beet Armyworm, Cabbage Looper, Colorado Potato Beetle, Granulate Cutworm, Hornworms, Vegetable Leafminer, Southern Armyworm, ~~Tomato Fruitworm and Tomato~~ Pinworm -- Use Pounce WSB at a rate of 1 to 2 bags (0.1 to 0.2 pound active) per acre. Do not apply more than 1.2 pound active ingredient acre per season. Can be applied up to the day of harvest. Do not apply to cherry tomatoes or other varieties which produce fruit less than one inch in diameter.

**Walnuts:** To control Navel Orangeworm, Codling Moth and Walnut Husk Fly--Use Pounce WSB at a rate of 2 to 4 bags (0.2 to 0.4 pound active) per acre. For full coverage application apply 1 bag (0.1 pound active) per 100 gallons (based on 400 gallons finished spray per acre), spray to runoff; or, for low volume application apply 2 to 4 bags (0.2 to 0.4 pound active) per acre (50 to 200 gallons finished spray per acre). For aerial application apply 0.2 to 0.4 pound active in a minimum of 20 gallons of finished spray per acre. Apply when insects appear. Do not apply more than 1.6 pounds active per acre per season. Do not apply within 1 day of harvest. Do not graze livestock in treated orchards or feed cover crops from treated orchards to livestock.

**Premises Spray**  
For agricultural use only.

Spray directly to walls and ceiling as residual surface treatment only. Do not treat manure or litter. Avoid contamination of feed and water. Do not apply directly to livestock or poultry.

Shake well before measuring.

| For Application in   | Target Insects  | Method of Applic. | Dilute                 | Applic. Rate                      |
|--|---|-------------------|------------------------|-----------------------------------|
| Dairies, Barns, Feedlots, Stables, Poultry Houses, Swine and Live-stock Houses | House Flies, Stable Flies and other Manure Breeding Flies. Also aids in the reduction of Cockroaches, Mosquitoes and Spiders. | Sprayer           | 1 bag to 11 gal. water | 1 gal. per 750 sq. ft. of surface |

Re-treat as necessary, but not more often than once every 2 weeks. The use of any residual fly spray should be supplemental with proper manure management and general sanitation to reduce or eliminate fly breeding site.

RATE CONVERSION CHART

| Pounds Active Per Acre | Water Soluble Bags Per Acre |
|------------------------|-----------------------------|
| 0.1                    | 1                           |
| 0.2                    | 2                           |
| 0.3                    | 3                           |
| 0.4                    | 4                           |
| 0.5                    | 5                           |

**Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

## **POUNCE WSB**

Cucurbit Vegetables: Balsam pear (Bitter Melon); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourds, edible; melons, including hybrids such as canteloupe, casaba, crenshaw, honeydew melons, honey balls, mango melon, muskmelon, Persian melon; pumpkin; squash, summer and winter; watermelon, including hybrids: To control leafminers and squash bug -- Use Pounce WSB at a rate of 2 bags (0.20 pound active) per acre. To control cabbage looper, cucumber beetle (adults), cutworms, melon worm, pickleworm, plant bugs (including Lygus and stink bugs), rindworms and squash vine borer -- Use Pounce WSB at a rate of 1 to 2 bags (0.10 to 0.20 pound active) per acre. Apply a minimum of 5 gallons of finished spray per acre by air or 20 gallons of finished spray per acre with ground equipment. Do not apply more than 1.6 pounds active ingredient per acre per season. Pounce WSB may be applied up to harvest.

Canteloupes: For general use directions, refer to the Cucurbit Vegetable crop grouping.

Pumpkins: For general use directions, refer to the Cucurbit Vegetable crop grouping.