



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

SEP 15 1994

CHRISTINE A. DIVELY
FMC CORPORATION
ACG SPECIALTY PRODUCTS
1735 MARKET STREET
PHILADELPHIA, PA 19103

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 04/15/94 Response to PR Notice 93-7
EPA Reg. No. 279-3106
COMMAND 50 WP HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division

Christine A. Dively
FMC CORP.
ACG SPECIALITY PRODUCTS
1735 MARKET STREET
PHILADELPHIA PA 19103

Comment for: EPA Reg Nr.279-3106
COMMAND 50 WP HERBICIDE

The following specific comments pertain to your WPS
labeling submission concerning the product
cited above:

Correct the typographical errors circled on your proposed
label.

Code 1331

Net Contents

COMMAND 50 WP

Herbicide

For Agricultural or Commercial Use Only
NOT FOR SALE OR USE IN CALIFORNIA

EPA Reg. No. 279-3106 FPA Est. No. 279

Active Ingredient

*Clomazone: 2-(2-Chlorophenyl)methyl-4,
4-dimethyl-3-isoxazolidinone 50.0%

Inert Ingredients 50.0%
100.00%

U.S. Patent No. 4,405,357

KEEP OUT OF REACH OF CHILDREN WARNING

AVISO

usted no entiende la etiqueta, busque a alguien para que se le explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If swallowed: Call a physician or Poison Control Center. Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash skin with plenty of soap and water. Get medical attention.

Emergency Assistance Call (800) 331-3148

Use other panels for additional precautionary information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning:

May be fatal if inhaled. Do not breathe dust. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing.

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAY 26 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

279-3106

FMC Corporation
Agricultural Chemical Group
Philadelphia, PA 19102

Personal Protective Equipment

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- For exposure in enclosed areas, a respirator with either an organic vapor-removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G)
- For exposure outdoors, dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statement: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6)), the handler's PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

TABLE OF CONTENTS

Read all information on pages 1-4 before referring to specific crop use.

	Page
Ingredient	1
Statements of Practical Treatment	1
Precautionary Statements	1
Directions for Use	1
Storage and Disposal	1
Sprayer Cleanup	2
General Mixing Instructions	2
Application Precautions	2
General Application Instructions	3
Command Geographic Areas	4
PUMPS	5
SUCKLING ANIMALS	5

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 50 WP herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

IMPORTANT

IF YOU DO NOT PERVE THE APPLICATION PRECAUTIONS SECTION OF THIS LABEL MAY RESULT IN INJURY TO DESIRABLE VEGETATION

- Desirable plants including some species of trees, shrubs, flowers, agronomic crops, and fruits and vegetables are sensitive to Command 50 WP herbicide.
- Foliar contact with spray drift or vapors may cause whitening of sensitive plants. Symptoms are generally temporary in nature but may persist on some plants.
- Carryover injury to approved rotational crops may result under extremely dry conditions. Choice of rotational crop hybrid, soil factors, and choice of other crop protection chemical, can impact the risk of injury to approved rotational crops. Refer to Rotational Cropping Precautions and Guidelines.

SPRAYER CLEANUP

Do not drain or flush equipment on or near desirable trees or other plants, or in areas where their roots may extend or in locations where the chemical may be washed or move into contact with their roots. Do not contaminate any body of water including irrigation water that may be used on other crops. Carefully follow sprayer clean up instructions noted below to prevent spray tank residues from damaging other crops.

Sprayer equipment should be thoroughly rinsed to remove residues of herbicide that might injure other subsequently sprayed crops. The steps below should be followed for the thorough rinsing of spray equipment following applications of Command 50 WP herbicide or tank mixes of Command 50 WP with other labeled products.

- 1) Drain any remaining spray solution from tank, pump, hoses and boom and discard in an approved manner (See Note that follows)

- 2) Clean tank and fittings by:
 - Thoroughly flush down the inside walls of the spray tank with a quantity of water equal to 1/2 of the total tank capacity and operating the pump to circulate this solution through the sprayer system for 15 minutes.
 - Washing down the outside surfaces of equipment.
 - Removing nozzle tip and screen from end nozzle in each boom section and allowing several gallons of rinsate solution to flush completely through boom (collect rinsate while flushing).
- 3) Thoroughly drain remaining rinsate solution from tank, pump and hoses. Combine with boom flushings and dispose of all rinsates from this first rinsing in an approved manner (see Note that follows).
 - When switching from water dilutions to applications utilizing crop oil or liquid fertilizer as a carrier, a small volume of crop oil or liquid fertilizer should be flushed through the tank, pump, hoses, and boom prior to the next use. Dispose of crop oil or liquid fertilizer rinsate in an approved manner (see Note for local, state and federal guidelines).
- 4) Remove the remaining nozzle tips, and screens and the line filter and wash in a pail of warm soapy water, thoroughly rinse and replace.
- 5) Hose down the inside walls of the spray tank a second time and circulate this solution using the same procedure as noted in #2 above.
- 6) If the next use of the sprayer will be for applying a preemergent or preplant incorporated pesticide on any crop for which Command 50 WP is registered, rinsate from this second rinsing may be utilized by diluting with the water for the next pesticide load;

HOWEVER

If the next use of the sprayer will be a postemergence applied pesticide on any crop, drain rinsate solution from this second rinsing. Retain rinsate solution for use only with a soil incorporated pesticide to be applied on a crop for which Command 50 WP is registered. Refill tank (after draining second rinsate solution) in accordance with post-emergence product label directions.

NOTE: Dispose of excess spray mixture and/or rinsate from first tank rinsing by application to cropland as described on this label. If excess spray mixture and/or rinsate from first rinsing cannot be disposed of according to label instructions, dispose of in compliance with local, state and federal guidelines. Contact your state pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance.

GENERAL MIXING INSTRUCTIONS

Care must be taken when mixing Command 50 WP herbicide. Avoid mixing in areas adjacent to desirable plants (See Section headed "IMPORTANT").

Command 50 WP Alone: Mix Command 50 WP with water or liquid fertilizer in the following manner. Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 50 WP, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

Tank Mixtures: Fill spray tank one-fourth to one third full with water, with agitator operating add the recommended amount of ingredients using the following order: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's). Mix thoroughly and fill tank one-half full continuing agitation. Add Command 50 WP herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Where use of a surfactant is recommended, add as the last ingredient to the spray tank. Maintain agitation during filling, mixing and application. When using drift reducing agents, follow specific product label instructions for order of addition to spray tank.

Fertilizer Spray Mixtures: Applications of Command 50 WP alone, or with recommended tank mixtures, in conjunction with fertilizer solutions, may be used unless use directions specifically state otherwise. Small quantities should be tested for compatibility by the following procedure before mixing tankful quantities.

- 1) Put 1 pint of fertilizer solution in a quart jar

2) Add the appropriate amount of herbicide based on the table below if more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's)

Herbicide	Rate/Acre	Amount to Be Added Per Pint of Fluid Fertilizer*
Wettable Powders or Dry Flowables (Dry Granules)	1/2 pound	1/4 level teaspoon
	1 pound	1/2 level teaspoons
	2 pounds	3 level teaspoons
Emulsifiable Concentrates or Flowables	3 pounds	4 1/2 level teaspoons
	1 pint	1/2 level teaspoon
	1 quart	1 level teaspoon
Flowables	2 quarts	2 level teaspoons
	3 quarts	3 level teaspoons

*Based on a spray volume of 25 gpa. For lower or higher spray volumes, adjust fertilizer quantity accordingly.

3) Close jar and shake well.

4) Watch mixture for several seconds, again in 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed.

If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below:

1) **Command 50 WP Alone:** Dilute with 2 parts of water and add slowly to spray tank of fertilizer with agitator running. Maintain agitation during application.

2) **Command 50 WP plus Tank Mixtures:** Dilute individual products with two (2) parts of water, then add to the spray tank of fertilizer, while maintaining agitation, using the following order—slurry of dry formulations (wetable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Continue agitation during application.

APPLICATION PRECAUTIONS

Geographic Restrictions: Command 50 WP must be soil incorporated in the following states, unless there is an approved SLN (24C) for counties within these states:

Connecticut	Maryland	Ohio
Delaware	Massachusetts	Pennsylvania
Florida	Michigan	Rhode Island
Georgia	Minnesota	Vermont
Indiana	New Hampshire	Virginia
Iowa	New Jersey	West Virginia
Kentucky	New York	Wisconsin
Maine		

For Soil Incorporated Applications: (Preplant Incorporated): See Geographic Restrictions above.

Plants such as the following may show symptoms of foliar whitening or yellowing if contacted by Command 50 WP herbicide. It is recommended that, prior to application, adjacent properties be checked and that spraying within 100 feet of such desirable plants be avoided.

Trees (Deciduous)	Trees (Evergreen)
Apple (inc. fruit & ornamental types)	Fir species
Ash (Green, White, Mountain)	Spruce species
Basswood	Shrubs & Vines
Boxelder	Azalea
Catalpa	Burningbush
Cherry (inc. fruit & ornamental types)	(Winged Euonymus)
Cottonwood	Grape
Elm	Honeysuckle
Ginkgo	Roses
Hackberry	Yew
Mulberry	Agronomic Crops
Peach	Alfalfa
Pear (inc. fruit & ornamental types)	Oats
Pecan	Vegetables and Flower plants
Poplar	
Russian olive	Others
Tree-of-Heaven	Ferns
Tulip tree	Herbs
Walnut trees	Strawberry
Willow species	Raspberry
	Blackberry

Apply Command 50 WP herbicide only to surfaces that will be soil incorporated. Do not apply Command 50 WP herbicide to non-field areas including fence rows, waterways, ditches, and roadsides.

Do not apply Command 50 WP within 1,000 feet of the areas listed below:

Towns and Subdivisions
Commercial Vegetable Production*
Commercial Nurseries
Commercial Greenhouses

*except pumpkins, succulent peas, sweet corn


For Preemergent Applications: See Geographic Restrictions above.

Except as restricted above (see Geographic Restrictions), Command may be used as a preemergent soil applied treatment without incorporation, however, additional precautions, beyond those required for a soil preplant incorporated treatment, must be taken as follows:

Do not apply Command 50 WP within 1,500 feet of the following areas: Towns and Subdivisions, Commercial Fruit or Vegetable Production, Commercial Greenhouses or Nurseries.

Prior to application, adjacent properties must be checked, and spraying within 1,500 feet of desirable plants must be avoided.

*except pumpkins, succulent peas, sweet corn



*Does Not Apply to Fallow Land Applications, or Counties for which an Approved SLN (24C) is on File

SPRAY DRIFT PRECAUTIONS

Non target spray drift of Command 50 WP herbicide should be avoided to prevent whitening of desirable vegetation. Drift is influenced by many factors which include wind speed, spray pressure, particle size, nozzle type, and boom height.

- Do not apply when weather conditions favor drift. If wind speeds exceed 10 miles per hour, a drift reducing additive must be used.
- A minimum spray volume of 10 gallons per acre is recommended with appropriate nozzle types and sizes that produce coarser sprays.
- The use of agriculturally approved drift reducing additives is recommended for application volumes of 15 to 40 gallons per acre when spraying in the proximity of desirable plants (see list above).
- The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre.
- Use minimum nozzle pressure and boom height while maintaining uniform spray pattern.
- Do not exceed 40 psi spray pressure.
- Selection and proper use of spray equipment is critical in minimizing spray drift. The following table suggests pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types.

Suggested Nozzle Types, Minimum Size and Recommended Pressure Ranges for Minimizing Drift

Nozzle	Pressure Range (PSI)	Minimum Flow Rate Within Pressure Range (GPM)	Minimum Nozzle Size
Flat-fan	15-30	0.3	#4*
LP-flat-fan	10-25	0.3	#3
Even flat-fan**	15-30	0.3	#4
Flood	10-25	0.3	#2.5*
Whirl-chamber	5-20	0.3	#5
Raindrop	15-40	0.15	#2
Wide angle			
full cone	15-40	0.3	#5

* Refers to tip number such as 8004 or LF 2.5

** Recommended for banded application

ROTATIONAL CROPPING PRECAUTIONS: Under some conditions, temporary whitening or yellowing of leaves may occur on approved rotational crops where undesirable soil residues of Command 50 WP herbicide exist.

Under abnormal conditions, carryover injury to rotational crops can occur. The following factors can contribute to increased risk of injury to rotational crops:

- 1) Exceeding label recommended rates.
- 2) Overapplication resulting from use of worn nozzles, excessive overlapping of spray swaths, failing to shut off spray booms when turning (end row areas), or slowing or stopping sprayer.
- 3) Soil with pH less than or equal to 5.9.
- 4) Extreme dryness in the four months following application.
- 5) Use of organophosphate soil insecticides followed by use of some postemergence corn herbicides.
- 6) Choice of rotational crop hybrid.

Additional recommendations to prevent rotational crop injury may be provided in the form of service bulletins for locations where risk of injury is significantly increased due to extremely dry conditions.

Refer to Rotational Guidelines and replanting instructions of specific crops for additional crop planting information.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 50 WP herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

GENERAL APPLICATION INSTRUCTIONS

Do not apply aerially or through irrigation equipment.

Ground Applications

Broadcast or Banded Applications: Apply Command 50 WP alone or in tank mix combination, by ground equipment using a finished spray volume of 10 to 40 gallons of water per acre. **NOTE:** The use of an agriculturally approved drift reducing additive is required at finished spray volumes of 10 to 15 gallons per acre. Use nozzles suitable for broadcast boom or banded application of herbicides. Coarse sprays are less likely to drift out of the target area than fine sprays. See "APPLICATION PRECAUTIONS" section for specific recommendations to reduce spray drift. For Command 50 WP tank mixtures with wettable powder or dry flowable formulations, nozzle screens and strainers should be no finer than 50 mesh.

labeled Application. Calculate the rate and volume required by using the following formula:

$$\frac{\text{Band width in inches} \times \text{Broadcast Rate Per Acre}}{\text{Row width in inches}} = \text{Band rate per acre}$$

$$\frac{\text{Band width in inches} \times \text{Broadcast Volume Per Acre}}{\text{Row width in inches}} = \text{Band volume per acre}$$

Follow broadcast minimum gallonage requirements adjusted to band width. Do not exceed 40 psi spray pressure

Refer to Spray Drift Precautions for suggested pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types. For banded applications an even flat-fan nozzle is recommended.

Soil Incorporated Application: Apply Command 50 WP to the soil surface and uniformly incorporate. Select equipment suited for proper incorporation of Command 50 WP applied prior to or after planting. Shallow incorporation normally results in better weed control than a deeper incorporation.


In states where soil incorporation is required (see Geographical Restrictions) application and immediate incorporation to a maximum depth of 3 inches is required unless the soil is dry. On dry soils, incorporation to a maximum depth of 3 inches must be completed within 8 hours of Command 50 WP herbicide application.

Soil must be in good tilth to allow for thorough mixing of the soil. Application on wet soils will increase the potential for off-site movement of Command 50 WP herbicide vapors and may result in poor soil incorporation and unsatisfactory weed control.

Cultivation After Planting

Soil treated with Command 50 WP may be shallow-cultivated, rotary-hoed or hand-hoed without reducing the weed control activity of Command 50 WP. Do not cultivate deeper than the Command 50 WP treated layer of soil since this may bring untreated soil to the surface and poor weed control may result.

COMMAND GEOGRAPHICAL AREAS For Rates, Weed Control, and Crop Rotation



* Southern area includes Missouri bootheel, which incorporates the following counties: Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott, and Stoddard

NORTHERN AREA ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur.

Refer to sections headed "Rotational Cropping Precautions" and "General Information"

NOTE: When using Command 50 WP with other registered herbicides, always refer to rotational restrictions and precautions on the other product's label

1.0 LB. (0.50 LB. A.I.)	1.5 LBS. (0.75 LB. A.I.)	2.0 LBS. (1.0 LB. A.I.)
ANY TIME Soybeans, Pumpkins, Peas	ANY TIME Soybeans, Pumpkins	ANY TIME Soybeans, Pumpkins
9 MONTHS Corn -- Field -- Pop -- Seed -- Sweet	9 MONTHS Same as 1.0 lb., plus Peas	9 MONTHS Same as 1.5 lbs., except Seed Corn and Sweet Corn require a 12 month rotation.
Cotton Cucurbits Dry Beans Peanuts Peppers Potatoes Rice Snap Beans Sorghum Sugar Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)		
12 MONTHS Cabbage* Tomatoes (all)* Wheat	12 MONTHS Same as 1.0 lb.	12 MONTHS Same as 1.5 lbs., plus Seed Corn Sweet Corn (except fresh market)
16 MONTHS All crops *Direct seeded and transplanted	16 MONTHS All crops	16 MONTHS All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 50 WP treatment. Do not allow livestock to graze on treated plants or feed treated plants to livestock.

SOUTHERN AREA ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur.

Refer to sections headed "Rotational Cropping Precautions" and "General Information"

NOTE: When using Command 50 WP with other registered herbicides, always refer to rotational restrictions and precautions on the other product's label

1.0 LB. (0.50 LB. A.I.)	1.5 to 2.0 LBS. (0.75 to 1.0 LB. A.I.)	2.5 LBS. (1.25 LB. A.I.)
ANY TIME Soybeans Pumpkins Peas	ANY TIME Soybeans Pumpkins	ANY TIME Soybeans
9 MONTHS Corn -- Field -- Pop -- Seed -- Sweet	9 MONTHS Same as 1.0 lb. plus Peas	9 MONTHS Same as 1.5 to 2.0 lbs., plus Pumpkins
Cotton Cucurbits Dry Beans Peanuts Rice Snap Beans Sorghum Sugar Beets Sweet Potatoes Tobacco (Transplanted)		
12 MONTHS All crops	12 MONTHS All crops	12 MONTHS All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 50 WP treatment. Do not allow livestock to graze on treated plants or feed treated plants to livestock.

PUMPKINS

GENERAL INFORMATION

Command 50 WP herbicide may be utilized as a preemergent soil applied treatment for the control of annual grass and broadleaf weeds in pumpkin.

Command 50 WP must be soil incorporated in the following states unless an approved 24(C) is on file for counties within that state (check with your local dealer): Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, and Wisconsin.

NOTE: When utilizing Command 50 WP as a soil incorporated treatment, incorporate Command to a depth of 1 inch or less and place the seed below the chemical barrier when planting.

Command 50 WP herbicide may be tank mixed with other pumpkin herbicides to broaden the weed control spectrum compared to the products applied alone. When applying mixtures of Command 50 WP herbicide plus other pumpkin herbicides, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures. Water or liquid fertilizer may be used as a carrier for Command 50 WP when applied alone, or when tank mixed with the herbicides listed above unless use directions specifically state otherwise.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 50 WP herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

REPLANTING INSTRUCTIONS

If initial seeding of pumpkins fails to produce a stand, pumpkins may be replanted in fields treated with Command 50 WP alone. Do not retreat field with a second application of Command 50 WP. When tank mixing with a labeled product refer to the pumpkin replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Guidelines on this label.

NOTE: Seed should be planted below the herbicide barrier to minimize early bleaching of seedlings. Not all varieties have been tested for tolerance to a Command treatment. Pea type and variety, soil type and amount of organic matter, rate of application, depth of incorporation, planting depth, amount of moisture throughout the growing season and other factors can impact the degree of sensitivity that may be observed.

COMMAND 50 WP APPLIED ALONE RATES AND WEEDS CONTROLLED

Soil Texture	Broadleaf Rates Per Acre*
Coarse (light) Soils (loamy sand, sandy loam)	1 pound
Medium Soils (loam, silt, silt loam, sandy clay, sandy clay loam)	1 to 1½ pounds
Fine (heavy) Soils (silty clay, clay loam, silty clay loam, clay)	1½ to 2 pounds

*Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

NORTHERN AREA—See Map

1.0 LB. (0.5 LB. A.I.)	1.5 LBS. (0.75 LB. A.I.)	2.0 LBS. (1.0 LB. A.I.)
GRASSES Suppression of some annual grasses, e.g., Barnyardgrass Crabgrass —Large —Smooth Goosegrass	GRASSES Barnyardgrass Crabgrass —Large —Smooth Field Sandbur Foxtail —Giant —Green —Robert —Yellow Goosegrass Panicum —Common —Fall —Texas Seedling Johnsongrass*	GRASSES All grasses controlled at 1.5 pounds, plus: Bermudagrass* Cupgrass —Southwestern Woolly Seedling Johnsongrass Shattercane* Wild Proso Millet*

BROADLEAVES

Lambsquarters*
Shepardspurse*
Spurred Anoda
Velvetleaf

BROADLEAVES

All broadleaves controlled at 1 pound plus:
Common Ragweed*
Gallega
Henbit
Jimsonweed*
Kochia
Lambsquarters
Prickly Sida
Purslane
Shepardspurse
Smartweed
—Lady thumb
—Pennsylvania
Venice Mallow

BROADLEAVES

All broadleaves controlled at 1.5 pounds, plus:
Cocklebur*
Common Ragweed
Jimsonweed
Plantain, Black Seeded
Smartweed
—Lady thumb
—Pennsylvania
Tropic Croton

*Partially controlled

SOUTHERN AREA—See Map

1.0 LB. (0.5 LB. A.I.)	1.5 LBS. (0.75 LB. A.I.)	2.0 LBS. (1.0 LB. A.I.)
GRASSES Suppression of some annual grasses, e.g., Barnyardgrass Crabgrass —Large —Smooth Goosegrass	GRASSES Barnyardgrass Broadleaf Signalgrass Crabgrass —Large —Smooth Field Sandbur* Foxtail —Giant —Green Goosegrass Panicum —Common —Fall —Texas Seedling Johnsongrass	GRASSES All grasses controlled at 1.5 pounds plus: Bermudagrass* Cupgrass —Southwestern Woolly Field Sandbur Itchgrass** Red Rice* Red Sprangletop
BROADLEAVES Lambsquarters* Shepardspurse* Spurred Anoda Velvetleaf	BROADLEAVES All broadleaves controlled at 1 pound plus: Common Ragweed* Jimsonweed* Lambsquarters Pennsylvania Smartweed* Prickly Sida Purslane Redweed Shepardspurse Venice Mallow	BROADLEAVES All broadleaves controlled at 1.5 pounds plus: Cocklebur Common Ragweed Dayflower Florida Beggarweed Florida Fourly Jimsonweed Kochia* Redvine* Tropic Croton Wild Poinsettia*

*Partially controlled

**For control of Itchgrass, do not incorporate Command 50 WP

Rotate to crops as listed under Rotational Crop Guidelines, otherwise crop injury may occur.

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 50 WP treatment. Do not allow livestock to graze on treated pumpkin vines or feed treated vines or vine trash to livestock.

SUCCULENT PEAS

GENERAL INFORMATION

Command 50 WP herbicide may be utilized as preemergent soil applied treatment for the control of annual grass and broadleaf weeds in peas.

Command 50 WP must be soil incorporated in the following states unless an approved 24(C) is on file for counties within that state (check with your local dealer): Connecticut, Delaware, Illinois, Indiana, Iowa, Kentucky, Maine, Maryland, Massachusetts, Michigan, Minnesota, New Hampshire, New Jersey, New York, Ohio, Pennsylvania, Rhode Island, Vermont, Virginia, West Virginia, and Wisconsin.

NOTE: When utilizing Command 50 WP as a soil incorporated treatment, incorporate Command to a depth of 1 inch or less and place the seed below the chemical barrier when planting.

Command 50 WP herbicide may be tank mixed with other herbicides registered for use in peas to broaden the weed control spectrum compared to the products applied alone. When applying mixtures of Command 50 WP plus other succulent pea herbicides, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixes. Water or liquid fertilizer may be used as a carrier for Command 50 WP when applied alone, or when tank mixed with other succulent pea herbicides unless directions specifically state otherwise.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command® 50 WP herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications, read and strictly follow all precautions and application instructions on this label.

REPLANTING INSTRUCTIONS

If initial seeding of peas fails to produce a stand, peas may be replanted in fields treated with Command 50 WP alone (or with recommended tank mixtures). Do not retreat field with a second application of Command 50 WP. When tank mixing with a labeled product refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Guidelines on this label.

NOTE: Seed should be planted below the herbicide barrier to minimize early bleaching of seedlings. Not all varieties have been tested for tolerance to a Command treatment. Pumpkin type and variety, soil type and amount of organic matter, rate of application, depth of incorporation, planting depth, amount of moisture throughout the growing season and other factors can impact the degree of sensitivity that may be observed.

COMMAND 50 WP APPLIED ALONE RATES AND WEEDS CONTROLLED

Soil Texture	Broadcast Rates Per Acre*
Coarse (light) Soils: (loamy sand, sandy loam)	¾ pound
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	¾ to 1 pound
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	1 pound

*Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

¾ POUND
(0.375 LB. A.I.)

GRASSES

Barnyardgrass*
Crabgrass (Large, Smooth)*
Foxtail (Green, Giant, Robust)*
Goosegrass*
Panicum (Fall, Common)*

BROADLEAVES

Galinsoga spp.*
Lambsquarters*
Spurred Anoda*
Velvetleaf*

1 POUND
(0.5 LB. A.I.)

GRASSES

Barnyardgrass*
Crabgrass (Large, Smooth)
Field Sandbur*
Foxtail (Green, Giant, Robust)
Goosegrass
Panicum (Fall, Common)
Seedling Johnsongrass*

BROADLEAVES

Galinsoga spp.*
Lambsquarters*
Prickly Sida*
Shepherdspurge*
Spurred Anoda
Velvetleaf
Venice Mallow*

*Partially Controlled

Rotate to crops as listed under Rotational Crop Guidelines, otherwise crop injury may occur.

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 50 WP treatment. Do not allow livestock to graze on treated pea vines or feed treated vines or vine trash to livestock.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

COMMAND and FMC—Trademarks of FMC Corporation
©1987 FMC Corporation

All rights reserved
(1331-4/30/91-A)