

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

5705453474  
18

NOV 15 1993

Edward T. Cherry, Ph.D  
FMC Corporation  
Agricultural Chemical Group  
1735 Market Street  
Philadelphia, PA 19103

Dear Dr. Cherry:

Subject: Application for Amendment/Extension of Conditional  
Registration

- Ammo® 2.5 EC Insecticide  
EPA Registration No. 279-3027
  - Ammo® 2.5 Miscible Insecticide  
EPA Registration No. 279-3046
  - Ammo® 2.5 Oil Insecticide  
EPA Registration No. 279-3044
  - Ammo® WSB Insecticide  
EPA Registration No. 279-3084
  - Fury® 1.5 EC Insecticide  
EPA Registration No. 279-3125
  - Fury® 1.5 EW Insecticide  
EPA Registration No. 279-3126
  - Pounce® 3.2 EC Insecticide  
EPA Registration No. 279-3013
  - Pounce® 25 WP Insecticide  
EPA Registration No. 279-3051
  - Pounce® WSB Insecticide  
EPA Registration No. 279-3083
  - Capture® 2 EC Insecticide/Miticide  
EPA Registration No. 279-3069
  - Capture® EC-CAL Insecticide/Miticide  
EPA Registration No. 279-3114
- Your Letter Dated November 15, 1993

The Agency accepts your application for amended registration incorporating the terms and conditions stipulated in the subject letter for purposes of extending the registration of the subject product.

Therefore in accordance with FIFRA Section 3(c)(7)(B)(C) the conditional registration of the subject product is extended for a period not to exceed November 15 1996.

CONCURRENCES

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

A stamped copy of the final labeling submitted November 15, 1993 is enclosed. We agree that this labeling including supplemental labeling will be revised to incorporate the revisions specified in paragraph 2 of the subject letter before distribution or use. Please submit five copies of revised finished labeling when printing is completed.

Sincerely yours,

George T. LaRocca  
Product Manager 13  
Insecticide-Rodenticide Branch  
Registration Division (H7505C)

)  
  
)  
  
Enclosure

Code 1370

**RESTRICTED USE PESTICIDE**

Toxic to fish and aquatic organisms.  
For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

Net Contents

# Capture® 2 EC-CAL

## Insecticide/Miticide For Sale and Use Only in California

EPA Reg. No. 279-3114

EPA Est. 279-

Active Ingredient:	By Wt.
Bifenthrin: (2 methyl[1,1'-biphenyl]-3-yl) methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)- 2,2-dimethyl-cyclopropanecarboxylate* .....	25.1%
Inert Ingredients:** .....	74.9%
	100.0%

\*Cis isomers 97% minimum, trans isomers 3% maximum.  
\*\*Contains xylene range aromatic solvents.  
This product contains 2 pounds active ingredient per gallon.  
U.S. Patent No. 4,238,505

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

**AVISO**

This label must be in the possession of the user at the time of application.

See other panels for additional precautionary information.

**BEST AVAILABLE COPY**

# FMC.

FMC Corporation  
Agricultural Chemical Group  
Philadelphia PA 19103

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

NOV 15 1993

8/93

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

279-3114

### STATEMENT OF PRACTICAL TREATMENT

**If swallowed:** Get medical attention. Call a Poison Control Center or physician promptly for advice. Describe Precautionary Statements and Note to Physician on the label. Do not induce vomiting unless advised by a physician or qualified medical advisor. Do not give anything by mouth to an unconscious person.

**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**If on Skin:** Wash with plenty of soap and water. Get medical attention.

**If in Eyes:** Flush with plenty of water. Call a physician if irritation persists.

**Note to Physician:**  
Pesticide Hotline (800) 858-7378. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

For Emergency Assistance Call (800) 331-3148.

### PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

**Warning**  
May be fatal if swallowed. Harmful if inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash contaminated clothing before reuse.

Applicators must wear long sleeve shirt and trousers. Mixers and loaders must wear long sleeve shirt, trousers, chemical resistant gloves and goggles, or face shield.

### Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

### Physical/Chemical Hazards

Do not use or store near heat or open flame.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Resistance.** Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

## STORAGE AND DISPOSAL

### Pesticide Storage

Do not freeze. Do not store below 40° F. If crystals are observed, warm material to above 60° F by placing container in warm location. Shake or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: 1-(800)-331-3148.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### Container Disposal

Metal or Plastic Container: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers.

U-Turn® Container: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

## Special Equipment

The registration of Capture 2 EC-CAL in California requires that the product be used in closed systems that meet the criteria for closed systems as established by the California Department of Food and Agriculture. The criteria and a list of the closed systems meeting the criteria are available through the California Department of Food and Agriculture.

## Application Instructions

Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations; higher rates under heavy insect pressure and for mite control. Arid climates generally require higher rates.

CROP	PEST	LB A/A	FL OZ/A	REMARKS
Cotton	European Corn Borer Soybean (Banded) Thrips Tobacco Thrips	0.02-0.1	1.3-6.4	Capture 2 EC may be applied in water or refined vegetable oil (soybean/cottonseed). Application in Water: Apply in a minimum of 2 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray.
	Boil Weevil Boltworm Cabbage Looper Cotton Aphid Cotton Fleahopper Cotton Leafperforator Cutworms Fall Armyworm Lygus Spp Plant Bugs Saltmarsh Caterpillar Southern Garden Leafhopper Silk Bugs Tobacco Budworm Whitefly Yellow Striped Armyworm	0.04-0.1	2.6-6.4	ULV Application: Apply the recommended rate of Capture 2 EC in refined vegetable oil in a minimum of 1 quart of finished spray per acre with aircraft calibrated to give adequate coverage. To Control Boil Weevil: Apply Capture 2 EC at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels. To Control Mites and Aphids: Apply when pests first appear. Repeat as necessary to maintain control. Higher rates will be required once a damaging threshold is established.
	Beet Armyworm Carline Spider Mite Pink Bollworm Two-spotted Spider Mite	0.05-0.1	3.8-6.4	

Do not apply more than 0.3 pound active per acre per season.

Do not make more than 3 applications per season.

Do not apply within 14 days of harvest.

Do not graze livestock in treated areas or cut treated crops for feed.

### Spray Drift Precautions—Cotton

Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Ammo®, Asana® XL, Baythroid®, Capture®, Danitol®, Karate®, Mustang®, Pounce® and Scout® X-TRA.

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes; reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Do not cultivate within 10' of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

For Capture 2 EC-CAL, do not make more than 3 applications per growing season.

## Reentry Statements

Do not apply this product in such a manner as to directly, or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter until spray has dried.

### Cotton Scouts

Do not enter treated areas for 7 days after application.

Tractor Drivers, Irrigators, Beekeepers, and Nonroutine Cultural Activities

May enter after spray has dried provided they wear protective clothing. Protective clothing consists of long sleeved shirt, long legged trousers, shoes, socks, cloth gloves, hat or other suitable head covering.

## Rotational Crops

Rotational crops may be planted no sooner than 30 days after last application. Straw may not be used for food or feed.

## Tank-Mixture

Capture 2 EC insecticide/mixture may be applied in tank mixtures with other products approved for use on cotton. Observe all restrictions and precautions which appear on the labels of these products.

**BEST AVAILABLE COPY**

**Dealers Should Sell in Original Packages Only.**


**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.



Ambush, Cymbush and Karate—Trademarks of ICI Americas Inc.  
Asana—Trademark of E.I. duPont de Nemours and Co., Inc.  
Baythroid—Trademark of Bayer AG  
Scout—Trademark of Hoechst Roussel Agri-Vet Company  
U-Turn, Capture, Pounce, Ammo and ~~FMC~~—Trademarks of FMC Corporation  
© 1987 FMC Corporation All rights reserved. (1370-6/28/91-A)