

RESTRICTED USE PESTICIDE
 Toxic to fish and aquatic organisms.
 For retail sale to and use only by certified applicators, or persons under their direct supervision and only for the uses covered by the certified applicator's certification.



**Insecticide/Miticide
 For Sale and Use Only in
 California**

EPA Reg. No. 279-3114

EPA Est. 279-

Active Ingredient:	By Wt.
Bifenthrin: (2 methyl[1,1'-biphenyl]-3-yl) methyl 3-(2-chloro-3,3,3-trifluoro-1-propenyl)- 2,2-dimethyl-cyclopropanecarboxylate*	25.1%
Inert Ingredients:**	74.9%
	100.0%

*Cis isomers 97% minimum, trans isomers 3% maximum.
 **Contains xylene range aromatic solvents.
 This product contains 2 pounds active ingredient per gallon.
 U.S. Patent No. 4,238,505

**KEEP OUT OF REACH OF CHILDREN
 WARNING
 AVISO**

This label must be in the possession of the user at the time of application.
 Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See other panels for additional precautionary information.



FMC Corporation
 Agricultural Chemical Group
 Philadelphia PA 19103

CaptureCal_3_12-18-01

ACCEPTED
 DEC 18 2001
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 279-3114

FIRST AID

If Swallowed: Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If In Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

Note to Physician:

This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

For Emergency Assistance Call (800) 331-3148.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

May be fatal if swallowed. Harmful if inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothing.

Wash thoroughly with soap and water after handling and before eating, drinking, or using tobacco. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Handlers who may be exposed to the dilute through application or other tasks must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as Barrier Laminar or Nitrile Rubber or Neoprene Rubber or Viton, and shoes plus socks.

Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear: long-sleeved shirt and long pants, chemical-resistant gloves, such as Barrier Laminar or Nitrile Rubber or Neoprene Rubber or Viton, shoes plus socks, and protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations
Users should:
 Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Resistance. Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. For applications to sites with restricted entry intervals (REIs) of 7 days or more, notify workers of applications by warning them orally and by posting warning signs at entrances of treated areas.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) listed in the chart below:

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemical-resistant gloves, such as Barrier Lamine or Nitrile Rubber or Neoprene Rubber or Viton, and Shoes plus socks.

Crop	Activity	REI
Eggplant	All	9 days
Artichokes	All	5 Days
Beans (for processing only)	All	24 hours
Beans (for fresh market)	All	9 days
Peas (for processing only)	All	24 hours
Peas (for fresh market)	All	9 days
Caneberries (for mechanically harvested)	All	24 hours
Caneberries (all others)	All	4 days
Head, Stem Brassica	All	5 days
Sweet Corn (for processing only)	All	24 hours
Sweet Corn (for fresh market)	All	18 days
Field Corn (Grown for seed)	Detasseling	18 days
Field Corn (Grown for seed)	Roguing	18 days
Field Corn (Grown for seed)	Other Activities	24 hours
Cotton	Scouting	7 days
Cotton	Other Activities	24 hours
Other Crops	All	24 hours

STORAGE AND DISPOSAL

Pesticide Storage

Do not freeze. Do not store below 40° F. If crystals are observed, warm material to above 60° F by placing container in warm location. Shake or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: 1-(800)-331-3148.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Metal or Plastic Container: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Do not cut or weld metal containers.

U-Tum® Container: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

Special Equipment

The registration of Capture 2 EC-CAL in California requires that the product be used in closed systems that meet the criteria for closed systems as established by the California Department of Food and Agriculture. The criteria and a list of the closed systems meeting the criteria are available through the California Department of Food and Agriculture.

Rotational Crops

Rotational crops may be planted no sooner than 30 days after last application. Straw may not be used for food or feed.

Tank-Mixture

Capture 2 EC-CAL insecticide/miticide may be applied in tank-mixtures with other products approved for use on cotton. Observe all restrictions and precautions which appear on the labels of these products.

Application Instructions

Rate of application is variable according to pest pressure, timing of sprays, and field scouting. Use lower rates under light to moderate infestations; higher rates under heavy insect pressure and for mite control. And climates generally require higher rates.

Use of Capture 2EC-CAL is prohibited in all coastal counties.

Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

For LEPA irrigation a minimum of 0.75 inch of water per acre is recommended. Where non-emulsified oils are used as the diluent, 1 to 2 pints per acre is recommended.

Results from utilizing chemigation have been variable and depend upon the set up and calibration of equipment. Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. Contact your State Agricultural Extension Service specialists, equipment manufacturers or other experts for consultation on the suitability of the equipment set up to obtain effective control of the target insect pests.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise. Failure to cease application during a mechanical stoppage may result in undesirable residues to adjacent areas.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Capture[®] 2 EC insecticide/miticide should be applied continuously for the duration of the water application. Capture 2 EC should be diluted in sufficient volume to ensure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation generally is not required when a suitable diluent is used. A diluent test should be conducted to ensure that phase separation will not occur during dilution and application. Failure to achieve a uniform dilution throughout the time of application may result in undesirable residues or less than desirable control.

BRASSICAS:

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Head and Stem Brassica Vegetables including: Broccoli Chinese Broccoli (gai lan, white flowering broccoli) Brussels Sprouts Cauliflower Cavalo broccoli Kohlrabi Cabbage Chinese Cabbage (napa) Chinese Mustard Cabbage (gai choy)	Cutworms Corn Earworm Saltmarsh Caterpillar Flea Beetles Imported Cabbageworm Whitefly Armyworms Loopers Crickets Diamondback Moth Lygus Spp.	0.033 - 0.10 0.08 - 0.10	2.1 - 6.4 5.12 - 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.5 lb. active ingredient (1 quart) per acre per season. Do not make more than 5 applications after bloom. Do not make applications less than 7 days apart. Do not apply within 7 days of harvest.

CANE BERRIES

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Caneberries Including: Blackberries Bingleberries Dewberries Lowberries Marionberries Ollalieberries Youngberries Loganberries Raspberries	Leafrollers Orange Tortrix Root Weevils Spider Mites	0.05 - 0.10 0.10	3.2 - 6.4 6.4	Apply by air or ground equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons per acre by air and 50 gallons per acre by ground). One application may be made pre-bloom and a second application may be made post bloom. Do not apply within 3 days of harvest. Do not apply within 4 days of harvest for caneberries that are hand-harvested.

CANOLA, CRAMBE, RAPESEED

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Canola Crambe Rapeseed	Cutworms Aphids Cutworms Diamondback Moth Loopers Other Lepidopterous Larvae Flea Beetle Flea Hopper Grasshopper Plant Bug Stink Bugs Seedpod Weevil Thrips Whitefly Armyworms	0.033 - 0.04	2.1 - 2.6	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.08 lb active ingredient (5.12 ounces) per acre per season. Do not make applications less than 14 days apart. Do not apply within 35 days of harvest.

COTTON

CROP	PEST	DOSAGE		REMARKS
		LB A/A	FL OZ/A	
Cotton	European Corn Borer Soybean (Banded) Thrips Tobacco Thrips Boll Weevil Bollworm Cabbage Looper Cotton Aphid Cotton Fleahopper Cotton Leafperforator Cutworms Fall Armyworm Lygus Spp. Plant Bugs Saltmarsh Caterpillar Southern Garden Leafhopper Stink Bugs Tobacco Budworm Whitefly Yellow Striped Armyworm Beet Armyworm Carmine Spider Mite Pink Bollworm Twospotted Spider Mite	0.02-0.1 0.04-0.1 0.06-0.1	1.3-6.4 2.6-6.4 3.8-6.4	Capture 2 EC-CAL may be applied in water or refined vegetable oil (soybean/cottonseed). Application in Water: Apply in a minimum of 5 gallons per acre with ground equipment or 1 gallon per acre by aircraft. When applying by air, 1 quart of emulsified oil may be substituted for one quart of water in the finished spray. To Control Boll Weevil: Apply Capture 2 EC-CAL at an interval of 3 to 4 days until pest numbers are reduced to acceptable levels. To Control Mites and Aphids: Apply when pests first appear. Repeat as necessary to maintain control. Higher rates will be required once a damaging threshold is established.

Do not apply more than 0.3 pound active per acre per season.
Do not make more than 3 applications per season.
Do not apply within 14 days of harvest.
Do not graze livestock in treated areas or cut treated crops for feed.

Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush[®], Ammo[®], Asana[®] XL, Baythroid[®], Capture[®], Danitol[®], Karate[®], Mustang, Pounce[®] and Scout[®] X-TRA.

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes; reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Do not cultivate within 10' of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

For Capture 2 EC-CAL, do not make more than 3 applications per growing season.

Cotton Scouts - Restricted-entry interval 7 days.

FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED (AT PLANT USE)

PEST	DOSAGE		REMARKS
Corn Rootworm Larvae Mexican Northern Southern Western	.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row	Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the Capture 2EC needs per acre. Apply in a minimum of 3 gallons of finished spray per acre. Mix Capture 2EC with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of Capture 2EC, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.
Amy Cutworm Cutworm Species Grubs Seed Corn Beetle Seed Corn Maggot True Armyworm or Armyworm Species Wireworm	.0023 to .0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row	Applications of Capture 2EC alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of Capture 2EC and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application. Do not apply to soil where there is greater than 30% cover of crop residue remaining. Do not apply within 30 days of harvest Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment. Do not apply more than 0.1 pound active per acre per season as an at plant application.

Row Spacings (inches)	40	38	36	30
Capture 2EC (pounds ai per acre)	0.060	0.064	0.069	0.080
Capture 2EC (formulated ounces per acre)	3.9	4.1	4.4	5.12

FIELD CORN (GRAIN AND SILAGE), POPCORN, FIELD CORN GROWN FOR SEED

PEST	DOSAGE	
	LB A/A	FL OZ/A
Aphids	0.033-0.10	2.1-6.4
Amy Cutworm		
Beet Armyworm		
Cereal Leaf Beetle		
Chinch Bug		
Common Stalk Borer		
Corn Earworm		
Corn Rootworm Adults		
Cucumber Beetle Adult		
Cutworm Species		
European Corn Borer		
Fall Armyworm		
Flea Beetle		
Grasshoppers		
Greenbug		
Japanese Beetle Adult		
Leafhopper		
Sap Beetle		
Southern Armyworm		
Southern Corn Leaf Beetle		
Southwestern Corn Borer		
Stinkbugs		
Tamished Plant Bug		
Thrip		
True Armyworm or Armyworm Species		
Webworms		
Western Bean Cutworm		
Yellowstriped Armyworm		
Banks Grass Mite		
Carmine Mite	0.08-0.10	5.12-6.4
Twospotted Spider Mite		

RESTRICTIONS—Corn

Do not apply more than 0.2 pound active per acre per season.

Do not apply within 30 days of harvest.

Do not graze livestock in treated areas or cut treated crops for feed within 30 days of the last application.

Use of ultra low volume (ULV) application on corn is prohibited.

Do not make aerial or ground applications to corn if heavy rainfall is imminent.

REMARKS—Corn

General: Apply in a minimum of 2-5 gallons of finished spray per acre by aircraft or in a minimum of 10 gallons per acre with ground equipment. To improve control by aircraft, use 5 gallons of finished spray per acre particularly when initial populations are heavier than normal. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

To control ear-attacking pests: Apply Capture 2 EC-CAL just before silking and repeat as necessary to maintain control.

Southwestern Corn Borer, European Corn Borer: Make application for corn borer control with initial application at or shortly before egg hatch.

For control of other insect pests: Apply when pests first appear and repeat as necessary.

For Control of Mites:

Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.

For Twospotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before wide-spread mite dispersal throughout the canopy.

Higher rates will be necessary for heavier initial populations and corn under heat or drought stress. Field experience with dimethoate at 0.5 lb. active per acre in tank mixture has demonstrated good control under these conditions.

**SWEET CORN
SWEET CORN GROWN FOR SEED (AT PLANT USE)**

PEST	DOSAGE		REMARKS
	LB/A/A	FL OZ/A	
Corn Rootworm Larvae Mexican Northern Southern Western	.0046 pound active per 1,000 linear feet of row	0.30 fluid ounces per 1,000 linear feet of row	Apply as a 5 to 7 inch T-band treatment over an open seed furrow. Position the spray nozzle behind the planter shoe, in front of the press wheel centered over the row. Use the table below to determine the Capture 2EC needs per acre. Apply in a minimum of 3 gallons of finished spray per acre. Mix Capture 2EC with water or fertilizer in the following manner. Fill the spray tank approximately one-half full with water or liquid fertilizer, add the proper amount of Capture 2EC, then add the rest of the water or fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture. Applications of Capture 2EC alone or in recommended tank mixtures, in conjunction with in furrow pop-up fertilizers may be used. A jar compatibility test should be performed with appropriate ratio of Capture 2EC and fertilizer to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application. Do not apply to soil where there is greater than 30% cover of crop residue remaining. Do not apply within 30 days of harvest. Do not graze livestock in treated area or cut treated crops for feed within 30 days of treatment. Do not apply more than 0.1 pound active per acre per season as an at plant application.
Army Cutworm Cutworm Species Grubs Seed Corn Beetle Seed Corn Maggot True Armyworm or Armyworm Species Wireworm	.0023 to .0046 pound active per 1,000 linear feet of row	0.15 to 0.30 fluid ounces per 1,000 linear feet of row	

Row Spacings (inches)	40	38	36	30
Capture 2EC (pounds ai per acre)	0.060	0.064	0.069	0.080
Capture 2EC (formulated ounces per acre)	3.9	4.1	4.4	5.12

**SWEET CORN
SWEET CORN GROWN FOR SEED**

PEST	DOSAGE	
	LB A/A	FL OZ/A
Aphids	0.033 - 0.10	2.1 - 6.4
Army Cutworm		
Aster leafhopper		
Beet Armyworm		
Cereal Leaf Beetle		
Chinch Bug		
Common Stalk Borer		
Corn Earworm		
Corn Rootworm Adults		
Corn Silky		
Cucumber Beetle Adult		
Cutworm Species		
European Corn Borer		
Fall Armyworm		
Flea Beetle		
Grasshoppers		
Greenbug		
Japanese Beetle Adult		
Leafhopper		
Sap Beetle		
Southern Armyworm		
Southern Corn Leaf Beetle		
Southwestern Corn Borer		
Stinkbugs		
Tarnished Plant Bug		
Thrip		
True Armyworm or Armyworm Species		
Webworms		
Western Bean Cutworm		
Yellowstriped Armyworm		
Banks Grass Mite	0.08 - 0.10	5.12 - 6.4
Carmine Mite		
Pacific Spider Mite		
Twospotted Spider Mite		

RESTRICTIONS—Sweet Corn

Do not make more than 0.2 pounds active ingredient (12.8 ounces formulated) per acre per season.

Do not apply within one day of harvest.

Do not apply within 18 days of harvest for sweet corn that is hand-harvested.

Do not graze livestock in treated areas or cut treated crops for feed within 1 day of the last application.

Use of ultra low volume (ULV) application on corn is prohibited.

Do not make aerial or ground applications to corn if heavy rainfall is imminent.

REMARKS—Sweet Corn

General: Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control.

To control ear-attacking pests: Apply Capture 2EC-CAL when silking begins and repeat as necessary to maintain control.

Southwestern Corn Borer, European Corn Borer: Make 2 applications for corn borer control with the initial application at or shortly before egg hatch.

For control of other insect pests: Apply when pests first appear and repeat as necessary.

For Control of Mites:

Apply for Banks Grass Mite control when colonies first form prior to leaf damage or discoloration and before dispersal above the bottom third of the plant.

For Twospotted Spider Mite and Carmine Mite control, apply when colonies first form prior to leaf damage or discoloration and before widespread mite dispersal throughout the canopy.

Higher rates will be necessary for heavier initial populations and corn under heat or drought stress.

ARTICHOKE

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Cnbrate Weevil Artichoke Plume Moth	0.10	6.4	Apply when pest population reaches damaging threshold and repeat as necessary to maintain control, but not more often than 15 day intervals. Application by ground: Apply a full cover spray in a minimum of 75 gallons of finished spray per acre. Application by air: Apply specified dosage in a minimum of 10 gallons per acre. A 5-day preharvest interval must be observed.

CUCURBITS:

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Chayote (fruit) Chinese wax gourd (Chinese preserving melon) Citron melon Cucumber Gherkin Gourd, edible (includes hyotan, cuozza, (Luffa spp.)) (includes hachima, Chinese okra) (Momordica spp.) (includes balsam apple, balsam pear, bitter melon, Chinese cucumber) Muskmelon (hybrids and/or cultivars of Cucumis melo) (includes true cantaloupe, casaba, cranshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon) Pumpkin (Cucurbita spp.) Squash, summer (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini) Squash, winter (includes butternut squash, calabaza, Hubbard squash (C. mixta; C. pepo)) (includes acorn squash, spaghetti squash) Watermelon (includes hybrids and/or varieties of Citullus spp.)	Aphids Cutworms Cabbage Looper Leafhoppers Cucumber Beetles Squash Bugs Armyworms Melonworm Pickleworm Plant bug Stink Bugs Rindworm Squash Vine Borer Corn Earworm Tobacco Budworm Grasshopper	0.04 - 0.10	2.6 - 6.4	Apply in a minimum of 5 gallons of finished spray per acre by air or in a minimum of 20 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.3 lb active ingredient (19.2 ounces formulated) per acre per season. Do not make more than two applications after bloom. Do not make applications less than 7 days apart. Do not apply within 3 days of harvest.
	Whitefly Twospotted Spider Mite Carmine Mite Banks Grass Mite Lygus spp.	0.08 - 0.10	5.12 - 6.4	

PEPPERS, BELL AND NON-BELL

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Aphids Armyworms Corn earworm Cucumber Beetles Cutworms European Corn Borer Flea Beetles Leafhoppers Leafminers Looper Pepper Weevil Stink bug spp Tobacco budworm Thrips Whitefly	0.033 to 0.1	2.1 to 6.4	Apply using sufficient water to obtain uniform coverage. Apply as needed. Apply with ground equipment using a minimum of 10 gallons of finished spray per acre or a minimum of 2 gallons per acre by aircraft. Do not make applications less than 7 days apart. Do not apply more than 0.20 pound active ingredient per acre per season. Do not apply within 7 days of harvest.
Lygus Spp. Broad Mite Carmine Mite Two Spotted Spider Mite	0.08 to 0.10	5.12 to 6.4	

SUCCULENT PEAS AND BEANS

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Pea (Pisum spp.): Dwarf pea, Edible-pod pea, English pea Garden pea, Green pea, Snow pea, Sugar snap pea Pigeon pea Bean (Phaseolus spp.): Broadbean (succulent) Lima bean (green) Runner bean, Snap bean, Wax bean Bean (Vigna spp.): Asparagus bean, Blackeyed pea Chinese longbean, Cowpea Moth bean, Southern pea Yardlong bean Jackbean Soybean (immature seed) Sword bean	Flea Beetle Grasshoppers Aster Leafhopper Leafhoppers Aphids Beet Armyworm Fall Armyworm Southern Armyworm Yellowstriped Armyworm Bean Leaf Beetle Cucumber Beetles Japanese beetle Adult Sap Beetle Plant Bug Stink Bugs Tarnished Plant Bug Alfalfa Caterpillar Cloverworm European Corn Borer Cutworms Western Bean Cutworm Corn Earworm Looper Corn Rootworm Adult Thrips Webworms Pea Weevil Pea Leaf Weevil Whitefly	0.025 - 0.10	1.6 - 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not apply more than 0.2 lb. active ingredient (12.8 ounces formulated) per acre per season. Do not apply within 3 days of harvest. Do not apply within 9 days of harvest for peas/beans that are hand-harvested.
	Banks Grass Mite Twospotted Spider Mite Carmine Mite Lygus Spp	0.08 - 0.10	5.12 - 6.4	

LETTUCE, HEAD

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Aphids Armyworms Cabbage Maggot Corn earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Grasshopper Imported Cabbageworm Leafhoppers Looper Salt Marsh Caterpillar Stink bug Spp. Thrips Tobacco Budworm Whitefly	0.033 to 0.10	2.1 to 6.4	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not make applications less than 7 days apart. A maximum of 0.5 lb. active ingredient may be applied per acre per season. Do not apply within 7 days of harvest.
Lygus Spp. Carmine Mite Two Spotted Spider Mite	0.08 to 0.10	5.12 to 6.4	

EGGPLANT

CROP	PEST	DOSAGE		REMARKS
		LB/A/A	FLOZ/A	
Eggplant	Colorado Potato Beetle Corn Earworm European Corn Borer Flea Beetle Cucumber Beetle Cabbage Looper Tomato Pinworm Tomato Hornworm Vegetable Leafminer Whitefly Armyworms Plant Bug Stink Bug Thrips	0.033 - 0.10	2.1 - 6.4	Apply in a minimum of 2 gallons of finished spray per acre by air or in a minimum of 10 gallons per acre with ground equipment. When applying by air, 1-2 quarts of emulsified oil may be substituted for 1-2 quarts of water in the finished spray. Thorough coverage is essential to achieve control. Do not make applications less than 7 days apart. Do not apply more than 0.2 lb active ingredient (12.8 ounces formulated) per acre per season. Do not apply within 9 days of harvest.
	Banks Grass Mite Twospotted Spider Mite Carmine Mite Pacific Spider Mite Lygus Spp.	0.08 - 0.10	5.12 - 6.4	

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Ambush, Cymbush and Karate—Trademarks of ICI Americas Inc.
Asana—Trademark of E.I. duPont de Nemours and Co., Inc.
Baythroid—Trademark of Bayer AG
Scout—Trademark of Hoechst Roussel Agri-Vet Company
U-Turn, Capture, Ammo, Mustang, and ~~FMC~~—Trademarks of FMC Corporation
© 2001 FMC Corporation All rights reserved. (777-9/3/99-A)

HOPS

PEST	DOSAGE		REMARKS
	LB/A/A	FLOZ/A	
Aphids Armyworms Cutworms Leafrollers Loopers	0.06 to 0.1	3.8 to 6.4	Do not exceed 0.1 lb ai per acre per application. Do not exceed 0.3 lb ai per acre per season.
Root Weevils	0.05 to 0.1	3.2 to 6.4	A spray interval of 21 days between applications must be maintained.
Twospotted spider mite	0.1	6.4	A 14 day pre-harvest interval must be observed. Application by ground: For best results, full coverage is essential. Early season recommend 100-150 gallons of spray per acre. Late season recommend 200-250 gallons of spray per acre. For Root Weevil control, make a directed spray to the base of the plant. Spray up the vine 3 feet and the soil surface 1.5 to 2 feet on either side of the plant. Application by air for late season control of twospotted spider mites: Apply no less than 16 oz. (0.1 lb ai) per application in a minimum of 10 gallons per acre.