

OCT - 3 2001

Ms. Nancy J. Hilton
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

Dear Ms. Hilton:

Subject: Amendment - Additional Uses-Sweet Corn, Rice, Leafy Vegetables (Except Brassica), Brassica Leafy Vegetables, Sugar Beets, Sugarcane, Corn (Field, Seed, Pop), Green Onions, and Alfalfa

Fury Technical Insecticide

EPA Registration No. 279-3124

Fury 1.5 EC Insecticide

EPA Registration No. 279-3125

Fury 1.5 EW Insecticide

EPA Registration No. 279-3126

Your submissions dated October 1, 2001

The Agency accepts your application for amended registration to add the subject crops to the subject labeling since you have agreed in writing to the following terms for conditional registration in accordance with FIFRA section 3(c)(7)(b)(c):

1. You will submit and/or cite all data required for registration/reregistration of your product under FIFRA sec.3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit responses required for reregistration of your product under FIFRA section 4.

2. The subject amendment is conditioned under the same terms and condition for data generation as stipulated in our November 15, 1993 letter for use of this product on cotton.

3. FMC Corporation agreed that the current synthetic pyrethroid mitigation measures are interim in nature and may be reconsidered or modified after review and evaluation of the Spray Drift Task Force.

4. You understand that this registration will expire on November 15, 2003. You further understand that it is EPA's stated intent to, by November 15, 2003, complete its review of all relevant data and other information that are available to the Agency, and to make FIFRA section 3(c)(5) or other appropriate regulatory decisions for cotton-use synthetic pyrethroids and other crops conditionally registered based on the Agency's review of such data/information and considering statutory and regulatory criteria for such decisions.

5. You will submit product information (pounds or gallons produced) for this product for the fiscal year in which the added uses are conditionally registered, in accordance with FIFRA section 29. The fiscal year begins October 1 and ends September 30. The production information will be submitted to the Agency no later than November 15 following the end of the preceding fiscal year.

The information should be submitted to:

U.S. Environment Protection Agency
Office of Pesticide Program (7504C)
Document Processing Desk
Ariel Rios Building
1200 Pennsylvania Avenue, NW
Washington, DC 20460

6. You have agreed to submit the data listed below, conducted in accordance with the 40CFR Part 158 Test Guidelines unless the Agency concludes that the additional information submitted (faxed) on September 25, 2001 negates the need to redo these studies:

| Title of Study | Guideline Reference No. | Date Due |
|---------------------------------|----------------------------|-------------------|
| Aerobic Aquatic Metabolism | 162-4 | September 1, 2002 |
| Anaerobic Aquatic Metabolism | 162-3 | September 1, 2002 |

*Negotiable pending further discussions between you and the Agency.

You agreed to submit two copies of the revised final printed label for the record.

You should note if you fail to satisfy any of the conditions imposed on this registration, e.g., you fail to submit the required data by the specified deadlines or the data submitted were not generated in accordance with the applicable test guidelines, EPA may issue a notice to cancel this use under FIFRA 6(e).

You should also note that, regardless of whether you satisfy all applicable conditions, this conditional registration will expire automatically on November 15, 2003. Sale and distribution of the subject product bearing labeling for subject uses after November 15, 2003 will be illegal.

Finally, once the required data have been submitted and evaluated, EPA will entertain an application to amend the registration of the subject product to allow the unconditional use on cotton and other crops without any special restrictions on the duration of the registration.

A copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (13)
Insecticide Branch
Registration Division (7505C)

Enclosure

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.



FURY®

Insecticide

EPA Reg. No. 279-3125

EPA Est. 279-FL-1

Active Ingredient:

* S-Cyano (3-phenoxyphenyl)methyl
(±) cis/trans 3-(2,2-dichloroethenyl)-
2,2 dimethylcyclopropane carboxylate**

By Wt.

| | |
|------------------------|--------|
| Inert Ingredients..... | 18.1% |
| | 81.9% |
| | 100.0% |

*Contains 1.5 pounds active ingredient per gallon.

**Cis/trans ratio: Max. 55% (±) cis and min. 45% (±) trans
U.S. Patent No. Pending

KEEP OUT OF REACH OF CHILDREN

WARNING

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

FMC

FMC Corporation
Agricultural Products Group
Philadelphia PA 19103

ACCEPTED
with COMMENTS
in EPA Letter Dated

Fury 1.5EC_3_9-28-01

OCT - 3 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

279-3125

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

Note to Physician: Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages by aspiration of the solvent.

For Emergency Assistance Call (800) 331-3148.

See other panels for additional precautionary information.

PRECAUTIONARY STATEMENTS**Hazards to Humans (and Domestic Animals)****Warning**

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. The active ingredient may cause sensitization reaction in some individuals.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Handlers who may be exposed to the dilute through application or other tasks must wear: Long-sleeved shirt and long pants, Chemical-resistant gloves, such as Barrier Laminate or Viton, and Shoes plus socks.

Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear: Long-sleeved shirt and long pants, Chemical-resistant gloves, such as Barrier Laminate or Viton, Shoes plus socks, and Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations**Users should:**

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Resistance. Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemical-resistant gloves, such as Barrier Laminate or Viton, and Shoes plus socks.

STORAGE AND DISPOSAL

Pesticide Storage

Store in a cool, dry, well-ventilated place. Do not store below -6.5 C (20 F). If solids are observed warm to above 4.4 C (40 F) and roll or shake containers to redissolve. Do not use near heat, open flame or hot surfaces. Store in original containers only. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrates or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. Keep out of reach of children and animals.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148.

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Sealed Containers: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Fury® insecticide should be applied continuously for the duration of the water application. Fury should be diluted in sufficient volume to insure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation is not required when a suitable diluent is used.

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

For cutworm control, Fury® insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control.

Rotational Crops

With the exception of the crops listed below, rotational crops should not be planted within 30 days of last application.

Tank-Mixture

Fury® insecticide may be applied in tank mixtures with other products approved for use on Alfalfa, Brassicas, Bulb Vegetables, Corn, Cotton, Head Lettuce, Pecans, Rice, and Sugarcane. Observe all restrictions and precautions which appear on the labels of these products. Test for compatibility of products before mixing.

Spray Drift Precautions

Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Ammo®, Asana® XL, Baythroid®, Capture®, Danitol®, Fury®, Karate®, Mustang®, and Scout® X-TRA.

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMANENT STREAMS, MARSHES OR NATURAL PONDS; ESTUARIES AND COMMERCIAL FISH FARM PONDS.

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes; reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made. For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Do not cultivate within 10' of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Maximum Usage When Applying Both Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either product when used alone, and do not apply more than the combined maximum seasonal total for both products as outlined in the table below.

| Crop | Maximum Seasonal Total for Either Product Used Alone (pounds active ingredient/acre) | | Maximum Seasonal Total When Applying Both Products to the Same Crop (pounds active ingredient/acre) |
|------------------------|--|---------------------|---|
| | Zeta-cypermethrin (Fury or Mustang) | Cypermethrin (Ammo) | Zeta-cypermethrin (Fury or Mustang) plus Cypermethrin (Ammo) |
| Cotton | 0.3 | 0.6 | 0.8 |
| Head and Stem Brassica | 0.3 | 0.6 | 0.6 |
| Leafy Brassica Greens | 0.3 | 0.4 | 0.4 |
| Head Lettuce | 0.3 | 0.6 | 0.6 |
| Bulb Vegetables | 0.25 | 0.5 | 0.5 |
| Pecans | 0.3 | 0.6 | 0.6 |

Alfalfa: Alfalfa grown for seed (Includes lucerne, sainfoin, holy clover, esparcet, birdsfoot trefoil and varieties and/or hybrids of these)

| Insects Controlled | Rate of Application | Method of Application |
|---|---|---|
| Alfalfa Caterpillar Alfalfa Looper Alfalfa Weevil Cutworms Egyptian Alfalfa Weevil (larvae & adult) Flea Beetles Green Cloverworm Meadow Spittlebug Potato Leafhopper Three Corned Alfalfa Hopper Velvetbean Caterpillar Webworms Blue Alfalfa Aphid* Green Peach Aphid* Pea Aphid* Spotted Alfalfa Aphid* | 2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre | Apply as insects appear in sufficient volume of water to ensure thorough coverage of foliage. Use higher recommended dosage for increased pest pressure or for increased residual pest control. Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment. ULV oil spray application is prohibited. Higher volumes of finished spray may improve insect control under high temperatures, when foliage is dense and/or when insect pressure is high. |
| Armyworms Grasshoppers Plant Bugs (including Lygus spp. & Stink Bugs) | 3.0 to 4.3 ounces (0.035 to 0.05 pound active) per acre | Follow appropriate spray drift precautions on this label. *Aphid control may be variable depending on species present and host-plant relationships. |

Do not make applications less than 7 days apart.

A maximum of 0.05 pounds active ingredient/acre may be applied per cutting and a maximum of 0.15 pounds active ingredient per acre per season.

Applications may be made up to 3 days of cutting or grazing or up to 7 days of harvesting seed.

Head and Stem Brassica Vegetables (1 day ph) including: Broccoli; Chinese Broccoli (gai lan, white flowering broccoli); Brussels Sprouts; Cauliflower; Cavalo broccoli; Kohlrabi; Cabbage; Chinese Cabbage (napa); Leafy Brassica Greens including: Broccoli Raab (rapini); Chinese cabbage (bok choy); Chinese Mustard Cabbage (gai choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens

| Insects Controlled | Rate of Application | Method of Application |
|---|---|--|
| Corn Earworm Cucumber Beetles Cutworm Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Saltmarsh Caterpillar Southern Cabbageworm Tobacco Budworm | 2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. |
| Alfalfa Looper Armyworms Cabbage Looper Cabbage Webworm Crickets Grasshoppers Ground Beetles Leafminers (adults) Lygus Bugs Onion Thrips Stinkbugs Wireworm (adults) Aphids* Whiteflies* | 3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre | Lower rates of Fury® should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations. In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. Follow appropriate spray drift precautions on this label. |

Do not make applications less than 7 days apart.

A maximum of 0.3 pounds active ingredient may be applied per acre per season.

Bulb Vegetables (Allium spp.) (7 day phi) including: Garlic; Garlic, Great-Headed (elephant); Green Eschalots; Japanese Bunching Onions; Leeks; Onion, Dry Bulb and Green; Onion, Welch; Shallots, Dry Bulb and Green; Spring Onion or Scallions

| Insects Controlled | Rate of Application | Method of Application |
|---|--|--|
| Onion Thrips | 3.2 to 4.3 ounces (0.0375 to 0.05 pound active) per acre | Apply in a minimum of 20 gallons per acre with ground equipment or in a minimum of 3 gallons per acre by aircraft. Begin applications when pests appear and repeat as necessary to maintain control. |
| Armyworms Cutworms Leafminers Onion Maggot Adults Stink Bugs Aphids* | 2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre | To control Onion Thrips: Use higher rates as population increases and avoid rescue situations. Use of a crop oil concentrate at 16 fluid ounces per acre is recommended. Follow appropriate spray drift precautions on this label. |
| *aids in control | | |
| Do not make applications less than 7 days apart. | | |
| Do not apply more than 0.25 pound active ingredient per acre per season. | | |
| Do not graze livestock in treated areas or cut treated crops for feed. | | |

Corn, Sweet (3 day phi)

| Insects Controlled | Rate of Application | Method of Application |
|--|---|--|
| Chinch Bug Corn Rootworm (Adult) Corn Silky Cutworms Flea Beetle Leafhoppers Japanese Beetle (Adult) Sap Beetle (adults) Tarnished Plant Bug | 2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre | Apply with ground or air equipment using sufficient water and application methods to insure thorough coverage of foliage. Apply in water using a minimum of 20 gallons of finished spray per acre with ground equipment and a minimum of 2 gallons per acre by air. |
| Armyworms Corn Borers Corn Earworm Grasshoppers Aphids* | 3.0 to 4.3 ounces (0.035 to 0.05 pound active) per acre | Follow appropriate spray drift precautions on this label. |
| *aids in control | | |
| Apply at minimum 3 to 5 day intervals or as needed for control. | | |
| A maximum of 0.3 pounds active ingredient per acre per season may be applied. | | |

Corn (Field), Field Corn Grown for Seed, Popcorn (At Plant Use)

| Insects Controlled | Rate of Application | Method of Application |
|--|--|--|
| Cutworms | 0.16 fluid ounces per 1,000 linear feet of row (0.002 pound active) per 1,000 linear feet of row | Apply as an in-furrow, band or T-band treatment using a minimum 4" band. Use table below to determine the Fury needs for each acre. Do not apply more than 0.20 pound active ingredient per acre per season including at-plant plus foliar applications of Fury. Do not apply within 30 days of harvest for grain and fodder (stover) and 60 days for forage (silage). |
| Row Spacings (inches) | 40 | 30 |
| Fury 1.5 EC (pounds ai per acre) | 0.024 | 0.036 |
| Fury 1.5 EC (formulated ounces per acre) | 2.05 | 3.08 |

Corn (Field), Field Corn Grown for Seed, Popcorn

| Insects Controlled | Rate of Application | Method of Application |
|---|--|---|
| Cutworms | 1.4 to 2.9 ounces (0.016 to 0.034 pound active) per acre | Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results. |
| Bean Leaf Beetle Cereal Leaf Beetle Corn Rootworm Beetle European Corn Borer Flea Beetle Grasshopper Hop Vine Borer Japanese Beetle (adult) Sap Beetle (adult) Southern Corn Leaf Beetle Southwestern Corn Borer Stalk Borer Stink Bug Spp. Tobacco Budworm Webworms Aphids* | 2.9 to 4.3 ounces (0.034 to 0.05 pound active) per acre | Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage (minimum of 2 gallons per acre by air and 10 gallons per acre by ground). Do not apply more than 0.20 pound active ingredient per acre per season including At-Planting plus foliar applications of Fury insecticide. Do not apply within 30 days of harvest for grain and fodder (stover) and 60 days for forage (silage). For chinch bug control, scout corn fields and make applications when bugs migrate from small grains or wild grasses to small corn. Direct spray to the base of plant. Repeat applications at 3 to 5 day intervals if needed. Fury may only suppress heavy infestations and/or subsequent migrations. |
| Armyworms (including Fall Armyworms) Chinch Bug | 3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre | Follow appropriate spray drift precautions on this label. |
| Corn Earworm ¹ Western Bean Cutworm ¹ Green Cloverworm Meadow Spittlebug | 1.9 to 4.3 ounces (0.022 to 0.05 pound active) per acre | *Control may be variable depending on species present and host-plant relationships. ¹ For control before the larva bores into the plant stalk or ear. |

Cotton (14 day phi)

| Insects Controlled | Rate of Application | Method of Application |
|--|---|---|
| Preemergent Use: Cutworms | 1.4 to 2.0 ounces (0.016 to 0.024 pound active) per acre | Use Fury* in the time period from 14 days prior to planting up to emergence of the crop. Apply as a broadcast spray by ground or air, banded (including T-band) or in-furrow spray using sufficient spray volume to achieve adequate coverage. Reduced volumes of water may be used with specialized equipment. Use a minimum of 1 gallon of water per acre by air. Use the higher rates of Fury* when incorporating into the soil. |
| Foliar Use: Cutworms Tobacco Thrips Soybean (banded) Thrips | 1.4 to 2.0 ounces (0.016 to 0.024 pound active) per acre | Fury* may be applied in water or refined vegetable oil. When water is used, apply a minimum of one gallon of finished spray per acre by air or five gallons of finished spray with ground equipment. When applying in water by air, one quart of emulsified oil may be substituted for one quart of water in the finished spray. When using oil, use a minimum of one quart per acre in the finished spray. Control of lepidopteran eggs may be achieved with proper timing of applications. |
| Foliar Use: Boll Weevil Cabbage Looper Cotton Bollworm Cotton Fleahopper Cotton Leaf Perforator European Corn Borer Fall Armyworm Pink Bollworm Saltmarsh Caterpillar Stink Bugs Tarnished Plant Bug Other Plant Bugs Tobacco Budworm Yellow Striped Armyworm | 2.8 to 3.8 ounces (0.033 to 0.045 pound active) per acre | Fury* may be injected into overhead sprinkler irrigation water provided 1) an anti-backflow check valve is present between the injection point and the water source, 2) a check valve is present in the line to prevent irrigation water from entering the chemical supply tank and 3) the irrigation injection system has interlocking on-off switches. |
| Foliar use: Lygus Bugs Beet Armyworm* Cotton Aphid** Whiteflies** | 3.0 to 4.3 ounces (0.035 to 0.050 pound active) per acre | For boll weevil control, apply Fury* at a 3 to 4 day interval until pest numbers are reduced to acceptable levels. Follow appropriate spray drift precautions on this label. |
| Foliar use: Grasshoppers | 3.2 to 4.3 ounces (0.0375 to 0.050 pound active) per acre | For control of grasshoppers, applications should be made based on careful field scouting. Treatment decisions should be made based on evidence of feeding damage and presence of grasshoppers in cotton. Loss of cotyledon leaves in seedling cotton should be considered more important than leaf loss in older cotton. Applications should be made on a broadcast basis since grasshopper are highly mobile. Adjust rates based on populations of grasshopper found in fields. Applications should be made on a three to five day schedule until grasshopper populations are under control or until foliage loss subsides. Increase application rates as grasshopper size and population density increases. |
| A maximum of 0.3 pound active ingredient may be applied per acre per season. Do not graze or feed cotton for forage. | | |
| * For control of beet armyworms only in the high plains of Texas, Arizona, and California. | | |
| ** Aids in control | | |

Lettuce, Head (5 day phi)

| Insects Controlled | Rate of Application | Method of Application |
|---|---|---|
| Corn Earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Saltmarsh Caterpillar Tobacco Budworm | 2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre | Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of Fury* should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations. |
| Armyworms Crickets Loopers Lygus Bugs Onion Thrips Stink Bugs | 3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre | In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. Follow appropriate spray drift precautions on this label. |
| A maximum of 0.3 pound active ingredient may be applied per acre per season. | | |

Pecans (21 day phi)

| Insects Controlled | Rate of Application | Method of Application |
|--|--|---|
| Black Pecan Aphid Hickory Shuckworm Pecan Nut Casebearer Pecan Weevil Yellow Pecan Aphid | 2.8 to 4.3 ounces (0.033 to 0.050 pound active) per acre | Applications at the lower rate should be made when pest populations are low. Rates should be increased as the pest pressure increases. Apply by ground equipment to the point of drip. Use 100 gallons of dilute spray per acre for smaller trees. For larger trees which require higher gallonage to achieve adequate coverage, apply in 200 to 300 gallons of water. In order to calculate the correct number of gallons of water needed to spray one acre of your trees to the point of drip, you may need to conduct a test. If you do not know how to conduct such a test with your equipment, you should request assistance from your equipment dealer. |
| Up to 0.3 pound active ingredient per acre season may be applied prior to shuck split. | | |
| Do not graze livestock in treated orchards or cut treated cover crops for feed. | | |

Rice (14 day phi)

| Insects Controlled | Rate of Application | Method of Application |
|--|---|---|
| Fall Armyworm Grasshopper Spp. Green Bug Leafhopper Spp. Rice Water Weevil (adult) True Armyworm Yellow-striped Armyworm Oat Birdcherry Aphid* | 3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre | Apply as needed based on pest thresholds determined by scouting practices. Refer to Extension Scouting guidelines for scouting techniques, pest thresholds and treatment timing and treatment intervals. Determine the need for repeat applications, usually at intervals of 7 days, by scouting. |
| Chinch Bug Rice Stink Bug | 2.8 to 4.3 ounces (0.033 to 0.05 pound active) per acre | Fury® can be safely applied in conjunction with approved rice herbicides. Apply by air or ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 5 gallons of water per acre. For increased control, crop oil concentrate at 16 fluid ounces per acre may be used. For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations. For control of rice water weevil in water seeded rice, make the first application after flooding, when scouting indicates the presence of adults and/or feeding scars. Application should usually begin when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations. Green bug is known to have many biotypes. Fury® may only provide suppression. If satisfactory control is not achieved with the first application of Fury, a resistant biotype may be present. Use alternate chemistry for control. Follow appropriate spray drift precautions on this label. |
| *aids in control Do not make applications less than 7 days apart. Do not release floodwater within 7 days of an application. A maximum of 0.20 pound active ingredient (1.1 pints) may be applied per acre per season. Do not use treated rice field for the aquaculture of edible fish and crustacea. Do not apply as an ultra-low volume (ULV) spray. | | |

Sugarcane (21 day phi)

| Insects Controlled | Rate of Application | Method of Application |
|---|--|--|
| Sugarcane (Foliar use) | 3.2 to 4.3 ounces (0.0375 to 0.05 pound active) per acre | Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results. Apply by air or ground equipment using sufficient water to obtain full coverage of foliage (minimum of 2 gallons per acre by air and 10 gallons per acre by ground). Follow appropriate spray drift precautions on this label. |
| Do not make applications less than 21 days apart. Do not apply more than 0.20 pound active ingredient per acre per season. | | |

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

FMC, Ammo, Capture, Fury, Mustang—trademarks of FMC Corporation
 Asana—trademark of E.I. duPont de Nemours & Company
 Ambush, Karate—trademarks of Zeneca, LTD.
 Baythroid—Trademark of Bayer Aktiengesellschaft
 Danitol—trademark of Sumitomo Chemical Company, LTD
 Scout—trademark of Hoechst Schering Agrevo S.A.
 © 1999 FMC Corporation All Rights Reserved

(1425-3/17/97-A)