

279-3125

3/10/2010

1027



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Ms. Nancy Hilton
FMC Corporation
1735 Market Street
Philadelphia, PA 19103

MAR 10 2010

Dear Ms. Hilton:

Subject: Amendment- at plant applications for legume vegetables (dried peas and beans) and sunflower (and associated crops
Fury Insecticide
EPA Registration Number 279-3125
Your submission dated June 17, 2010

The amendment referred to above, submitted in connection with registration under the FIFRA section 3(c) (7) (a), is acceptable provided you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA sec. 3(c) (5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.

2. Make the label change listed below before you release the product for shipment bearing the amended labeling:

- a. You must clarify the restriction statement on legume vegetables and sunflowers. Change "Maximum amount per season: Do not apply more than -lb ai/A per season" with "Maximum amount per season: Do not apply more than -lb ai/A per season including at-plant foliar application." Without such a statement it could be misunderstood that the at-plant application could be made in addition to the seasonal rate allowed for foliar application.
- b. Update your label for storage and disposal in accordance with PR Notice 2007-4.

3. You agree that the subject registrations are conditional under the same terms and conditions for data generation as stipulated in the Agency's November 15, 1993 and November 15, 2004 letters for use of these products on cotton.

4. You agree that the current synthetic pyrethroid mitigation measures are interim in nature and may be reconsidered or modified after review and evaluation of the Spray drift Task Force data.

2027

5. FMC understands that it is the US EPA intent to complete the reviews of all relevant data and other information that are available to the agency, and to make FIFRA 3(c)(5) or other appropriate regulatory decisions for cotton- use synthetic pyrethroids and other crops conditionally registered based on the Agency's review of such data/information and considering statutory and regulatory criteria for such decisions.

A stamped copy of the label is enclosed for your records. Please submit three (3) copies of the amended labeling bearing the above revisions before you release the product for shipment.

If you have any questions you may contact Linda A. DeLuise at 703.305.5428.

Sincerely yours,



f Richard Gebken
Product Manager (10)
Insecticide Branch
Registration Division (7505P)


Enclosure

30f27

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.



RECEIVED
MAR 10 2010
FBI LABORATORY

MAR 10 2010

Under the Federal Insecticide, Fungicide, and Rodenticide Act
It is prohibited for the manufacture
registered under EPA Reg. No.
279-3125

FURY® Insecticide

EPA Reg. No. 279-3125

EPA Est. 279-FL-1

Active Ingredient:

By Wt.

* S-Cyano (3-phenoxyphenyl)methyl (±) cis/trans 3-(2,2-dichloroethenyl)- 2,2 dimethylcyclopropane carboxylate**	18.1%
Inert Ingredients	81.9%
	100.0%

**Contains 1.5 pounds active ingredient per gallon.

**Cis/trans ratio: Max. 55% (±) cis and min. 45% (±) trans
U.S. Patent No. Pending

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.



FMC Corporation
Agricultural Products Group
1735 Market Street
Philadelphia PA 19103

4 of 27

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

Note to Physician: Vomiting should be supervised by a physician or the professional staff because of the possible pulmonary damages by aspiration of the solvent.

For Emergency Assistance Call (800) 331-3148.

See other panels for additional precautionary information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. The active ingredient may cause sensitization reaction in some individuals.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Handlers who may be exposed to the dilute through application or other tasks must wear: Long-sleeved shirt and long pants, Chemical-resistant gloves, such as Barrier Laminate or Viton, and Shoes plus socks.

Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear: Long-sleeved shirt and long pants, Chemical-resistant gloves, such as Barrier Laminate or Viton, Shoes plus socks, and Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

This pesticide is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

Physical/Chemical Hazards

Do not use or store near heat or open flame.

50427

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Resistance. Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, Chemical-resistant gloves, such as Barrier Laminate or Viton, and Shoes plus socks.

STORAGE AND DISPOSAL

Pesticide Storage

Store in a cool, dry, well-ventilated place. Do not store below -6.6°C (20°F). If solids are observed warm to above 4.4°C (40°F) and roll or shake containers to redissolve. Do not use near heat, open flame or hot surfaces. Store in original containers only. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

Keep out of reach of children and animals.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC: (800) 331-3148.

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

Container Disposal

Metal Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

Plastic Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Sealed Containers: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must also contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Fury® insecticide should be applied continuously for the duration of the water application. Fury® should be diluted in sufficient volume to insure accurate application over the area to be treated. Use the

7 of 27

appropriate amount of water to carry the product to the target pest.
Agitation is not required when a suitable diluent is used.

GENERAL INSTRUCTIONS

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting. Do not exceed maximum allowable rate.

Preventive Use

For cutworm, armyworm, or stalk borer control, Fury® insecticide may be applied before, during, or after planting. For soil-incorporated applications, use higher rates for improved control. Do not exceed maximum allowable rate.

Rotational Crops

With the exception of the crops listed below, rotational crops should not be planted within 30 days of last application.

Tank-Mixture

Fury® Insecticide may be applied in tank mixtures with other products approved for use on Alfalfa and Nongrass Animal Feeds; Berries; Brassica Vegetables; Bulb Vegetables; Canola (Rapeseed); Corn; Cotton; Cucurbit Vegetables; Fruiting Vegetables; Grapes; Grass Forage, Fodder and Hay and Grass Grown for Seed; Leafy Vegetables; Legume Vegetables; Peanut; Pome Fruits; Rice; Root and Tuber Vegetables; Sorghum; Soybeans; Stone Fruits; Sugarcane; Sunflower; Tree Nuts; and Wheat. Observe all restrictions and precautions which appear on the labels of these products. Test for compatibility of products before mixing.

BUFFER ZONES

Vegetative Buffer Zones

Construct and maintain a minimum 10-foot-wide vegetative filter strip of grass or other permanent vegetation between the field edge and down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing zeta-cypermethrin onto fields where a maintained vegetative buffer strip of at least 10 feet exists between the field and down gradient aquatic habitat.

For guidance, refer to the following publication for information on constructing and maintaining effective buffers: *Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. USDA, NRCS. 2000. Fort Worth, Texas. 21pp.*
<http://www.in.nrcs.usda.gov/technical/agronomy/newconbuf.pdf>

Buffer Zone for Ground Application (groundboom, overhead chemigation, or airblast) – Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for ULV Aerial Application – Do not apply within 450 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Buffer Zone for Non-ULV Aerial Application – Do not apply within 150 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, estuaries, and commercial fish ponds).

Spray Drift Requirements

Wind Direction and Speed

Only apply this product if the wind direction favors on-target deposition. Do not apply when the wind velocity exceeds 15 mph.

Temperature Inversion

Do not make aerial or ground applications into temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

Droplet Size

Use only Medium or coarser spray nozzles (for ground and non-ULV aerial application) according to ASAE (S572) definition for standard nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

Additional Requirements for Ground Applications

Wind speed must be adjacent to the application site on the upwind side, immediately prior to application.

For ground boom applications, apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

Additional Requirements for Aerial Applications

The spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of the wing span or 80% rotor diameter.

9a27

Flight speed and nozzle orientation must be considered in determining droplet size.

Spray must be released at the lowest height consistent with pest control and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Maximum Usage When Applying Both Zeta-Cypermethrin and Cypermethrin Products to the Same Crop Within the Same Season.

Do not apply more than the maximum seasonal total for either product when used alone, and do not apply more than the combined maximum seasonal total for both products as outlined in the table below.

Crop	Maximum Seasonal Total for Either Product Used Alone (pounds active ingredient/acre)		Maximum Seasonal Total When Applying Both Products to the Same Crop (pounds active ingredient/acre)
	Zeta-cypermethrin (Fury or Mustang)	Cypermethrin (Ammo)	Zeta-cypermethrin (Fury or Mustang) plus Cypermethrin (Ammo)
Cotton	0.3	0.6	0.6
Head and Stem Brassica	0.3	0.6	0.6
Leafy Brassica Greens	0.3	0.4	0.4
Head Lettuce	0.3	0.6	0.6
Bulb Vegetables	0.25	0.5	0.5
Pecans	0.3	0.6	0.6

* Fury or Mustang (1.5 EC or 1.5 EW); Mustang MAX (0.8 EC or 0.8 EW).

** Any cypermethrin product approved for crop use, including Ammotm

Maximum Seasonal Usage and PHI (Pre-Harvest Interval) for Fury Labeled Crops

Crop	Maximum Seasonal Total/Acre for Fury		PHI (days)
	Lbs AI	Fl oz	
Alfalfa and Nongrass Animal Feeds (Forage, Fodder, Straw and Hay) Group	0.05/cutting	4.3	3 (cutting or grazing) 7 (harvesting seed)
	0.15/season	12.9	
Berries	0.3	25.8	1
Brassica Vegetables	0.3	25.8	1
Bulb Vegetables	0.25	21.5	7
Citrus	0.2	17.2	1
Corn, sweet	0.3	25.8	3
Corn, field, seed, pop	0.2	17.2	30 (grain & stover) 60 (forage)
Cotton	0.3	25.8	14
Cucurbit Vegetables	0.3	25.8	1
Fruiting Vegetables	0.3	25.8	1
Grapes	0.3	25.8	1
Grass Forage, Fodder, and Hay Group and Grass Grown for Seed	0.05/cutting	4.3	0 (Forage and Hay) 7 (Straw and Seed Screenings)
	Hay 0.20/season	17.2	
	Forage, Straw & Seed Screenings 0.25/season	21.5	
Leafy Vegetables	0.3	25.8	1
Legume Vegetables	0.3	25.8	1 (succulent shelled or edible-podded) 21 (dried shelled)
Oilseed Commodities:			
Canola (Rapeseed)	0.3	25.8	7
Safflower	0.15	12.9	14
Sunflower	0.25	21.5	30
Peanut	0.3	25.8	7
Pome Fruits	0.3	25.8	14
Rice and Wild Rice	0.2	17.2	14
Root and Tuber Vegetables (except Sugar Beet)	0.3	25.8	1
Sorghum	0.25	21.5	14 (grain & fodder (stover)) 45 (forage (silage))
Soybeans	0.3	25.8	21

10 of 27

Stone Fruits	0.3	25.8	14
Sugarcane	0.2	17.2	21
Tree Nuts	0.25	21.5	7
Wheat	0.25	21.5	14

The REI (Restricted Entry Interval) is 12 hours for all labeled crops.
Refer to the crop specific use directions for detailed information on application timing and any use restrictions

Nongrass Animal Feeds (Forage, Fodder, Straw and Hay) Group including: Alfalfa; Alfalfa grown for seed (Includes lucerne, sainfoin, holy clover, esparcet, birdsfoot trefoil and varieties and/or hybrids of these); Velvet Bean; Clover; Kudzu; Lespedeza; Lupin; Sainfoin; Trefoil; Vetch; Crown Vetch; and Milk Vetch.

Insects Controlled	Rate of Application	Method of Application
Alfalfa Caterpillar Alfalfa Looper Alfalfa Weevil Cutworms Egyptian Alfalfa Weevil (larvae & adult) Flea Beetles Green Cloverworm Hornworms Meadow Spittlebug Potato Leafhopper Velvetbean Caterpillar Webworms Blue Alfalfa Aphid ¹ Green Peach Aphid ¹ Pea Aphid ¹ Spotted Alfalfa Aphid ¹ Threecornered Alfalfa Hopper	2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre	Apply as insects appear in sufficient volume of water to ensure thorough coverage of foliage. Use higher recommended dosage for increased pest pressure or for increased residual pest control. Do not exceed maximum allowable rate. Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment. ULV oil spray application is prohibited. Higher volumes of finished spray may improve insect control under high temperatures, when foliage is dense and/or when insect pressure is high.
Armyworms Grasshoppers Plant Bugs (including Lygus spp. & Stink Bugs)	3.0 to 4.3 ounces (0.035 to 0.05 pound active) per acre	
Do not make applications less than 7 days apart. A maximum of 4.3 ounces of product or 0.05 pounds of active ingredient/acre may be applied per cutting and a maximum of 12.9 ounces of product per acre or 0.15 pounds of active ingredient per acre per season. Applications may be made up to 3 days of cutting or grazing or up to 7 days of harvesting seed. ¹ Aphid control may be variable depending on species present and host-plant relationships.		

Berries Crop Group (1 Day PHI) including: blackberry; loganberry, red and black raspberry; blueberry, highbush and lowbush; currant; elderberry; gooseberry; huckleberry; and cultivars and/or hybrids of these.

Insects Controlled	Rate of Application	Method of Application
Leafrollers Orange Tortrix Root Weevils	4.3 ounces (0.05 pounds active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate. Apply by ground and air equipment using sufficient water to obtain full coverage of foliage (minimum of 20 gallons by ground and 2 gallons by air).

11/06/27

Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season.

Do not make applications less than seven days apart.

Head and Stem Brassica Vegetables (1 day PHI) including: Broccoli; Chinese Broccoli (gai lan, white flowering broccoli); Brussels Sprouts; Cauliflower; Cavalo broccolo; Kohlrabi; Cabbage; Chinese Cabbage (napa); Chinese Mustard Cabbage (gai choy).

Leafy Brassica Greens (1 day PHI) including: Broccoli Raab (rapini) ; Chinese cabbage (bok choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens; Turnip Greens.

Insects Controlled	Rate of Application	Method of Application
Corn Earworm Cucumber Beetles Cutworm Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Saltmarsh Caterpillar Southern Cabbageworm Tobacco Budworm	2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of Fury® should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations.
Alfalfa Looper Armyworms Cabbage Looper Cabbage Webworm Crickets Grasshoppers Ground Beetles Leafminers (adults) Lygus Bugs Onion Thrips Stinkbugs Wireworm (adults) Aphids* Whiteflies*	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required.
Do not make applications less than 7 days apart. A maximum of 25.8 ounces of product or 0.3 pounds of active ingredient may be applied per acre per season. *Aids in control		

Bulb Vegetables (Allium spp.) (7 day PHI) including: Garlic; Garlic, Great-Headed (elephant); Green Eschalots; Japanese Bunching Onions; Leeks; Onion, Dry Bulb and Green; Onion, Welsh; Shallots, Dry Bulb and Green; Spring Onion or Scallions)

Insects Controlled	Rate of Application	Method of Application
Onion Thrips	3.2 to 4.3 ounces (0.0375 to 0.05 pound active) per acre	Apply in a minimum of 20 gallons per acre with ground equipment or in a minimum of 3 gallons per acre by aircraft. Begin applications when pests appear and repeat as necessary to maintain control.
Armyworms Cutworms Leafminers Onion Maggot Adults Stink Bugs Aphids*	2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre	To control Onion Thrips: Use higher rates as population increases and avoid rescue situations. Use of a crop oil concentrate at 16 fluid ounces per acre is recommended. Do not exceed maximum allowable rate.
Do not make applications less than 7 days apart. Do not apply more than 21.5 ounces of product or 0.25 pound of active ingredient per acre per season. Do not graze livestock in treated areas or cut treated crops for feed. *Aids in control		

Citrus Fruits Crop Group (1 Day PHI) including: Calamondin (*Citrus mitis*; *Citrofortunella mitis*); Citrus citron (*Citrus medica*); Citrus hybrids (*Citrus* spp.) (includes chironja, tangelo, tangor); Grapefruit (*Citrus paradisi*); Kumquat (*Fortunella* spp.); Lemon (*Citrus jambhiri*, *Citrus limon*); Lime (*Citrus aurantiifolia*); Mandarin (tangerine) (*Citrus reticulata*); Orange, sour (*Citrus aurantium*); Orange, sweet (*Citrus sinensis*); Pummelo (*Citrus grandis*, *Citrus maxima*); and Satsuma mandarin (*Citrus unshiu*).

Insects Controlled	Rate of Application	Method of Application
Asian Cockroach Beet Armyworm Blue-Green Citrus Root Weevils Cutworms Diaprepes Root Weevil Fire Ants Fuller Rose Beetle Glassy-Winged Sharpshooter Grasshopper Katydid Leafhoppers Leafrollers Leafminers* Little Leaf Notcher Loopers Orange Tortrix Orangedog Caterpillars Plantbugs Psyllids Thrips Whiteflies	4.3 ounces (0.05 pound active) per acre	Apply by ground equipment using sufficient water to obtain full coverage of foliage in a minimum of 20 gallons for concentrate spray or a minimum of 100 gallons for dilute spray. Apply by air in a minimum of 10 gallons per acre. Begin applications when pest activity is noted.
Do not apply more than 17.2 ounces of product or 0.20 pounds of active ingredient per acre per season. Do not make applications less than fourteen days apart.		

Corn, Sweet (3 day PHI)

Insects Controlled	Rate of Application	Method of Application
Chinch Bug Corn Rootworm (Adult) Corn Silkfly Cutworms Flea Beetle Leafhoppers Japanese Beetle (Adult) Sap Beetle (adults) Tarnished Plant Bug	2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre	Apply with ground or air equipment using sufficient water and application methods to insure thorough coverage of foliage. Apply in water using a minimum of 20 gallons of finished spray per acre with ground equipment and a minimum of 2 gallons per acre by air.
Armyworms Corn Borers Corn Earworm Grasshoppers Aphids*	3.0 to 4.3 ounces (0.035 to 0.05 pound active) per acre	
Apply at minimum 3 to 5 day intervals or as needed for control. A maximum of 28.5 ounces of product or 0.3 pounds of active ingredient per acre per season may be applied. *Aids in control		

Corn (Field), Field Corn Grown for Seed, Popcorn (At Plant Use)

Insects Controlled	Rate of Application	Method of Application
Cutworms	0.16 fluid ounces per 1,000 linear 4' band.	Apply as an in-furrow, band or T-band treatment using a minimum 4' band. Use table below to

Fury EC At Plant Use Amendment 06-17-09

	feet of row (0.002 pound active) per 1,000 linear feet of row	determine the Fury®needs for each acre. Do not apply more than 0.20 pound active ingredient per acre per season including at-plant plus foliar applications of Fury®. Do not apply within 30 days of harvest for grain and fodder (stover) and 60 days for forage (silage).		
Row Spacings (inches)		40	30	20
Fury 1.5 EC (pounds ai per acre)		0.024	0.036	0.048
Fury 1.5 EC (formulated ounces per acre)		2.05	3.08	4.10

Corn (Field), Field Corn Grown for Seed, Popcorn

Insects Controlled	Rate of Application	Method of Application
Cutworms	1.4 to 2.9 ounces (0.016 to 0.034 pound active) per acre	Make applications when insect populations reach economic threshold levels. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results. Do not exceed maximum allowable rate.
Bean Leaf Beetle Cereal Leaf Beetle Corn Rootworm Beetle European Corn Borer Flea Beetle Grasshoppers Hop Vine Borer Hornworms Japanese Beetle (adult) Sap Beetle (adult) Southern Corn Leaf Beetle Southwestern Corn Borer Stalk Borer Stink Bug Spp. Tobacco Budworm Webworms Aphids*	2.9 to 4.3 ounces (0.034 to 0.05 pound active) per acre	Apply by air or by ground equipment using sufficient water to obtain full coverage of foliage (minimum of 2 gallons per acre by air and 10 gallons per acre by ground). Do not apply more than 0.20 pound active ingredient per acre per season including At-Planting plus foliar applications of Fury® insecticide.
Armyworms (including Fall Armyworms) Chinch Bug	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	Do not apply within 30 days of harvest for grain and fodder (stover) and 60 days for forage (silage).
Corn Earworm ¹ Western Bean Cutworm ¹ Green Cloverworm Meadow Spittlebug	1.9 to 4.3 ounces (0.022 to 0.05 pound active) per acre	For chinch bug control, scout corn fields and make applications when bugs migrate from small grains or wild grasses to small corn. Direct spray to the base of plant. Repeat applications at 3 to 5 day intervals if needed. Fury® may only suppress heavy infestations and/or subsequent migrations.
*Control may be variable depending on species present and host-plant relationships.		
¹ For control before the larva bores into the plant stalk or ear.		

Cotton (14 day PHI)

Insects Controlled	Rate of Application	Method of Application
Preemergent Use: Cutworms	1.4 to 2.0 ounces (0.016 to 0.024 pound active) per acre	Use Fury® in the time period from 14 days prior to planting up to emergence of the crop. Apply as a broadcast spray by ground or air, banded (including T- band) or in-furrow spray using sufficient spray volume to achieve adequate coverage. Reduced volumes of water may be used with specialized equipment. Use a minimum of 1 gallon of water per acre by air. Use the higher rates of Fury® when incorporating into the soil.

140f27

Foliar Use: Cutworms Tobacco Thrips Soybean (banded) Thrips	1.4 to 2.0 ounces (0.016 to 0.024 pound active) per acre	Fury® may be applied in water or refined vegetable oil. When water is used, apply a minimum of one gallon of finished spray per acre by air or five gallons of finished spray with ground equipment. When applying in water by air, one quart of emulsified oil may be substituted for one quart of water in the finished spray. When using oil, use a minimum of one quart per acre in the finished spray.
Foliar Use: Boll Weevil Cabbage Looper Cotton Bollworm Cotton Fleahopper Cotton Leaf Perforator European Corn Borer Fall Armyworm Pink Bollworm Saltmarsh Caterpillar Stink Bugs Tarnished Plant Bug Other Plant Bugs Tobacco Budworm Yellow Striped Armyworm	2.8 to 3.8 ounces (0.033 to 0.045 pound active) per acre	Control of lepidopteran eggs may be achieved with proper timing of applications.
Foliar use: Lygus Bugs Beet Armyworm* Cotton Aphid** Whiteflies**	3.0 to 4.3 ounces (0.035 to 0.05 pound active) per acre	Fury® may be injected into overhead sprinkler irrigation water provided 1) an anti- backflow check valve is present between the injection port and the water source, 2) a check valve is present in the line to prevent irrigation water from entering the chemical supply tank and 3) the irrigation injection system has interlocking on- off switches.
Foliar use: Grasshoppers	3.2 to 4.3 ounces (0.0375 to 0.05 pound active) per acre	For boll weevil control, apply Fury® at a 3 to 4 day interval until pest numbers are reduced to acceptable levels. For control of grasshoppers, applications should be made based on careful field scouting. Treatment decisions should be made based on evidence of feeding damage and presence of grasshoppers in cotton. Loss of cotyledon leaves in seedling cotton should be considered more important than leaf loss in older cotton. Applications should be made on a broadcast basis since grasshopper are highly mobile. Adjust rates based on populations of grasshopper found in fields. Applications should be made on a three to five day schedule until grasshopper populations are under control or until foliage loss subsides. Increase application rates as grasshopper size and population density increases.
A maximum of 25.8 ounces of product or 0.3 pounds of active ingredient may be applied per acre per season. Do not graze or feed cotton for forage. * For control of beet armyworms only in the high plains of Texas, Arizona, and California. ** Aids in control		

Canola, Crambe, Rapeseed, Borage, Cuphea, Echium, Flax, Gold
of Pleasure, Hare's-Ear Mustard, Lesquerella, Lunaria,
Meadowfoam, Milkweed, Mustard, Oil Radish, Poppy Seed,
Sesame, and Sweet Rocket (7 Day PHI).

Insects Controlled	Rate of Application	Method of Application
-----------------------	------------------------	--------------------------

Aphids Cutworms Diamondback Moth Loopers Lepidopterous Larvae Flea Beetle Flea Hoppers Grasshopper Plant Bug Stink Bugs Seedpod Weevil Thrips Whitefly Armyworms	4.3 ounces (0.05 pounds active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season. Do not make applications less than seven days apart.		

Cucurbit Vegetables Group (1 day PHI) including: **Chayote (fruit); Chinese Waxgourd** (Chinese Preserving Melon); **Citron Melon; Cucumber; Gherkin; Gourd (edible)** (including hyotan, cucuzza, hechima, Chinese okra); **Mormordica spp.** (includes balsam apple, balsam pear, bitter melon, Chinese cucumber); **Muskmelon** (hybrids and/or cultivars of *Cucumis melo*) (includes true cantaloupe, cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, and snake melon); **Pumpkin; Summer Squash** (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow, zucchini); **Winter Squash** (includes butternut squash, calabaza, hubbard squash, acorn squash, and spaghetti squash); **Watermelon** (includes hybrids and varieties).

Insects Controlled	Rate of Application	Method of Application
Cutworm spp.	1.4 to 4.3 ounces (0.016 to 0.05 pounds active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate.
Cabbage Looper Cucumber Beetle spp. (adult) Leafhopper spp. Melonworm Pickleworm Rindworm Squash Bug Squash Vine Borer	3.0 to 4.3 ounces (0.035 to 0.05 pounds active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Aphid spp. ^{1,2} Armyworm, Beet ^{1,2} Corn Earworm Leafminer Plant Bug spp. Stinkbug spp.	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	Do not make applications less than 7 days apart.
Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season. ¹ Aids in control. ² See resistance statement under "Directions For Use" section.		

Fruiting Vegetables (except Cucurbits) (1 day PHI) including: **Eggplant; groundcherry** (*Physalis* spp.); **okra; pepino** (Melon pear); **pepper** (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); **tomatillo; tomato.**

Insects Controlled	Rate of Application	Method of Application
--------------------	---------------------	-----------------------

Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier Colorado Potato Beetle Corn Borer, European Corn Borer, Southwestern Corn Earworm Cucumber Beetle Cutworm spp. Flea Beetle Garden Webworm Green Stink Bug Hornworms Leafminer spp. Leafhopper spp. Meadow Spittlebug Pepper Maggot (adults) Pepper Weevil Plant Bug spp. Tobacco Budworm Tomato Fruitworm Tomato Pinworm	2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Aphid spp. ^{1,2} Armyworm, Beet ² Armyworm, Fall Cabbage Looper Grasshoppers Lygus Bugs Brown Stink Bug Tomato Psyllid Thrips spp. ^{1,2} Whitefly spp. ^{1,2}	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre.	
Do not make applications less than 7 days apart. Do not apply more than 25.8 ounces of product or 0.3 pounds active ingredient per acre per season. ¹ Aids in control ² See resistance statement		

Grape (1 Day PHI).

Insects Controlled	Rate of Application	Method of Application
Asian Lady Bird Beetle Lady Bird Beetle Cutworm species	2.15 to 4.3 ounces (0.025 to 0.05 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate.
Eastern Grape Leafhopper Variegated Leafhopper Western Grape Leafhopper Grape Berry Moth Japanese Beetle (adult)	4.3 ounces (0.05 pounds active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not apply more than 25.8 ounces of product or 0.30 pounds of active ingredient per acre per season. Do not make applications less than seven days apart.		

Grass Forage, Fodder, and Hay Group and Grass Grown for Seed and Pasture and Rangeland (0 day PHI for forage and hay; 7 day PHI for straw and seed screenings) including: bahiagrass, barnyardgrass, bentgrass, Bermudagrass, Kentucky bluegrass, big bluestem, smooth brome grass, buffalograss, reed canarygrass, centipedegrass, crabgrass, cupgrass, dallisgrass, sand dropseed, Kentucky fescue, meadow foxtail, eastern gamagrass, side-oats grama, guinea grass, Indian grass, Johnsongrass, lovegrass, napiergrass, oatgrass, orchardgrass, pangolagrass, paspalum, redtop, Italian ryegrass, St. Augustine grass, sprangletop, squirreltailgrass, stargrass, switchgrass, timothy, crested wheatgrass, wildrye grass and zoysia grass.

170927

Also included are sudangrass and sorghum forages and their hybrids.

Insects Controlled	Rate of Application	Method of Application
Alfalfa Caterpillar Alfalfa Looper Alfalfa Weevil Cutworms Egyptian Alfalfa Weevil (larvae & adult) Flea Beetles Green Cloverworm Hornworms Meadow Spittlebug Potato Leafhopper Velvetbean Caterpillar Webworms Blue Alfalfa Aphid ¹ Green Peach Aphid ¹ Pea Aphid ¹ Spotted Alfalfa Aphid ¹ Threecornered Alfalfa Hopper	2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre	Apply as insects appear in sufficient volume of water to ensure thorough coverage of foliage. Use higher recommended dosage for increased pest pressure or for increased residual pest control. Do not exceed maximum allowable rate. Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment.
Armyworms Cereal Leaf Beetle Chinch Bug Grass Mealybug Grasshoppers Plant Bugs (including Lygus spp. & Stink Bugs)	3.0 to 4.3 ounces (0.035 to 0.05 pound active) per acre	ULV oil spray application is prohibited. Higher volumes of finished spray may improve insect control under high temperatures, when foliage is dense and/or when insect pressure is high.
Do not make applications less than 7 days apart for forage and hay; not less than 17 days for straw and seed screenings. Do not spray livestock. Allow application to dry before letting livestock graze on treated area. A maximum of 4.3 ounces of product or 0.05 pounds of active ingredient per acre may be applied per cutting. For hay, a maximum of 17.2 ounces of product or 0.20 pounds of active ingredient per acre per season may be applied. For forage, straw, and seed screenings, a maximum of 21.5 ounces of product or 0.25 pounds active ingredient per acre per season may be applied. Applications may be made up to 0 days for forage and hay; 7 days for straw and seed screenings. ¹ Aphid control may be variable depending on species present and host-plant relationships.		

Leafy Vegetables (except Brassica) (1 day PHI): Amaranth (leafy amaranth, Chinese spinach, tampala); Arugula (Roquette); Cardoon; Celery; Celery, Chinese; Celtuce; Chervil; Chrysanthemum, edible-leaved and garland; Cilantro (not for use on cilantro grown for seed or coriander); Corn salad; Cress, garden; Cress, upland (yellow rocket, winter cress); Dandelion; Dock (sorrel); Endive (escarole); Fennel, Florence (finocchio); Lettuce, head and leaf; Orach; Parsley; Purslane, garden; Purslane, winter; Radicchio (red chicory); Rhubarb; Spinach (including New Zealand and vine, Malabar spinach, Indian spinach); Swiss chard.

Insects Controlled	Rate of Application	Method of Application
Corn Earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Saltmarsh Caterpillar Tobacco Budworm ^{2,3} Aphid spp. ^{2,3} Whitefly spp. ^{1,2}	2.4 to 4.3 ounces (0.028 to 0.05 pound active) per acre	Apply in water as necessary for insect control using a minimum of 10 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of Fury should be used under light to moderate insect pressure. Higher rates should be used

180227

Armyworms Ground Beetles Crickets Loopers Lygus Bugs Onion Thrips Stink Bugs Wireworm (adults)	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	to control heavy to extremely heavy insect populations. In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required.
Do not make applications less than 7 days apart. A maximum of 25.8 ounces of product or 0.3 pounds of active ingredient may be applied per acre per season. ¹ Aids in control ² See resistance statement under "Directions For Use" section ³ Aphid control may be variable depending on species present and host-plant relationships.		

Legume Vegetables

Legume Vegetables - Dried (Except Soybeans) (At-plant Application)

Dried Shelled Pea and Bean (except Soybean)

Dried cultivars of bean (*Lupinus* spp.) includes Grain Lupin, Sweet
 Lupin, White Lupin, and White Sweet Lupin; *Phaseolus* spp. Includes
 Field Bean, Kidney Bean, Lima Bean (dry), Navy Bean, Pinto Bean,
 Tepary Bean; Bean (*Vigna* spp.) includes Adzuki Bean, Blackeyed
 Pea, Catjang, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice
 Bean, Southern Pea, Urd Bean; Broad Bean (dry); Chickpea; Guar;
 Lablab Bean; Lentil; Pea (*Pisum* spp.) includes Field Pea; Pigeon Pea.

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. White grub Wireworm spp.	4.3 fl oz/ A (0.05 lbs ai/ A)	For Cutworm spp: Apply at planting on the soil surface in a 5 – 7 inch band in a minimum of 2 – 7 gallons per acre or broadcast in a minimum of 10 gallons per acre. For White grubs and Wireworms: Apply in- furrow or in a 3 – 4 inch T- Band (band over the open furrow) at planting in a minimum of 2 – 7 gallons per acre.
PHI: Do not apply within 21 days of harvest for dried shelled peas or beans. Maximum Amount per Application: A maximum of 4.3 ounces of product or 0.05 pounds of active ingredient per acre. Maximum Amount per Season: Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season including at- plant plus foliar applications.		
Row spacing (inches)	Fl oz/ 1000 linear feet	Lbs ai/ 1000 linear feet
30	0.247	0.00287
20	0.164	0.0019
15	0.123	0.0014

Legume Vegetables (Foliar use)

Legume Vegetables - Succulent and Dried (except Soybeans)

1 day PHI for succulent shelled or edible-podded peas or beans
 21 day PHI for dried shelled peas or beans

**Succulent Edible-Podded Peas, Succulent Shelled Peas and Dried
 Shelled Peas (*Pisum* spp.) including:**

Dwarf Pea; Edible-pod Pea; Snow Pea; Sugar Snap Pea; Pigeon
 pea; English Pea; Garden Pea; Green Pea; Lentil.

**Succulent Edible-Podded Beans, Succulent Shelled Beans, and
 Dried Shelled Beans including:**

Runner Bean; Snap Bean; Wax Bean; Asparagus Bean;
 Chinese Longbean; Moth Bean; Yardlong Bean; Jackbean;
 Soybean (immature seed); Swordbean; Lima Bean; Broad Bean
 (Fava Bean); Blackeyed Pea; Southern Pea; Grain Lupin; Sweet
 Lupin; White Lupin; White Sweet Lupin; Field Bean; Kidney

19027

Bean; Navy Bean; Pinto Bean; Tepary Bean; Adzuki Bean; Catjang; Crowder Pea; Moth Bean; Mung Bean; Rice Bean; Urd Bean; Chickpea (Garbanzo Bean); Guar; Lablab bean. (Foliar use)

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Thistle Caterpillar (Painted Lady) Saltmarsh Caterpillar Silverspotted Skipper	1.4 to 4.3 ounces (0.016 to 0.05 pound active) per 1000 linear ft.	Apply as required by scouting, usually at intervals of 5 or more days. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate.
Alfalfa Caterpillar Armyworm, Southern Armyworm, True Armyworm, Yellow-Striped Bean Leaf Beetle Blister Beetle spp. Colorado Potato Beetle Corn Borer, European Corn Borer, Southwestern Corn Earworm Corn Rootworm Beetle (adult) Cowpea Curculio Cucumber Beetle Flea Beetle Green Cloverworm Ground Beetles Imported Cabbageworm Japanese Beetle Leaf Skeletonizer spp. Leafhopper spp. Leafminer spp. Mexican Bean Beetle Pea Weevil Pea Leaf Weevil Plant Bug spp. Potato Leafhopper Seedcorn Beetle Seedcorn Maggot (adult) Spittlebug Three-Cornered Alfalfa Hopper Tobacco Budworm ² Velvetbean Caterpillar Webworm spp. Woolly Bear Caterpillar	3.0 to 4.3 ounces (0.035 to 0.05 pound active) per 1000 linear ft.	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Aphid spp. ^{1,2} Armyworm, Beet ² Armyworm, Fall Grasshoppers Lesser Cornstalk Borer ¹ Looper spp. ² Stink Bug spp. Thrips spp. Whitefly spp. ^{1,2}		
Do not make applications less than 5 days apart. Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per 1000 linear ft per season. ¹ Aids in control ² See resistance statement under "Directions For Use" section		

Peanut (7 day PHI)

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Green Cloverworm Velvetbean Caterpillar Red-necked Peanut Worm	1.4 to 4.3 ounces (0.016 to 0.05 pounds active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic

Bean Leaf Beetle Leafhopper spp. Southern Corn Rootworm (adult) Vegetable Weevil Whitefringed Beetle (adult)	1.9 to 4.3 ounces (0.022 to 0.05 pounds active) per acre	threshold levels. Do not exceed maximum allowable rate.
Aphid spp. ^{1,2} Armyworm, Beet ^{1,2} Armyworm, Fall ^{1,2} Corn Earworm Grasshopper spp. Lesser Cornstalk Borer ^{1,2} Soybean Looper ^{1,2} Stink Bug spp. ² Tobacco Thrips ²	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). Do not make applications less than 14 days apart.
Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season. Do not graze livestock in treated areas. Do not use treated vines or hay for animal feed. ¹ Aids in control. ² See resistance statement under "Directions For Use" section.		

Pome Fruit Group (14 day PHI) including: Apple; Crabapple; Loquat; Mayhaw; Pear; Oriental Pear; and Quince.

Insects Controlled	Rate of Application	Method of Application
Apple Maggot Codling Moth European Apple Sawfly Green Fruitworm Japanese Beetle Lesser Appleworm Oblique Banded Leafroller Oriental Fruit Moth Pandemis Leafroller Pear Psylla Plum Curculio Potato Leafhopper Redbanded Leafroller Rosy Apple Aphid Spirea Aphid Spotted Tentiform Leafminer Stink Bugs Tarnished Plant Bug Tufted Apple Bud Moth Variegated Leafroller White Apple Leafhopper	1.4-4.3 ounces (0.016-0.05 pounds active) per acre	Begin applications at delayed dormant through first cover as common to the production areas and the target pest species. Apply in a full season spray program. Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (for ground application use a minimum of 20 gallons for concentrate spray or a minimum of 100 gallons for dilute spray; for air application use a minimum of 10 gallons). Do not make applications less than 7 days apart. Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.
Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season. Do not apply as a ULV spray. Do not feed or allow livestock to graze on cover crops from treated orchards.		

Rice and Wild Rice (14 day PHI)

Insects Controlled	Rate of Application	Method of Application
--------------------	---------------------	-----------------------

21 of 27

Fall Armyworm Grasshoppers Green Bug Leafhopper Spp. Mexican Rice Borer Rice Stalk Borer Rice Water Weevil (adult) Sugarcane Borer True Armyworm Yellow-striped Armyworm Oat Birdcherry Aphid* Wild Rice Worm	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	Apply as needed based on pest threshold levels determined by scouting practices. Refer to Extension Scouting guidelines for scouting techniques, pest thresholds and treatment timing and treatment intervals. Determine the need for repeat applications, usually at intervals of 7 days, by scouting. Do not exceed maximum allowable rate.
Chinch Bug Rice Stink Bug	2.8 to 4.3 ounces (0.033 to 0.05 pound active) per acre	<p>Fury® can be safely applied in conjunction with approved rice herbicides.</p> <p>Apply by air or ground equipment using sufficient water to obtain full coverage of foliage. When applying by air, apply in a minimum of 5 gallons of water per acre. For increased control, crop oil concentrate at 16 fluid ounces per acre may be used.</p> <p>For control of rice water weevil in dry seeded rice, make a foliar application as indicated by scouting for the presence of adults and/or feeding scars, usually within a time-frame of 0-5 days after permanent flood establishment. Do not exceed 10 days from starting permanent flood until insecticide application unless scouting indicates weevils have not been previously present. Adults may also be treated at later stages of rice development to reduce overwintering populations.</p> <p>For control of rice water weevil in water seeded rice, make the first application after flooding when scouting indicates the presence of adults and/or feeding scars. Application should usually begin when rice has emerged 0.5 inch above the waterline. Under conditions of prolonged migration into the field, start field scouting for rice water weevil adults and/or feeding scars 3-5 days after the initial treatment and, if needed, apply a second application within 7-10 days of the first application. Adults may also be treated at later stages of rice development to reduce overwintering populations.</p> <p>Green bug is known to have many biotypes. Fury® may only provide suppression. If satisfactory control is not achieved with the first application of Fury, a resistant biotype may be present. Use alternate chemistry for control.</p>
Do not make applications less than 7 days apart. Do not release floodwater within 7 days of an application. A maximum of 17.2 ounces of product or 0.20 pounds of active ingredient (1.1 pints) may be applied per acre per season. Do not use treated rice field for the aquaculture of edible fish and crustacea. Do not apply as an ultra-low volume (ULV) spray. *Aids in control		

Root and Tuber Vegetables Group 1 (except Sugar Beet) (1 day PHI) including: Arracacha; Arrowroot; Artichoke (Chinese and Jerusalem); Garden Beet; Edible Burdock; Edible Canna; Carrot;

Fury EC At Plant Use Amendment 06-17-09

Cassava (Bitter and Sweet); Celeriac (Celery Root); Chayote (Root); Turnip-Rooted Chervil; Chicory; Chufa; Dasheen (Taro); Ginger; Ginseng; Horseradish; Leren; Turnip-Rooted Parsley; Parsnip; Potato; Oriental Radish (Daikon); Radish; Rutabaga; Salsify (Oyster Plant); Black Salsify; Spanish Salsify; Skirret; Sweet Potato; Tanier (Cocoyam); Turmeric; Turnip; Yam Bean; and Yam (True).

Insects Controlled	Rate of Application	Method of Application
Cutworm spp.	1.4 to 4.3 ounces (0.016 to 0.05 pounds active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds levels. Do not exceed maximum allowable rate.
Cabbage Looper Cucumber Beetle European Corn Borer Flea beetle spp. Leafhopper spp. Southern Corn Rootworm (adult) Vegetable Weevil Whitefringed Beetle (adult)	1.9 to 4.3 ounces (0.022 to 0.05 pounds active) per acre	
Aphid spp. ^{1,2} Armyworm, Beet ^{1,2} Armyworm, Yellowstriped Cabbage Maggot Colorado Potato Beetle ² Grasshopper spp. Imported Cabbageworm Potato Leafhopper Tarnished Plant Bug	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	Do not make applications less than 4 days apart.
<p>Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season.</p> <p>Leaves of Root and Tuber Vegetables cannot be used for food or feed.</p> <p>Aids in control.</p> <p>² See resistance statement under "Directions For Use" section.</p>		

Safflower (14 day phi)

Insects Controlled	Rate of Application	Method of Application
Cutworms	4.3 ounces (0.05 pound active) per acre	<p>Apply as needed based on pest thresholds determined by scouting practices. Refer to Extension Scouting guidelines for scouting techniques, pest thresholds and treatment timing and treatment intervals. Determine the need for repeat applications, at a minimum of 14 day intervals, by scouting. Do not exceed maximum allowable rate.</p> <p>Apply with ground or air equipment using sufficient water and application methods to insure thorough coverage of foliage. Apply in water using a minimum of 15 gallons of finished spray per acre.</p>
A maximum of 12.9 ounces of product or 0.15 pounds active ingredient per acre per season may be applied.		

Sorghum (Grain) and Millet (14 day PHI for grain and stover; 45 day PHI for forage):

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Sorghum Midge	1.4 to 4.3 ounces (0.016 to 0.05 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon

23 of 27

Armyworm, Fall Armyworm, Southern Armyworm, True Armyworm, Yellow-Striped Corn Borer, European Corn Borer, Southwestern ³ Corn Earworm Flea Beetle spp. Hornworms Stink Bug spp. Webworm spp.	1.9 to 4.3 ounces (0.022 to 0.05 pound active) per acre	insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air). The addition of one to two quarts of emulsified oil per acre to the spray solution may improve spray deposition and insect control. For sorghum midge control, begin applications when 25% of the sorghum heads have emerged and are in tip bloom. Repeat applications at 10-day intervals if needed. For chinch bug control, begin applications when bugs migrate from small grains or grass weeds to small sorghum. Direct spray to the base of plants with sufficient spray volume to penetrate the soil/stem interface, leaf collars, and sheaths.
Aphid spp. ^{1,2} Armyworm, Beet ² Chinch Bug False Chinch Bug Grasshopper spp. Lesser Cornstalk Borer ³ Thrips spp. ^{1,2} Whitefly spp. ^{1,2}	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	
Do not make applications less than 10 days apart. Do not apply more than 21.5 ounces of product or 0.25 pounds of active ingredient per acre per season. ¹ Aids in Control ² See resistance statement under "Directions For Use" section ³ For control before the larva bores into the plant stalk.		

Soybeans (21 day PHI):

Insects Controlled	Rate of Application	Method of Application
Cutworm spp. Painted Lady (Thistle) Caterpillar Saltmarsh Caterpillar Silverspotted Skipper	1.4 to 4.3 ounces (0.016 to 0.05 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate.
Alfalfa Caterpillar Armyworm, Southern Armyworm, True Armyworm, Yellow-Striped Bean Leaf Beetle Blister Beetle spp. Colorado Potato Beetle Corn Borer, European Corn Earworm Corn Rootworm Beetle (adult) Cowpea Curculio Cucumber Beetle European Corn Borer Flea Beetle Green Cloverworm Hornworms Imported Cabbageworm Japanese Beetle Leaf Skeletonizer spp. Leafhopper spp. Leafminer spp. Mexican Bean Beetle Pea Weevil Plant Bug spp. Potato Leafhopper Seedcorn Maggot (adult) Soybean Aphid Spittlebug Three-Cornered Alfalfa Hopper Tobacco Budworm ² Velvetbean Caterpillar Webworm spp. Woolly Bear Caterpillar	3.0 to 4.3 ounces (0.035 to 0.05 pound active) per acre	Apply with either aerial or ground equipment using sufficient spray volume to obtain full coverage of the plant and foliage. Use a minimum of 2 gallons of finished spray per acre by air or 10 gallons of finished spray per acre by ground. The addition of one to two quarts of emulsified oil per acre to the spray solution may improve spray deposition and insect control.

24 of 27

Aphid spp. ^{1,2} Armyworm, Beet Armyworm, Fall Grasshopper spp. Lesser Cornstalk Borer ¹ Looper spp. ² Stink Bug spp. Thrips spp. Whitefly spp. ^{1,2}	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	
Do not make applications less than 7 days apart. Do not graze or harvest treated soybean forage, straw, or hay for livestock feed. Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season. ¹ Aids in control ² See resistance statement under "Directions For Use" section		

Stone Fruit Group (14 day PHI) including: Apricot; Cherry (Sweet and Tart); Nectarine; Peach; Plum (including Chickasaw Plum, Damson Plum, and Japanese Plum); Plumcot; and Prune (fresh).

Insects Controlled	Rate of Application	Method of Application
American Plum Borer Black Cherry Aphid Cherry Fruit Fly Green Fruitworm Leafrollers Leafhoppers Lesser Peach Tree Borer Peach Tree Borer Peach Twig Borer Plum Curculio Oriental Fruit Moth Rose Chafer Stink Bugs Tarnished Plant Bug Tufted Apple Budmoth Western Cherry Fruit Fly	1.4-4.3 ounces (0.016-0.05 pounds active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (for ground application use a minimum of 20 gallons for concentrate spray or a minimum of 100 gallons for dilute spray; for air application use a minimum of 10 gallons). Do not make applications less than 7 days apart.
Do not apply more than 25.8 ounces of product or 0.3 pounds of active ingredient per acre per season. Do not apply as a ULV spray. Do not feed or allow livestock to graze on cover crops from treated orchards.		

Sugarcane (21 day PHI)

Insects Controlled	Rate of Application	Method of Application
Sugarcane Borer Mexican Rice Borer	3.2 to 4.3 ounces (0.0375 to 0.05 pound active) per acre	Make applications when insect populations reach economic thresholds. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results. Do not exceed maximum allowable rate. Apply by air or ground equipment using sufficient water to obtain full coverage of foliage (minimum of 2 gallons per acre by air and 10 gallons per acre by ground).
Do not make applications less than 21 days apart. Do not apply more than 17.2 ounces of product or 0.20 pounds of active ingredient per acre per season.		

Sunflower, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Stokes Aster, Tallowwood, Tea Oil Plant, and Vernonia (30 day PHI) (At-plant Application)

Insects Controlled	Rate of Application	Method of Application
--------------------	---------------------	-----------------------

Cutworm spp. White Grub Wireworm	4.3 fl oz/A (0.05 lbs ai/A)	<p>For White grubs and Wireworms: Apply in-furrow or in a 3 – 4 inch T-Band (band over the open furrow) at planting in a minimum of 3 – 5 gallons per acre.</p> <p>For Cutworm spp: Apply at planting on the soil surface in a 5 – 7 inch band or broadcast in a minimum of 3 – 5 gallons per acre.</p>
<ul style="list-style-type: none"> • PHI: Do not apply within 30 days of harvest. • Maximum Amount per Application: Do not apply more than 0.05 lb ai/A per application. • Maximum Amount per Season: Do not apply more than 0.25 lb ai/A per season. Do not make more than 5 applications at the maximum rate per season. • Grazing: Do not graze livestock in treated areas or cut treated crops for feed. 		

Sunflower, Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Stokes Aster, Tallowwood, Tea Oil Plant, and Vernonia (30 day PHI) (Foliar use)

Insects Controlled	Rate of Application	Method of Application
Thistle Caterpillar (Painted Lady) Cutworm species	1.4 to 4.3 ounces (0.016 to 0.05 pound active) per acre	Apply with ground or air equipment using sufficient water and application methods to insure thorough coverage of foliage. Apply in a minimum of 2 gallons of finished spray per acre by aerial equipment or 10 gallons per acre by ground equipment. Begin applications when pest appears and repeat as necessary to maintain control. Do not make applications less than 7 days apart. Use higher recommended dosage for increased residual pest control. Do not exceed maximum allowable rate.
Sunflower Beetle Sunflower Moth Sunflower Maggot Stem Weevil (adult) Grasshopper species Leafhopper species Head-Clipper Weevil (adult) Red Sunflower Seed Weevil (adult) Grey Sunflower Seed Weevil (adult) Saltmarsh Caterpillar Banded Sunflower Moth Armyworm Sunflower Butterfly Woolly Bear Caterpillar Japanese Beetle Webworm species	2.72 to 4.3 ounces (0.032 to 0.05 pound active) per acre	
Long-Horned Beetle (Dectes Stem Borer adult) Beet Armyworm Fall Armyworm Stink Bug Species Pale striped Flea Beetle	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	
Do not apply more than 21.5 ounces of product or 0.25 pounds of active ingredient per acre per season. Do not make more than five applications at the maximum application rate per season. Do not graze livestock in treated areas or cut treated crops for feed Avoid applications when honey bees are actively foraging by applying during the early morning or evening hours.		

Tree Nuts Group (7 Day PHI) including: almond; beech nut; Brazil nut; butternut; cashew; chestnut; chinquapin; filbert (hazelnut); hickory nut; macadamia nut; pecan; and walnut (black and English).

Insects Controlled	Rate of Application	Method of Application
--------------------	---------------------	-----------------------

Black Pecan Aphid Codling Moth Filbert Worm Hickory Shuckworm Leaf-footed Bugs Navel Orangeworm Oblique-banded Leafroller Peach Twig Borer Pecan Leaf Casebearer Pecan Nut Casebearer Pecan Phylloxera Pecan Weevil Plant Bugs Stink Bugs Walnut Aphid Walnut Husk Fly Yellow Pecan Aphid	3.4 to 4.3 ounces (0.04 to 0.05 pounds active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic threshold levels. Do not exceed maximum allowable rate. Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Do not apply more than 21.5 ounces of product or 0.25 pounds active ingredient per acre per season.		
Do not make applications less than seven days apart.		

Wheat and Triticale (14 day PHI for grain, forage, and hay):

Insects Controlled	Rate of Application	Method of Application
Cutworm spp., including Army Cutworm Painted Lady (Thistle) Caterpillar	1.4 to 4.3 ounces (0.016 to 0.05 pound active) per acre	Apply as required by scouting. Timing and frequency of applications should be based upon insect populations reaching locally determined economic thresholds. Do not exceed maximum allowable rate.
Armyworm, Southern Armyworm, True Armyworm, Yellow-Striped Cereal Leaf Beetle Flea Beetle spp. Pale Western Cutworm Plant Bug spp. Spittlebug Webworm spp.	1.9 to 4.3 ounces (0.022 to 0.05 pound active) per acre	Apply by ground or air equipment using sufficient water to obtain full coverage of foliage (minimum of 10 gallons by ground and 2 gallons by air).
Aphid spp. ^{1,2} Armyworm, Beet ² Armyworm, Fall Chinch Bug Grass Sawfly Grasshopper spp. Greenbug ^{1,2} Stink Bug spp. Thrips spp. ^{1,2} Wheat Stem Sawfly (adult) ¹ Whitefly spp. ^{1,2}	3.4 to 4.3 ounces (0.04 to 0.05 pound active) per acre	For chinch bug control, begin applications when bugs migrate from small grains or grass weeds. Apply sufficient spray volume to penetrate the soil/stem interface, leaf collars, and sheaths.
Do not make applications less than 14 days apart. Do not apply more than 21.5 ounces of product or 0.25 pounds of active ingredient per acre per season.		
¹ Aids in Control ² See resistance statement under "Directions For Use" section		

Dealers Must Sell in Original Packages Only.
Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. EXCEPT AS WARRANTED BY THIS LABEL, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR

PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) seller or FMC, and buyer assumes the risk of any such use.

To the extent consistent with applicable law FMC or seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

FMC, Ammo, Capture, Fury, Mustang--trademarks of FMC Corporation

Asana--trademark of E.I. duPont de Nemours & Company

Ambush, Karate--trademarks of Zeneca, LTD.

Baythroid--Trademark of Bayer Aktiengesellschaft

Danitol--trademark of Sumitomo Chemical Company, LTD

Scout--trademark of Hoechst Schering Agrevo S.A.

© 2009 FMC Corporation All Rights Reserved