

Reg # 279-3126

PM-13

193

Code 1495

**RESTRICTED USE PESTICIDE**

Toxic to fish and aquatic organisms  
For retail sale to, and use only by, Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Net Contents

# Mustang™ 1.5 EW Insecticide

EPA Reg. No. 279-3126

EPA Est. 279-FL-1

**Active Ingredient**

* s-Cyano (3-phenoxyphenyl)methyl (±) cis/trans 3-(2,2-dichloroethenyl)- 2,2 dimethylcyclopropane carboxylate**	17.1%
Inert Ingredients	<u>82.9%</u>
	100.0%

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans (and Domestic Animals)**

**Warning**

May be fatal if swallowed. Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. The active ingredient may cause sensitization reaction in some individuals. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Get medical attention if irritation persists.

Applicators must wear long sleeve shirt and trousers. Mixers/loaders must wear long sleeve shirts, trousers, chemical resistant gloves and safety glasses.

Clothing which has been drenched or heavily contaminated should be disposed of in accordance with State or local regulations.

**Environmental Hazards**

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SHAKE WELL BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

**STATEMENT OF PRACTICAL TREATMENT**

**If swallowed:** Call physician or Poison Control Center. Drink plenty of water and induce vomiting by touching the back of the throat with a finger or by giving ipecac. Do not induce vomiting or give anything by mouth to an unconscious person. Do not administer milk, cream, animal or vegetable fats or alcohol as they enhance adsorption.

**If on skin:** Wash thoroughly with soap and water. Get medical attention if irritation occurs and persists.

**If in eyes:** Flush thoroughly with water. If irritation occurs and persists, contact physician.

**If inhaled:** Remove to fresh air. If breathing difficulty or discomfort occurs, obtain medical attention.

**Note to Physician:**

Skin contact (vapor or powder) may be followed by transient tingling or numbness, usually of the face, but this subsides without treatment. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive.

For Emergency Assistance Call (800) 331-3148

See other panels for additional precautionary information.

DATE APR 14 1993



FMC Corporation  
Agricultural Chemical Group  
Philadelphia PA 19103

6/92

**BEST AVAILABLE COPY**

## STORAGE AND DISPOSAL

### Pesticide Storage

Do not freeze. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Contain spills. Call FMC: (800) 331-3148.

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### Container Disposal

**Metal Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

**Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Returnable/Refillable Sealed Containers:** Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

## Chemigation Use Directions

Apply this product only through sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must also contain a functional, automatic, recirculating check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Mustang™ 1.5 EW Insecticide should be applied continuously for the duration of the water application. Mustang™ 1.5 EW should be diluted in sufficient volume to insure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of

irrigation water is recommended. Agitation is not required when a suitable diluent is used.

## GENERAL INSTRUCTIONS

### Reentry Statements

Do not treat areas while unprotected humans or domestic animals are present in the treatment area.

Do not allow entry into treated areas without protective clothing until sprays have dried.

State reentry regulations will apply if they are more restrictive than those stated on this label. Consult with State regulatory officials.

### Rotational Crops

Rotational crops may not be planted within 30 days after last application.

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

CROP	PEST	DOSAGE		REMARKS
		LB A/A	FL OZ/A	
Cotton	Cutworms Soybean (banded) Thrips	0.016 to 0.024	1.37 to 2.05	Mustang™ 1.5 EW may be applied in water with an emulsified refined vegetable oil. When water is used, apply a minimum of one gallon of finished spray per acre by air or five gallons of finished spray with ground equipment. When applying in water by air, one quart of emulsified oil may be substituted for one quart of water in the finished spray.  Control of lepidopteran eggs may be achieved with proper timing of applications.  Mustang™ 1.5 EW may be injected into overhead sprinkler irrigation water provided 1) an antibackflow check valve is present between the injection port and the water source, 2) a check valve is present in the line to prevent irrigation water from entering the chemical supply tank and 3) the irrigation injection system has interlocking on-off switches.
	Tobacco Thrips			
	Boll Weevil Cabbage Looper Cotton Bollworm Cotton Flea hopper Cotton Leaf Perforator European Corn Borer Fall Armyworm Pink Bollworm Saltmarsh Caterpillar Tarnished Plant Bug Tobacco Budworm Yellow Striped Armyworm Other Plant Bugs	0.033 to 0.045	2.82 to 3.83	
	Beet Army- worm Lygus Bugs Cotton Aphid* White Flies*	0.035 to 0.050	2.99 to 4.26	For boll weevil control, apply Mustang™ 1.5 EW at a 3 to 4 day interval until pest numbers are reduced to acceptable levels.  A maximum of 0.3 lb. active ingredient may be applied per acre per season.  Mustang™ 1.5 EW may not be applied within 14 days of harvest.  Do not graze or feed cotton for forage.

\*Aids in control.

### Cotton Spray Drift Precautions

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

Do not make aerial or ground applications during temperature inversions.

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. Boom length must not exceed 75% of wing span or rotor diameter.

Do not apply by ground equipment within 25 feet, or by air within 75 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season. Synthetic pyrethroid products include Ambush®, Ammo®, Asana® X1 Baythroid®, Capture®, Cymbush®, Mustang™, Karate®, Pource® and Scout® X-TRA.

**BEST AVAILABLE COPY**

CROP	PEST	DOSAGE*		REMARKS
		LB A/A	FLOZ/A	
Lettuce (Head)	Corn earworm Cucumber Beetles Cutworms Diamondback Moth Flea beetles Imported Cabbageworm Leafhoppers Lygus Bugs Saltmarsh Caterpillar Tobacco Budworm	0.028 to 0.050	2.39 to 4.26	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air.  Lower rates of Mustang™ 1.5 EW should be used with light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations.  A maximum of 0.3 lbs. active ingredient may be applied per acre per season. Do not apply within 5 days of harvest.
	Amyworms Crickets Loopers Onion Thrips Stinkbugs	0.040 to 0.050	3.41 to 4.26	*In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required.
Pecans	Black Pecan Aphid Hickory Shuckworm Pecan Nut Casebearer Pecan Weevil Yellow Pecan Aphid	0.033 to 0.050	2.82 to 4.26	Applications at the lower rate should be made when pest populations are low. Rates should be increased as the pest pressure increases.  Apply by ground equipment to the point of drip. Use 100 gallons of dilute spray per acre for smaller trees. For larger trees which require higher gallonage to achieve adequate coverage, apply in 200 to 300 gallons of water. In order to calculate the correct number of gallons of water needed to spray one acre of your trees to the point of drip, you may need to conduct a test. If you do not know how to conduct such a test with your equipment, you should request assistance from your equipment dealer.  Up to 0.3 lb active ingredient per acre per crop season may be applied prior to shuck split.  Do not apply within 21 days of harvest.  Do not graze livestock in treated orchards or cut treated cover crops for feed.

## Tank-Mixture

Mustang™ 1.5 EW insecticide may be applied in tank mixtures with other products approved for use on cotton. Observe all restrictions and precautions which appear on the labels of these products.

### Dealers Should Sell in Original Packages Only.

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Ambush, Cymbush and Karate—Trademarks of ICI Americas Inc.  
Asana—Trademark of E.I. duPont de Nemours and Co., Inc.  
Baythroid—Trademark of Bayer AG  
Scout—Trademark of Hoechst Roussel Agri-Vet Company  
Ammo, Capture, Mustang, Pounce and **FMC**—  
Trademarks of FMC Corporation

(1495-6/1/92-A)

**BEST AVAILABLE COPY**