

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (75-767) WASHINGTON, DC 20460	EPA REGISTRATION NO.	DATE OF ISSUANCE
	279-3151	JAN 9 1995
	TERM OF ISSUANCE	
NOTICE OF PESTICIDE: <input type="checkbox"/> REGISTRATION <input type="checkbox"/> REREGISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	conditional	
	NAME OF PESTICIDE PRODUCT	
Command 2 ME Herbicide		

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

FMC Corporation
 Agricultural Chemical Group
 1735 Market Street
 Philadelphia, PA 19103

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase, "EPA Registration No. 279-3151" and submit five (5) copies of your final printed labeling before you release the product for shipment.

Update label to read; "Do not contaminate water when disposing of equipment washwaters" where appropriate.

PRODUCT CHEMISTRY

3. The CSF dated 9/21/94 (Basic Formulation) is not considered acceptable for the following reasons (refer to Confidential Appendix A for details):

a. The nominal concentration for the AI does not agree with the label claim as required in PR Notice 91-2.

ATTACHMENT The certified limits for the AI should be based on the nominal concentration per 40 CFR 158.175.

SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------

7505C/HICKS

BEST COPY AVAILABLE

19/15

- c. The total weight in Block #17 is in error.
- d. The flash point should be specified in Block #9.
4. The weight of the product should be placed on the label in accordance with 40 CFR 156.10(d), preferably on the bottom of the front panel.
5. The label claim for the AI does not agree with the nominal concentration specified in the CSF dated 9/21/94.
6. Series 61-1: The structural formula shown on page 4 of MRID No. 43380901 should be corrected to reflect the "2-chlorophenyl" compound.
7. Series 61-2: The EPA Reg. No. for the AI specified on page 9 of MRID No. 43380902 should be corrected to "279-3052".
8. Series 62-3: Please provide sample calculations to support the percentage AI in the sample for the GC method referenced in MRID No. 43380902.
9. Series 63-17: Storage stability tests for at least one year under ambient warehouse conditions must be conducted on the commercially packaged product. The data need not be submitted to EPA unless one or more of the deficiencies outlined in PR Notice 92-5 have been observed. The data should be available whenever EPA may need to require it.

PRECAUTIONARY LABELING

10. The acute dermal and dermal irritation tests are considered to be supplementary data, and the submission of new tests is recommended. The quality assurance statements are considered incomplete. (see attached review)
11. The precautionary statement must include:
 - a. Harmful if inhaled. Avoid breathing vapor or spraymist. Remove contaminated clothing and wash before reuse.

The label may have to be revised upon the presentation and acceptance of the outstanding data.

BEST COPY AVAILABLE

Code

Net Contents



2ME

Microencapsulated Herbicide

For Agricultural or Commercial Use Only
NOT FOR SALE OR USE IN CALIFORNIA

EPA Reg. No. 279-XXXX

EPA Est. 279-


Active Ingredient:	By Wt.
Clomazone: 2-(2-Chlorophenyl)methyl-4, 4-dimethyl-3-isoxazolidinone	22.5%
Inert Ingredients:	77.5%
	100.0%

Contains 2.0 pounds of active ingredient per gallon
U.S. Patent No. 4,405,357

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)



FMC.

FMC Corporation
Agricultural Chemical Group
Philadelphia PA 19103

12/94-DRAFT

FIRST AID

If in eyes: Flush eyes with plenty of water. Contact a medical doctor if irritation occurs and persists.

If swallowed: Contact a medical doctor or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with a finger, or by giving syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of water. Contact a medical doctor if irritation occurs and persists.

For Emergency Assistance Call (800) 331-3148

See other panels for additional precautionary information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Warning

Harmful if inhaled. Avoid breathing vapor or spray mist.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pants; Chemical-resistant gloves, such as Barrier Laminate or Viton; Shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from the area treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

TABLE OF CONTENTS	
Read all information on pages 1-6 before referring to specific crop use.	
	Page
Ingredients	1
First Aid	1
Precautionary Statements	1
Directions for Use	2
Storage and Disposal	2
Sprayer Cleanup	3
Spray Drift Precautions	5
General Mixing Instructions	3
General Application Precautions	3
General Application Instructions	4
Command Geographical Areas	6
ROW CROPS	6
Cotton	6
Soybeans	7

Physical/Chemical Hazards

Do not use or store near heat or open flame.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

This product is for use only in a monitored commercial program conducted under FMC supervision.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: if the product is soil-injected or soil-incorporated, the Worker Protection Standard under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls; Chemical-resistant gloves, such as Barrier Laminate or Viton; Shoes plus socks.

STORAGE AND DISPOSAL

Pesticide Storage

Do not freeze. Do not store below 40°F. If solid crystals are observed, warm material to above 60°F by placing container in warm location. Shake or roll container periodically to redissolve solids.

Keep out of reach of children and animals. Store in original containers only. Store in a dry place. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC (800) 331-3148.

To confine spill: Dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Sealed Containers

Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

IMPORTANT

BEFORE YOU OBSERVE THE GENERAL APPLICATION PRECAUTION SECTION OF THIS LABEL, PLEASE READ THE FOLLOWING INFORMATION:

Some species of trees, shrubs, flowers, and ornamentals, and vegetables are sensitive to Command 2ME herbicide. Contact with foliage or vapors may cause injury to these plants. Symptoms are generally temporary if the herbicide is washed off some plants.

Plants in the following may show symptoms of injury if they are in contact with Command 2ME herbicide:

<ul style="list-style-type: none"> Apple (the full range of varieties) Aspen Bamboo Basswood Birch Cherry (the full range of varieties) Cornus Crabapple Elm Ginkgo Japanese Maple Juniper Laburnum Lilac Maple Pear (the full range of varieties) Pine Poplar Prunella Rose Spirea Yew 	<ul style="list-style-type: none"> Tree (Evergreen) Fir Species Spruce species
<ul style="list-style-type: none"> Shrub & Vine Azalea Burning Bush Hydrangea Grape Honeysuckle Rose Yew 	<ul style="list-style-type: none"> Agonomic Crops Alfalfa Oats
<ul style="list-style-type: none"> Vegetable and Flower Plants 	<ul style="list-style-type: none"> Others Ferns Herbs Strawberry Raspberry Blackberry

Do not apply Command 2ME herbicide to non-field areas including fence rows, waterways, ditches, and roadsides.

SPRAYER CLEANUP

Do not drain or flush equipment on or near desirable trees or other plants, or in areas where their roots may extend or in locations where the chemical may be washed or move into contact with their roots. Do not contaminate any body of water including irrigation water that may be used on other crops. Carefully follow sprayer clean-up instructions noted below to prevent spray tank residue from damaging other crops.

Sprayer equipment should be thoroughly rinsed to remove residues of herbicide that might injure other subsequently sprayed crops. The steps below should be followed for the thorough rinsing of spray equipment following applications of Command 2ME herbicide or tank mixes of Command 2ME with other labeled products.

- 1) Drain any remaining spray solution from tank, pump, hoses and boom and discard in an approved manner (See Note that follows).
- 2) Clean tank and fittings by:
 - Thoroughly hosing down the inside walls of the spray tank with a quantity of water equal to 1/3 of the total tank capacity and operating the pump to circulate this solution through the sprayer system for 15 minutes.
 - Washing down the outside surfaces of equipment.
 - Removing nozzle tip and screen from end nozzle in each boom section and allowing several gallons of rinsate solution to flush completely through boom (collect rinsate while flushing).
- 3) Thoroughly drain remaining rinsate solution from tank, pump and hoses. Combine with boom flushing and dispose of all rinsates from this first rinsing in an approved manner (see Note that follows).
 - When switching from water dilutions to applications utilizing crop oil or liquid fertilizer as a carrier, a small volume of crop oil or liquid fertilizer should be flushed through the tank, pump, hoses, and boom prior to the next use. Dispose of crop oil or liquid fertilizer rinsate in an approved manner (see Note for local, state and federal guidelines).
- 4) Remove the remaining nozzle tips, and screens and the line filter and wash in a pail of warm soapy water, thoroughly rinse and replace.
- 5) Hose down the inside walls of the spray tank a second time and circulate this solution using the same procedure as noted in #2 above.
- 6) If the next use of the sprayer will be for applying a preemergent or preplant incorporated pesticide on any crop for which Command 2ME is registered, rinsate from this second rinsing may be utilized by diluting with the water for the next pesticide load;

HOWEVER

If the next use of the sprayer will be a postemergence applied pesticide on any crop, drain rinsate solution from this second rinsing. Retain rinsate solution for use only with a soil incorporated pesticide to be applied on a crop for which Command 2ME is registered. Refill tank (after draining second rinsate solution) in accordance with post-emergence product label directions.

NOTE: Dispose of excess spray mixture and/or rinsate from first tank rinsing by application to cropland as described on this label. If excess spray mixture and/or rinsate from first rinsing cannot be disposed of according to label instructions, dispose of in compliance with local, state and federal guidelines. Contact your state pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA regional office for guidance.

GENERAL MIXING INSTRUCTIONS

Care must be taken when mixing Command 2ME herbicide. Avoid mixing in areas adjacent to desirable plants (See Section headed "IMPORTANT")

Command 2ME Alone: Mix Command 2ME with water or liquid fertilizer in the following manner: Fill the spray tank one-half to three-fourths full with water or liquid fertilizer, add the proper amount of Command 2ME, then add the rest of the water or liquid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform spray mixture.

Tank Mixtures: Fill spray tank one-fourth to one-third full with water, with agitator operating add the recommended amount of ingredients using the following order: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's). Mix thoroughly and fill tank one-half full continuing agitation. Add Command 2ME herbicide to tank while maintaining agitation. Complete filling the sprayer tank with water. Where use of a surfactant is recommended, add as the last ingredient to the spray tank. Maintain agitation during filling, mixing and application. When using drift reducing agents, follow specific product label instructions for order of addition to spray tank.

Fertilizer Spray Mixtures: Applications of Command 2ME alone, or with recommended tank mixtures, in conjunction with fertilizer solutions may be used unless use directions specifically state otherwise. Small quantities should be tested for compatibility by the following procedure before mixing tankful quantities.

- 1) Put 1 pint of fertilizer solution in a quart jar.
- 2) Add the appropriate amount of herbicide based on the table at the top of page 3. If more than one product is to be used, add each separately using the following sequence: dry formulations (e.g., wettable powders, dry flowables) first, liquid suspensions (e.g., flowables) next and finally liquids (e.g., EC's).

Herbicide	Rate/Acre	Amount To Be Added Per Pint of Fluid Fertilizer*
Wettable Powders or Dry Flowables (Dry Granules)	1/2 pound 1 pound 2 pounds 3 pounds	1/4 level teaspoon 1/2 level teaspoon 3/4 level teaspoon 1 level teaspoon
Emulsifiable Concentrates or Flowables	1 pint 1 quart 2 quarts 3 quarts	1/2 level teaspoon 1 level teaspoon 2 level teaspoons 3 level teaspoons

*Based on a spray volume of 25 gpa. For lower or higher spray volumes, adjust fluid fertilizer quantity accordingly.

- 3) Close jar and shake well.
- 4) Watch mixture for several seconds, again in 5 minutes and again after 30 minutes. If herbicide/fertilizer combination remains mixed or can be remixed readily (i.e., does not permanently separate, foam, gel or become lumpy), the mixture is compatible and can be mixed in full volumes and sprayed. If the mixture is compatible, prepare spray by adding fertilizer solution to the tank first, then follow directions noted below:
 - 1) **Command 2ME Alone:** Dilute with 2 parts of water and add slowly to spray tank of fertilizer with agitator running. Maintain agitation during application.
 - 2) **Command 2ME plus Tank Mixtures:** Dilute individual products with two (2) parts of water, then add to the spray tank of fertilizer, while maintaining agitation, using the following order—slurry of dry formulations (wetable powders, dry flowables) first, diluted liquid formulations (EC's, flowables) second. Continue agitation during application.

GENERAL APPLICATION PRECAUTIONS

Do not apply aerially or through irrigation equipment.

For Preemergent Applications:

Command 2ME must be used as a preemergent soil applied treatment without incorporation. Application precautions must be taken as follows:

Do not apply Command 2ME herbicide to non-cropland areas including roads, waterways, ditches, and borders.

Do not apply Command 2ME within 1,500 feet of the following areas: Town and Subdivisions, Commercial Fruit/Nut or Vegetable Production, Commercial Greenhouses or Nurseries.

Sensitive plants may show symptoms of foliar whitening or yellowing if contacted by Command 2ME herbicide. Prior to application, adjacent properties must be checked, and spraying within 1,500 feet of desirable plants must be avoided.

Refer to individual crop use directions for additional requirements.

©1991 Dow Agrosciences

SPRAY DRIFT PRECAUTIONS

Non-target spray drift of Command 2ME herbicide should be avoided to prevent whitening of desirable plants. Drift is influenced by many factors which include wind speed, spray pressure, particle size, nozzle type, and boom height. The use of an agriculturally approved drift reducing additive is recommended. Follow specific product label instructions for the order of addition of all products to the spray tank. Small quantities should be tested for compatibility before mixing tankful quantities.

- Do not apply when weather conditions favor drift.
- Do not apply in winds above 10 miles per hour.
- A minimum spray volume of 10 gallons per acre is recommended with appropriate nozzle types and sizes that produce coarser sprays.
- Use minimum nozzle pressure and boom height while maintaining uniform spray pattern.
- Do not exceed 30 psi spray pressure.
- Selection and proper use of spray equipment is critical in minimizing spray drift. The following table suggests pressures, flow rates, and nozzle sizes for drift reduction using various nozzle types.
- When moving spray equipment to noncontiguous sites, do not allow spray solution to spray or drip from tanks, hoses, fittings or spray nozzles and tips.

Suggested Nozzle Types, Minimum Size and Recommended Pressure Ranges for Minimizing Drift (See Special Precautions in the cotton use section or further restrictions)

Nozzle	Pressure Range (PSI)	Minimum Flow Rate Within Pressure Range (GPM)	Minimum Nozzle Size
Flat-fan	15-30	0.3	#4*
LP-flat-fan	10-25	0.3	#3
Even flat-fan**	15-30	0.3	#4
Flood	10-25	0.3	#2.5*
Whirl-chamber	5-20	0.3	#5
Raindrop	15-30	0.15	#2
Wide angle full cone	15-30	0.3	#5

* Refers to tip number such as 8004 or LF2.5.
 ** Recommended for banded application.

ROTATIONAL CROPPING PRECAUTIONS

Under some conditions, temporary whitening or yellowing of leaves may occur on approved rotational crops where undesirable soil residues of Command 2ME exist.

Under abnormal conditions, carryover injury to rotational crops can occur. The following factors can contribute to increased risk of injury to rotational crops:

- 1) Exceeding label recommended rates.
- 2) Overapplication resulting from use of worn nozzles, excessive overlapping of spray swaths, failing to shut off spray booms when turning (end row areas), or slowing or stopping sprayer.
- 3) Soil with pH less than or equal to 5.9.
- 4) Extreme dryness in the four months following application.
- 5) Use of organophosphate soil insecticide followed by use of some postemergence corn herbicides.
- 6) Choice of rotational crop hybrid.

Additional recommendations to prevent rotational crop injury may be provided in the form of service bulletins for location where risk of injury is significantly increased due to extremely dry conditions.

Refer to Rotational Crop Guidelines and Replanting Instructions of specific crops for additional crop planting information.

SPECIAL PRECAUTION

Off-site movement of spray drift or vapors of Command 2ME herbicide can cause foliar whitening or yellowing of some plants. Prior to making applications read and strictly follow all precautions and application instructions on this label.

GENERAL APPLICATION INSTRUCTIONS

Ground Applications

Broadcast or Banded Applications: Apply Command 2ME alone or in tank mix combinations by ground equipment using a finished spray volume of 10 to 40 gallons of water per acre. NOTE: The use of an agriculturally approved drift reducing additive is recommended. Use nozzles suitable for broadcast boom or banded application of herbicides. Coarse sprays are less likely to drift out of the target area than fine sprays. See "APPLICATION PRECAUTIONS" section for specific recommendations to reduce spray drift. For Command 2ME tank mixtures with wettable powder or dry flowable formulations, nozzle screens and strainers should be no finer than 50-mesh.

Banded Applications—Calculate the rates and volumes required by using the following formulas:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Rate Per Acre} = \text{Band rate per acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast Volume Per Acre} = \text{Band volume per acre}$$

Cultivation After Planting

Soil treated with Command 2ME may be shallow-cultivated, rotary-hoed or hand-hoed without reducing the weed control activity of Command 2ME. Avoid deep cultivation which may bring untreated soil to the surface and result in poor weed control.

COMMERCIAL IMPREGNATION AND APPLICATION OF COMMAND 2ME ON DRY BULK FERTILIZERS

Command herbicide may be impregnated on dry bulk fertilizers. When applied as directed Command/dry bulk fertilizer mixtures provide weed control equal to that provided by the same rates of Command applied in water.

Impregnation: Apply using a minimum of 200 pounds of dry bulk fertilizer per acre and up to a maximum of 450 pounds per acre with the recommended amount of Command 2ME herbicide per acre. Use a closed rotary-drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform, fine spray pattern over the tumbling fertilizer for thorough coverage. The physical properties of fertilizers vary, particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Command provides a satisfactory, dry mixture. If the absorptive capacity is inadequate use of a highly absorptive powder is required to provide a dry, flowable mixture. Microcel E (Johns-Manvill Products Corporation) is a recommended absorbent powder. Generally less than 2% by weight of Microcel E is required. DO NOT impregnate Command 2ME onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the herbicide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with Command.

The amount of Command actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with Command 2ME herbicide should be applied immediately, not stored. All state regulations, labeling, etc., of the mixtures are the responsibility of the seller.

For those rates not listed in the following table, calculate the amount of Command to be impregnated on a ton of dry bulk fertilizer using the following formula:

$$\frac{200 \text{ pounds dry fertilizer per acre}}{\text{PINTS of Command per acre (recommended rate for soil texture)}} \times \text{PINTS of Command per acre} = \text{PINTS of Command per ton of fertilizer}$$

RATE CHART FOR IMPREGNATION OF DRY BULK FERTILIZERS WITH COMMAND 2ME HERBICIDE

PINTS OF COMMAND 2ME PER TON OF FERTILIZER


Fertilizer Rate Lbs./Acre	Command 2ME Rate Per Acre		
	3.0 PT. (.75 lbs A.I.)	4.0 PT. (1 lb A.I.)	5.0 PT. (1.25 lbs A.I.)
200	30	40	50
250	24	32	40
300	20	26 2/3	33 1/3
350	17	23	28
400	15	20	23
450	13 1/2	18	22

If fertilizer materials are excessively dusty, use diesel oil or other suitable additive to reduce dust prior to impregnation as dusty fertilizer will result in poor distribution during application. Crop injury and/or poor weed control may occur where the impregnated fertilizer is not uniformly applied.

SPECIAL PRECAUTIONS FOR DRY BULK FERTILIZER IMPREGNATION AND APPLICATION WITH COMMAND 2ME

- All equipment used to apply Command must be thoroughly cleaned immediately following use to ensure no contamination results which could cause injury to non-labeled crops or desirable vegetation. Refer to "Sprayer Cleanup" section for additional details and disposal of rinsates.
- Equipment used to physically transport Command treated fertilizer, including boots, augers, conveyers, bins, etc. must be covered to prevent loss of fine particles and subjected to cleanup procedures previously described.
- Mixing of Command herbicide and dry fertilizer must be conducted in a blender which will ensure that airborne particles and potential for volatilization are avoided.
- Bulk containers must be tightly covered while the product is being transported and applied to reduce chances of Command volatilization or product loss.
- It is recommended that Dry Bulk Fertilizer impregnation operations using Command not be conducted within 1/4 mile of residential areas or areas where Command symptomology on desirable vegetation would not be tolerated.

COMMAND GEOGRAPHICAL AREAS For Rates, Weed Control, and Crop Rotation




* Southern area includes Missouri bootheel which incorporates the following counties: Butler, Dunklin, Mississippi, New Madrid, Pemiscot, Scott, and Stoddard.

COTTON

GENERAL INFORMATION

Command 2ME Herbicide may be applied, as a preemergent banded or broadcast treatment in conventional tillage or conservation tillage systems for the control of annual grasses and broadleaf weeds in cotton. Do not incorporate Command 2ME or use in a tank mixture with any herbicide that requires soil incorporation.



REQUIREMENTS FOR PLANTING TIME APPLICATIONS

Do not apply Command 2ME Herbicide to cotton unless either disulfoton or phorate organophosphate insecticide is applied in-furrow with the seed at planting time at a minimum of 0.75 pound per acre of active ingredient. Do not reduce the application rate of the organophosphate insecticide when Command 2ME herbicide is applied as a banded treatment. Failure to apply either disulfoton or phorate insecticides with Command in accordance with in-furrow label use directions can result in crop phytotoxicity (bleaching) and/or stand reduction. Combinations of at planting systemic granular carbamate and organophosphate insecticides in conjunction with Command may result in injury to cotton. Crop injury may occur with higher Command rates on sandy soils. Diuron is not recommended at planting when Command is used as plant injury may result.

Refer to the insecticide product labels for appropriate in-furrow application directions and maximum use rates. Monitor application equipment to insure accurate and uniform placement of the insecticide.

PREEMERGENT BANDED APPLICATIONS

For preemergent banded treatments, refer to the directions for banded applications in the GENERAL APPLICATION INSTRUCTIONS section.

REPLANTING INSTRUCTIONS

If the initial planting of cotton fails to produce a uniform stand, cotton may be replanted in fields treated with Command 2ME alone. Do not retreat fields with a second application of Command 2ME. When tank mixing with a labeled product, refer to the replant instructions for that product. Do not replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP GUIDELINES on the Command 2ME label. Where a tank mix is used, refer to the product's labels for any additional replant instructions.

If replanting is required follow the directions under REQUIREMENTS FOR PLANTING TIME APPLICATIONS noted above.

Command 1.5ME Herbicide Applied Alone

RATES AND WEEDS CONTROLLED¹

2.0 PT. ² (0.50 LB. A.I.)	3.0 PT. (0.75 LB. A.I.)	4.0 PT. (1.0 LB. A.I.)	5.0 PT. (1.25 LB. A.I.)
GRASSES	GRASSES	GRASSES	GRASSES
Suppression of some annual grasses	Barnyardgrass Broadleaf Signalgrass Crabgrass —Large —Smooth Foxtail —Giant —Green Goosegrass Panicum —Common —Fall —Texas Seedling Johnsongrass Field Sandbur*	All grasses controlled at 1.5 pints plus: Cupgrass —South-western —Woolly Field Sandbur Bermudagrass* Red Rice Itchgrass ³	All grasses controlled at 2.0 pints plus: Foxtail —Purple —Robust —Yellow Junglerice* Shattercane* Wild Proso Millet*

2.0 PT. ² (0.50 LB. A.I.)	3.0 PT. (0.75 LB. A.I.)	4.0 PT. (1.0 LB. A.I.)	5.0 PT. (1.25 LB. A.I.)
BROADLEAVES	BROADLEAVES	BROADLEAVES	BROADLEAVES
Velvetleaf Spurred Anoda	Velvetleaf Spurred Anoda plus: Common Ragweed* Jimsonweed* Lambsquarter Pennsylvania Smartweed* Prickly Sida Purslane Redweed Venice Mallow	All broadleaves controlled at 1.5 pints plus: Cocklebur Common Ragweed Dayflower Florida Beggarweed Florida Pusley Jimsonweed Kochia* Redvine* Tropic Croton Wild Poinsettia ⁴	All broadleaves controlled at 2.0 pints plus: Balloonvine* Black Nightshade* Curly Dock Joint Vetch* Kochia Morningglory —Pitted Pennsylvania Smartweed Pigweed —Redroot* —Smooth* Prostrate Spurge

*Partially Controlled
¹Higher label rates (a minimum of 4 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, a minimum application rate of 5 pints per acre for cocklebur or common ragweed control is recommended.
²For heavy weed pressure or heavy soils use 4 pints per acre to control Velvetleaf or Spurred Anoda.
³For control of itchgrass, do not incorporate Command 4 EC. For high weed pressure and heavy soils use 5 pints.
⁴Use in AR/LA/MS/TX/OK only—for high pressure and heavy soils use 5 pints.

COMMAND 2ME HERBICIDE MAY BE APPLIED WITH OTHER PREEMERGENT HERBICIDES REGISTERED FOR THESE USES ON COTTON.

Command 2ME may be applied in combination with other herbicides registered for use in cotton to broaden the weed control spectrum. Refer to the labels of the other herbicides for additional information. Carefully observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur. Refer to sections headed "Rotational Cropping Precautions" and "General Information."

Note: When using Command 2ME with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

2.0 PT. (0.50 LB. A.I.)	3.0 to 4.0 PT. (0.75 to 1.0 LB. A.I.)	5.0 PT. (1.25 LB. A.I.)
ANYTIME	ANYTIME	ANYTIME
Cotton* Soybeans Peppers Pumpkins Peas	Cotton* Soybeans Pumpkins Peppers	Cotton* Soybeans
9 MONTHS	9 MONTHS	9 MONTHS
Corn —Field —Pop —Seed —Sweet Cotton Cucurbits Dry Beans Peanuts Potatoes Pumpkins Rice Snap Beans Sorghum Suga Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)	Same as 2.0 pint, plus: Peas	Same as 3.0 to 4.0 pints, plus: Pumpkins
12 MONTHS	12 MONTHS	12 MONTHS
All crops	All crops	All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 2ME treatment.

*Refer to the section "Requirements for Planting Time Applications" and "Replanting Instructions."

RESTRICTIONS

- Do not apply Command within 65 days of harvest.
- Do not apply more than 1.25 pounds active ingredient per acre per season.
- Do not allow livestock to graze on treated cotton forage or trash, or feed treated cotton forage or trash to livestock.

SOYBEANS

GENERAL INFORMATION

Command 2ME herbicide may be utilized as a preemergent soil applied treatment for the control of annual grass and broadleaf weeds in soybeans.

Command 2ME herbicide may be tank mixed with other preemergent soybean herbicides to broaden the weed control spectrum compared to the products applied alone. Command 2ME may be tank mixed with Canopy®, Dual®, Lasso®, Lexone®, Lorox®, Lorox Plus®, Preview®, Prowl®, Scepter®, Sencor®, Sonalan®, Squadron®, or other registered herbicides. Observe all precautions instructions, and rotational cropping guidelines of each product's label when tank mixing, including all references to potential carryover and crop injury warnings or restrictions.

Water or liquid fertilizer may be used as a carrier for Command 2ME when applied alone, or when tank mixed with the herbicides listed above, unless use directions specifically state otherwise.

REPLANTING INSTRUCTIONS

If initial seeding of soybeans fails to produce a stand, soybeans may be replanted in fields treated with Command 2ME alone (or with recommended tank mixtures). Do not retreat field with a second application of Command 2ME. When tank mixing with a labeled product refer to the soybean replant instructions for that product. Do not replant treated fields with any crop at intervals which are inconsistent with the Rotational Crop Guidelines on this label.

Northern Area—See Map
COMMAND® 2ME HERBICIDE
APPLIED ALONE
RATES AND WEEDS CONTROLLED¹

2.0 PT. ² (0.50 LB. A.I.)	3.0 PT. (0.75 LB. A.I.)	4.0 PT. (1.0 LB. A.I.)
GRASSES Suppression of some annual grasses	GRASSES Barnyardgrass Crabgrass —Large —Smooth Field Sandbur Foxtail —Giant —Green —Robust —Yellow Goosegrass Paricum —Common —Fall —Texas Seedling Johnsongrass [*]	GRASSES All grasses controlled at 3.0 pints plus: Bermudagrass [*] Broadleaf Signalgrass Cupgrass —Southwestern —Woolly [*] Seedling Johnsongrass Shattercane [*] Wild Proso Millet [*]
BROADLEAVES Velvetleaf Spurred Anoda	BROADLEAVES Velvetleaf Spurred Anoda plus: Common Ragweed [*] Galinsoga Jimsonweed [*] Lambsquarters Prickly Sida Purslane Smartweed [*] —Ladysthumb [*] —Pennsylvania [*] Venice Mallow	BROADLEAVES All broadleaves controlled at 3.0 pints plus: Black Seeded Plantain Cocklebur [*] Common Ragweed Jimsonweed Kochia Smartweed —Ladysthumb —Pennsylvania Tropic Croton

^{*}Partially Controlled
¹Higher label rates (a minimum of 4.0 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types.
²For heavy weed pressure or heavy soils use 2.5 pints per acre to control Velvetleaf or Spurred Anoda.
NOTE: When using Command 2ME more than 30 days prior to planting, the highest label rate should be used.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur. Refer to sections headed "Rotational Cropping Precautions" and "General Information."

NOTE: When using Command 1.5ME with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

2.0 PT. ² (0.50 LB. A.I.)	3.0 PT. (0.75 LB. A.I.)	4.0 PT. (1.0 LB. A.I.)
ANYTIME Soybeans Cotton [*] Peppers Pumpkins Pears	ANYTIME Soybeans Cotton [*] Peppers Pumpkins	ANYTIME Soybeans Cotton [*] Peppers Pumpkins
9 MONTHS Corn —Field —Pop —Seed —Sweet	9 MONTHS Same as 2.0 pint, plus: Pears	9 MONTHS Same as 4.0 pints, except Seed Corn and Sweet Corn require a 12 month rotation.
Cotton Cucurbits Dry Beans Peanuts Peppers Potatoes Rice Snap Beans Sorghum Sugar Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)		
[*] Refer to the "Requirements for Planting Time Applications" and "Replanting Instructions" in the Cotton Directions for Use section.		
12 MONTHS Cabbage Tomatoes (all) Wheat	12 MONTHS Same as 2.0 pint.	12 MONTHS Same as 3.0 pints, plus: Seed Corn Sweet Corn
16 MONTHS All crops	16 MONTHS All crops	16 MONTHS All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 2ME treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

Southern Area—See Map

**COMMAND® 2ME HERBICIDE
APPLIED ALONE
RATES AND WEEDS CONTROLLED¹**

2.0 PT. ² (0.50 LB. A.I.)	3.0 PT. (0.75 LB. A.I.)	4.0 PT. (1.0 LB. A.I.)	5.0 PT. (1.25 LB. A.I.)
GRASSES Suppression of some annual grasses	GRASSES Barnyardgrass Broadleaf Signalgrass Crabgrass —Large —Smooth Foxtail —Giant —Green Goosegrass Panicum —Common —Fall —Texas Seedling Johnsongrass Field Sandbur*	GRASSES All grasses controlled at 3.0 pints plus: Cupgrass —South-western —Woolly Field Sandbur Bermudagrass* Red Rice Ichgrass ³	GRASSES All grasses controlled at 4.0 pints plus: Foxtail —Purple —Robust —Yellow Junglerice* Shattercane* Wild Proso Millet*
BROADLEAVES Velvetleaf Spurred Anoda	BROADLEAVES Velvetleaf Spurred Anoda plus: Common Ragweed* Jimsonweed* Lamb'squarter Pennsylvania Smartweed* Prickly Sida Purslane Redweed Venice Mallow	BROADLEAVES All broadleaves controlled at 3.0 pints plus: Cocklebur Common Ragweed Dayflower Florida Beggarweed Florida Pusley Jimsonweed Kochia* Redvine* Tropic Croton Wild Poinsettia ⁴	BROADLEAVES All broadleaves controlled at 4.0 pints plus: Balloonyvine* Black Nightshade* Curly Dock Joint Vetch* Kochia Morningglory —Pitted Pennsylvania Smartweed Pigweed —Redroot* —Smooth* Prostrate Spurge

* Partially Controlled

¹ Higher label rates (a minimum of 4 pints per acre) are recommended for fields that have previously been in conservation programs, or where heavy weed pressure is anticipated. Select lower to higher rates based on lighter to heavier soil types. For fields that have previously been in conservation programs, a minimum application rate of 5 pints per acre for cocklebur or common ragweed control is recommended.

² For heavy weed pressure or heavy soils use 2.5 pints per acre to control Velvetleaf or Spurred Anoda.

³ For control of Ichgrass, do not incorporate Command 2ME. For high weed pressure and heavy soils use 5 pints.

⁴ Use in ARL/AMS/TX/OK only—for high pressure and heavy soils use 5 pints.

NOTE: When using Command 2ME more than 30 days prior to planting, the highest label rate should be used.

ROTATIONAL CROP GUIDELINES

Rotate to Crops as listed below, otherwise crop injury may occur.

Refer to sections headed "Rotational Cropping Precautions" and "General Information."

NOTE: When using Command 1.5ME with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

2.0 PT. (0.50 LB. A.I.)	3.0 to 4.0 PT. (0.75 to 1.0 LB. A.I.)	5.0 PT. (1.25 LB. A.I.)
ANYTIME Soybeans Cotton* Peppers Pumpkins Peas	ANYTIME Soybeans Cotton* Peppers Pumpkins	ANYTIME Soybeans Cotton*
9 MONTHS Corn —Field —Pop —Seed —Sweet	9 MONTHS Same as 2.0 pint, plus: Peas	9 MONTHS Same as 3.0 to 4.0 pints, plus: Pumpkins
Cotton Cucurbits Dry Beans Peanuts Potatoes Pumpkins Rice Snap Beans Sorghum Sugar Beets Sweet Potatoes Tobacco Tomatoes (Transplanted)		

12 MONTHS	12 MONTHS	12 MONTHS
All crops	All crops	All crops

Cover crops, however, may be planted anytime but stand reductions may occur in some areas. Do not graze or harvest for food or feed cover crops planted less than 9 months after Command 2ME treatment. Do not allow livestock to graze on treated soybean vines or feed treated vines or vine trash to livestock.

* Refer to the "Requirements for Planting Time Applications" and "Replanting Instructions" in the Cotton Directions for Use section.

**COMMAND® 2ME HERBICIDE
APPLIED IN COMBINATIONS FOR SOYBEANS**

Command 2ME herbicide may be tank mixed with other soybean herbicides to broaden the weed control spectrum compared to the products applied alone. In the sections that follow FMC recommends a combination of products and application rates to broaden weed control spectrum. FMC makes no claims as to weed control for any combination of rates or products other than as recommended in this label. Additional recommendations for product use with other herbicides not included in the following sections may become available subsequent to the date of publication for this label. Please check with your dealer for more recent FMC technical bulletins and label supplements. When applying mixtures of Command 2ME with other herbicides, observe all precautions and limitations appearing on the labels of each product. Do not incorporate tank mixtures of Command 2ME and other products. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixes.

**Command 2ME Herbicide plus Canopy® Herbicide—
Northern & Southern Areas**

Tank mix combination for broadened weed control spectrum.

Soil Texture	Broadcast Rates Per Acre*		
	Command North	Command South	Canopy
(1/2** to 5% Organic Matter)			
Coarse (light) Soils: (loamy sand, sandy loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	4 to 6 oz.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	5 to 8 oz.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	4.0 to 5.0 pints	6 to 8 oz.

* Select lower to higher rates within the range to correspond to the coarser to finer soil types within the textural group.

Rate within the range should correspond to anticipated weed pressure.

** Do not apply to soils less than 1/2% Organic Matter.

16

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Cocklebur**
- Copperleaf, hophornbeam*
- Hemp sesbania
- Morning-glory* (Annual, Ivy-leaf, Entireleaf, Pitted, Tall)
- Pigweed (Redroot, Smooth)
- Sicklepod*

* Partially Controlled

** Large seeded weeds, germinating deep in the soil, such as morningglory, sicklepod and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a postemergence herbicide for season long control.

When applying mixtures of Command 2ME plus Canopy, observe all cautions and limitations appearing on the labels of each product. Special attention must be paid to sections regarding planting depth, phosphate pesticide interaction, previous pesticides, and sensitive soybean varieties on the Canopy label. Observe application precautions rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Dual® 8 E Herbicide—Northern & Southern Areas

Tank mix combination for broadened weed control spectrum.

Broadcast Rates Per Acre

Soil Texture	Command		Dual 8 E
	North	South	
Coarse (light) Soils: (loamy sand, sandy loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	1½ to 2 pints
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	2 to 2½ pints
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	4.0 to 5.0 pints	2 to 2½ pints

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Nutsedge
- Red Rice*
- Witchgrass

* Use higher rates in areas with heavy pressure.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Black Nightshade
- Carpetweed
- Pigweed

When applying mixtures of Command 2ME plus Dual, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Lasso® or Lasso® Micro-Tech—Northern & Southern Areas

A tank mix combination of Command 2ME herbicide and Lasso or Lasso Micro-Tech herbicide will broaden the weed control spectrum compared to either product used alone.

Broadcast Rates Per Acre

Soil Texture	Command		Lasso/Lasso Micro-Tech
	North	South	
Coarse (light) Soils: (loamy sand, sandy loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	2 to 2½ quarts
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	2½ to 3 quarts
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	4.0 to 5.0 pints	2½ to 3 quarts

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., sand, loamy sand, sandy loam) within a textural group.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Black Nightshade
- Carpetweed
- Pigweed

When applying mixtures of Command 2ME herbicide plus Lasso or Lasso Micro-Tech, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Lorox®

A preemergence tank mix combination of Command 1.5ME herbicide and Lorox herbicide will broaden the weed control spectrum compared to either product alone.

The tank mix combination may not be incorporated.

Broadcast Rates Per Acre*

Soil Texture**	Command		Lorox DF*
	North	South	
Coarse (light) Soils: (loamy sand**, sandy loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	¾ to 1 lb.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	1 to 1½ lbs.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	4.0 to 5.0 pints	1½ to 2 lbs.

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

** Plant seed at least 1½ inches deep or Lorox injury to soybeans may result. Refer to Lorox label for other planting and application precautions, one pint of Lorox L equals one pound of Lorox DF, use equivalent rates.

** Do not use on sand, loamy sand, gravelly soils, or exposed subsoils or soils with less than 1% organic matter. Use only on loamy sand with over 1% organic matter.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Carpetweed
- Chickweed
- Pigweed (Redroot, Smooth)
- Wild Mustard
- Wild Radish

When applying mixtures of Command plus Lorox observe all cautions and limitations appearing on the labels of each product. Observe application precautions, rotational crop/grazing and feeding restrictions and replanting instructions sections of this label when using tank mixtures.

Command 2ME Herbicide plus Lorox Plus®

A preemergence tank mix combination of Command 2ME herbicide and Lorox Plus herbicide will broaden the weed control spectrum compared to either product alone.

The tank mix combination may not be incorporated.

Broadcast Rates Per Acre*

Soil Texture**	Command		Lorox Plus
	North	South	
Coarse (light) Soils: (loamy sand**, sandy loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	9 to 12 oz.
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	10 to 14 oz.
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	4.0 to 5.0 pints	11 to 14 oz.

- * Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- ** Plant seed at least 1 1/2 inches deep or Lorox injury to soybeans may result. Refer to Lorox Plus label for other planting and application precautions.
- ** Do not use on sand, loamy sand, gravelly soils, or exposed subsoils or soils with less than 1% organic matter. Use only on loamy sand with over 1% organic matter.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Black Nightshade*
- Carpetweed
- Cocklebur**
- Giant Ragweed*
- Morning-glory* (Annual, Ivy-leaf, Entireleaf, Pitted, Tall)
- Mustards
- Pigweed (Redroot, Smooth)

* Partially Controlled

- ** Large seeded weeds, germinating deep in the soil, such as morningglory, sicklepod and cocklebur or weeds with subsequent flushes may require a cultivation or an application of a postemergence herbicide for season long control.

When applying mixtures of Command plus Lorox Plus observe all cautions and limitations appearing on the labels of each product. Observe application precautions, rotational crop/grazing and feeding restrictions and replanting instructions sections of this label when using tank mixtures.

Command 2ME Herbicide plus Metribuzin (Sencor® or Lexone®)—Northern Area Only

A tank mix combination of Command 1.5ME herbicide and metribuzin will broaden the weed control spectrum compared to either product used alone.

Broadcast Rates Per Acre*

Soil Texture**	Command		Sencor DF* Lexone DF
	Command	Sencor DF* Lexone DF	
Coarse (light) Soils: (loamy sand**, sandy loam)	3.0 to 4.0 pints	1/4 to 1/2 lb.	
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	1/4 to 1/2 lb.	
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	1/3 to 1/2 lb.	

- * Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- ** For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.
- ** Do not use Command 1.5ME in combination with metribuzin on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Carpetweed
- Pigweed* (Redroot, Smooth)
- Wild Mustard

- * Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 2ME plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME herbicide plus Metribuzin (Sencor® or Lexone®)—Southern Area Only

A tank mix combination of Command 1.5ME herbicide and metribuzin will broaden the weed control spectrum compared to either product used alone.

Broadcast Rates Per Acre*

Soil Texture**	Command		Sencor DF* Lexone DF
	Command	Sencor DF* Lexone DF	
Coarse (light) Soils: (loamy sand**, sandy loam)	4.0 to 5.0 pints	1/4 to 1/2 lb.	
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	4.0 to 5.0 pints	1/4 to 1/2 lb.	
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	4.0 to 5.0 pints	1/3 to 1/2 lb.	

- * Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.
- ** For pints Sencor 4 or Lexone 4L, multiply pounds DF by 1.5.
- ** Do not use Command 1.5ME in combination with metribuzin on soils with less than 0.5% organic matter, on sand, or on loamy sand with less than 2% organic matter, or on soils having a calcareous surface area or a pH of 7.5 or higher as metribuzin injury to soybeans may occur. Use 1/2 to 1 pound metribuzin on silty clay and heavy clay soils of the Mississippi Delta.

NOTE: Metribuzin injury to soybeans may occur under excess moisture conditions, cool weather, or compacted soil areas.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:

- Carpetweed
- Hemp sesbania (Coffee-weed)
- Pigweed (Redroot, Smooth)
- Sicklepod**
- Wild Mustard

- * Reduced Competition
- * Use the higher rate of metribuzin for a given soil type.

When applying mixtures of Command 2ME plus metribuzin, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Preview®—Northern Area Only

A tank mix combination of Command 2ME herbicide and Preview herbicide will broaden the weed control spectrum compared to either product used alone.

Broadcast Rates Per Acre*

Soil Texture	Command		Preview
	Command	Preview	
Coarse (light) Soils: (loamy sand, sandy loam)	3.0 to 4.0 pints	5 to 6 oz.	
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	6 to 7 oz.	
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	6 to 8 oz.	

- * Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

Do not use Command 2ME in combination with Preview on soils with less than 0.5% organic matter, or on soils with pH greater than 6.8 or on soils having a calcareous surface area as Preview injury to rotational crops may occur.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
 Burcucumber*
 Carpetweed
 Cocklebur*
 Copperleaf, hophornbeam
 Giant Ragweed*
 Morning-glory* (Annual, Tall, Ivyleaf, Pitted, Entireleaf)
 Mustards
 Pigweed
 Spurge, spotted
 Sunflower*

* NOTE: Control of morningglory, sunflower, cocklebur and giant ragweed may be erratic ranging from poor to excellent depending upon soil temperature, time of weed seed germination, depth of weed seed in the soil, and the amount and timing of soil moisture. Control may be improved with timely cultivation.

When applying mixtures of Command 2ME plus Preview, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Prowl® 4 EC Herbicide—Northern & Southern Areas

A tank-mix combination of Command 2ME and Prowl will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre*		
	Command		Prowl 4 EC
	North	South	
Coarse (light) Soils: (loamy sand, sandy loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	1 to 1½ pints
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	3.0 to 4.0 pints	4.0 to 5.0 pints	2 pints
Heavy (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	3.0 to 4.0 pints	4.0 to 5.0 pints	2 to 2½ pints

* Select lower to higher rates within the ranges noted based on lighter to heavier soil types (e.g., loam, silt loam, sandy clay, sandy clay loam) within a textural group.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
 Carpetweed
 Pigweed (Redroot, Smooth)

When applying mixtures of Command 2ME plus Prowl, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Scepter® Herbicide—Northern Area Only

See Scepter label for appropriate geographic and rainfall restrictions.

A tank mix combination of Command 1.5ME and Scepter will broaden the weed control spectrum of either product used alone.

NOTE: The Scepter label does not allow rotation to Corn, Wheat, Barley, or Oats prior to 18 months after last Scepter application when Scepter is applied preemergent within the Northern Use Area as defined on the Scepter label. See Scepter label for appropriate geographic restrictions.

Soil Texture	Broadcast Rates Per Acre*	
	Command	Scepter
All Soil Types	3.0 to 4.0 pints	½ to ½ pint

* Use higher use rates, within the range stated above, when heavier weed pressure is anticipated.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
 Black Nightshade
 Cocklebur
 Common Sunflower
 Giant Ragweed*
 Pigweed spp.

* Partially controlled

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT make additional Command 2ME or Command-Scepter® tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

When applying mixtures of Command 2ME plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Scepter Herbicide—Southern Area Only

See Scepter label for appropriate geographic and rainfall restrictions.

A tank mix combination of Command 2ME and Scepter will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre*	
	Command	Scepter
All Soil Types	4.0 to 5.0 pints	½ to ¾ pint

* Use higher use rates, within the range stated above, when heavier weed pressure is anticipated.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
 Black Nightshade
 Cocklebur
 Common Sunflower
 Pigweed spp.
 Wild Poinsettia

In the event of a crop loss due to weather conditions, soybeans can be replanted. DO NOT make additional Command 2ME or Command-Scepter tank mix applications.

NOTE: Adequate soil moisture is important for optimum tank mix applications.

When applying mixtures of Command 2ME plus Scepter, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

Command 2ME Herbicide plus Sonalan® EC Herbicide—Northern Area Only

A tank mix combination of Command ME and Sonalan will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre*	
	Command	Sonalan EC
Coarse (light) Soils: (loamy sand, sandy loam)	4.0 to 5.0 pints	1¼ to 2 pints
Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	4.0 to 5.0 pints	1¾ to 2½ pints
Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	4.0 to 5.0 pints	2¼ to 3 pints

* Select lower to higher rates within the range to correspond to the coarser to finer soil types within the textural group. Rates within the range should correspond to anticipated weed pressure.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
 Carpetweed
 Pigweed (Redroot, Smooth)

When applying mixtures of Command 2ME plus Sonalan, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 2ME Herbicide plus Squadron—
Northern Area Only**

See Squadron label for appropriate geographic restrictions.

A tank mix combination of Command 2ME and Squadron, will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Squadron*
All Soil Types	3.0 to 4.0 pints	1 1/2 to 2 1/4 pint

* Do not exceed 1 1/2 pints in Northern Area (as defined on the Squadron label). Use higher rate of Squadron in areas of heavy cocklebur infestation.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
 Black Nightshade
 Cocklebur
 Common Sunflower
 Giant Ragweed*
 Pigweed spp.

* Partially controlled

When applying mixtures of Command 2ME plus Squadron observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**Command 2ME Herbicide plus Squadron—
Southern Area Only**

See Squadron label for appropriate geographic restrictions.

A tank mix combination of Command 2ME and Squadron will broaden the weed control spectrum of either product used alone.

Soil Texture	Broadcast Rates Per Acre	
	Command	Squadron*
All Soil Types	4.0 to 5.0 pints	2 1/4 to 3 pints

* Use higher rate of Squadron in areas of heavy pigweed and cocklebur infestation.

WEEDS CONTROLLED:

Grass Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled.

Broadleaf Weeds

All species listed under the section Command Applied Alone Rates and Weeds Controlled, plus the following:
 Black Nightshade
 Cocklebur
 Common Sunflower
 Pigweed spp.
 Wild Poinsettia

When applying mixtures of Command 2ME plus Squadron, observe all cautions and limitations appearing on the labels of each product. Observe all application precautions, rotational guidelines and replanting instructions of each product's label when using tank mixtures.

**COMMAND 2ME HERBICIDE APPLICATION
FOLLOWED BY POSTEMERGENCE TREATMENTS**

Postemergence herbicides approved for use on soybeans may be applied following treatments of Command 2ME alone or in tank mix combinations for control of emerged weeds as listed on the postemergence product labels. Read and follow the precautionary statements, directions for use, rates of application and all other information appearing on the product labels.

DO NOT apply Classic, Pursuit or Scepter as a postemergence treatment after Command 2ME plus Canopy, Preview, Pursuit Plus, Scepter, or Squadron in a tank mix treatment, unless the other product labels specifically recommend such a treatment, as those other herbicides share a common mode of action and may cumulatively cause injury to soybeans or rotational crops.

**COMMAND 2ME HERBICIDE FOLLOWED BY
ACIFLUORFEN (BLAZER® 2L OR TACKLE®) HERBICIDE**

Following an application of Command 2ME or tank mixture with Command 2ME, a postemergence application of acifluorfen will control the following emerged broadleaf weeds:

Broadleaf Weeds

Black Nightshade
 Cocklebur
 Hemp sesbania
 Pigweed (Redroot)
 Pitted Morning-glory
 Purple Moonflower
 Wild Mustard

Apply acifluorfen at a rate of 1 pint per acre with .25% crop oil concentrate to actively growing weeds at no more than the 4-leaf growth stage (do not count cotyledonary leaves but only the fully developed true leaves). This timing generally correlates to soybean growth stages of the first to third trifoliate leaves.

NOTE: Application of Command 2ME herbicide generally retards the development rate of weeds which may extend the period where acifluorfen can be effectively applied.

Read and follow all precautions, restrictions, and warnings on all product labels.

**COMMAND 2ME HERBICIDE FOLLOWED BY
SCEPTER® HERBICIDE**

Following an application of Command 2ME or tank mixtures with Command 2ME, a postemergence application of Scepter herbicide will control the following emerged broadleaf weeds:

Broadleaf Weeds

Cocklebur, common
 Pigweed
 Wild Poinsettia

Apply Scepter after crop emergence but before weeds exceed a height of 12 inches (6 inches for Wild Poinsettia). Apply at a broadcast rate of 1/3 pint per acre. Apply when weeds are actively growing. DO NOT apply Scepter postemergence when soybeans and weeds have been subjected to stress conditions such as temperature and moisture extremes. The total amount of Scepter must not exceed 1/3 pint per acre per season in northern areas or 2/3 pint per acre per season in southern areas as defined on the Scepter label. Squadron includes the same active ingredient as Scepter.

For postemergence applications, the addition of a nonionic surfactant or crop oil concentrate is required. The nonionic surfactant should contain at least 80% active ingredient and should be applied at a rate of 2 pints per 100 gallons of spray mixture. Apply the crop oil concentrate (COC) at the rate stated on the COC label.

Read and follow all precautionary statements and restrictions, directions for use and all other information appearing on the Scepter label.

15 7 15

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

COMMAND and ~~PRIME~~—Trademarks of FMC Corporation
CANOPY, GLEAN, LEXONE, LOROX, LOROX PLUS and PREVIEW—
Trademarks of E.I. duPont de Nemours and Co., Inc.
OPTION—Trademark of Hoechst-Roussel Agri-Vet Company
SONALAN, TREFLAN—Trademarks of DowElanco
LASSO and ROUNDUP—Trademarks of Monsanto Company
SENCOR—Trademark of Bayer AG
PROWL, SCEPTER, SQUADRON, TRI-SCEPT—Trademarks of American Cyanamid Company
DUAL—Trademark of Ciba-Geigy Corporation
BLAZER—Trademark of BASF Corporation
TACKLE—Trademark of Rhone-Poulenc, Inc.
©1994 FMC Corporation

All rights reserved
(1584-1/14/94-A)

15